

are sub-micron intergrowths of allanite with a bastnäsite-type mineral phase.

### Acknowledgements

Electron microprobe facilities at Amsterdam were provided by the Free University (Amsterdam) and by WACOM, a working group for analytical geochemistry subsidized by The Netherlands Organization for the Advancement of Pure Research (NWO).

### References

- Baker, J. H., Hellingwerf, R. H., and Oen, I. S. (1988) *Geol. Mijnb.*, **67**, 121–38.
- Bayliss, P., Mazzi, F., Munno, R., and White, T. J. (1989) *Mineral. Mag.*, **53**, 565–9.
- Borodin, L. S., Bykova, A. B., Kapitonova, T. A., and Pyatenko, Yu. A. (1960) *Dokl. Acad. Sci. USSR, Earth Sci. Sect.*, **134**, 1022–4.
- Deer, W. A., Howie, R. A., and Zussman, J. (1986) *Rock-forming Minerals*, 2nd ed., Vol. 1B, 151–79. Longman Scientific & Technical.
- Fowler, M. B. and Williams, C. T. (1986) *Mineral. Mag.*, **50**, 326–8.
- Frondel, J. W. (1975) *Lunar Mineralogy*, 323. New York, Wiley Interscience.
- Gavelin, S. (1989) *Geol. Fören. Stockh. Förh.* **111**, 213–27.
- Geijer, P. (1927) *Sver. Geol. Unders.*, Ser. C, no. 343.
- (1961) *Arkiv. Mineral. Geol.*, **3**, 99–105.
- Gieré, R. (1986) *Contrib. Mineral. Petrol.*, **93**, 459–70.
- (1990) *Terra Nova*, **2**, 60–7.
- and Williams, C. T. (1991) Gac-MAC-SEG 16, Abstracts, May 1991, A44.
- Igelström, L. J. (1871) *Mineralogisk Vägvisare i Värmland*. Filipstad.
- Platt, R. G., Wall, F., Williams, C. T., and Woolley, A. R. (1987) *Mineral. Mag.*, **51**, 253–63.
- Purtscheller, F. and Tessardi, R. (1985) *Ibid.*, **49**, 523–9.
- Ringwood, A. E. (1985) *Ibid.*, **49**, 159–76.
- Rickard, D. T. (1988) *Geol. Mijub.*, **67**, 139–55.
- Tegengren, F. R. (1924) Sveriges ädlare malmer och bergwerk. *Sver. Geol. Unders.*, Ser. Ca, 17.
- Williams, C. T. and Gieré, R. (1988) *Schweiz. Mineral. Petr. Mitt.*, **68**, 133–40.

[Manuscript received 10 September 1989;  
revised 15 July 1991]

### Addendum

After the manuscript was accepted for publication, the description of a new mineral, dissakisite, the Mg-analogue of allanite was brought to our attention. (Grew, E. S., et al. 'Dissakisite-(Ce)', a new member of the epidote group and the Mg-analogue of allanite-(Ce) from Antarctica'. *Amer. Mineral.*, **76**, 1990–7.) It is apparent that the chemistry, optical properties and mineral paragenesis of allanite-(Ce) described here from Koberg is similar to that of the new mineral dissakisite-(Ce).