

The role of H₂O in Martian magmatic systems

BJORN O. MYSEN,^{1,*} DAVID VIRGO,¹ ROBERT K. POPP,² AND CONSTANCE M. BERTKA¹

¹Geophysical Laboratory, Carnegie Institution of Washington, 5251 Broad Branch Road, NW, Washington, D.C. 20015, U.S.A.

²Department of Geology and Geophysics, Texas A & M University, College Station, Texas 77843, U.S.A.

ABSTRACT

We have estimated the water content of Mars' interior by using analyzed water contents of kaersutite inclusions from shergottites nakhlites chassignites (SNC) meteorites in conjunction with an experimentally-derived crystal-chemical model of kaersutite amphibole. This model predicts quantitatively the relationships between iron oxidation and hydrogen deficiency in the kaersutite. The H₂O content of the magma from which the kaersutites in SNC meteorites could have crystallized is in the 100–1000 ppm range. That H₂O content leads to an estimated water content of 1–35 ppm for a Martian mantle that could have been the source rock for such magmas.

INTRODUCTION

Information on the H₂O abundance in the Martian interior is required to characterize the accretional and degassing history of the planet (McSween and Harvey 1993; Carr and Wänke 1992; Dreibus and Wänke 1985, 1987). The H₂O abundance also affects the composition of partial melts (Kushiro 1972; Stolper 1980; Treiman 1985; Longhi and Pan 1989; Johnson et al. 1991) and the physical properties of the magma [density, compressibility, and viscosity (Lange 1994; Schulze et al. 1996; Kushiro 1976)]. Those properties, in turn, govern the dynamics of magma ascent and the style of eruption.

Previous estimates of H₂O contents of the Martian interior vary widely, ranging from several tens of ppm (Dreibus and Wänke 1985) to perhaps 2000–3000 ppm (Reid and Bunch 1975; McSween and Harvey 1993; Harvey and McSween 1992). The lower estimates are based on solar element abundances and abundance ratios in shergottites nakhlites chassignites (SNC) meteorites and in carbonaceous chondrites. The SNC meteorites most likely are of Martian origin (Bogard and Johnson 1983; Becker and Pepin 1984; McSween 1985). The high estimates derive from the existence of hydrous minerals, such as kaersutitic amphibole, coexisting with glass as inclusions in olivine and pyroxene in SNC meteorites (Harvey and McSween 1992; Treiman 1985; Johnson et al. 1991) coupled with the assumption of H₂O saturation of the kaersutite [2(OH) pfu]. Kaersutite is a Ti-rich amphibole with the formula, Ca₂(Na,K)(Mg,Fe²⁺,Fe³⁺)₄(Ti,Al)(Si₆Al₂O₂₂(O,OH,F)₂ (Deer et al. 1963). In its structure there are two tetrahedral sites (T1 and T2) containing Si and Al, and five octahedral sites (two M1, two M2, and one M3 site) containing Fe³⁺, Fe²⁺, and Mg in the M1 and M3 sites and Ti and Al in the M2 site. There are three O sites (O1, O2, and O3)

with 2(OH,O,F) in the O3 site associated with the M3 octahedral site (Pechar et al. 1989).

Amphiboles that are water-saturated [that is 2(OH) pfu ~2 wt% H₂O in these kaersutites] most likely would crystallize from a magma that contains several weight percent H₂O. A Martian mantle source of such magma would probably contain several thousand ppm H₂O (Harvey and McSween 1992) or several times the concentration estimated for the interior of the Earth (Jambon and Zimmermann 1990).

The H₂O contents of kaersutite in Chassigny and Zagami SNC meteorites are, however, 0.1–0.2 wt% (Watson et al. 1994), or only 5–10% of the H₂O content of kaersutite with 2(OH) pfu (Table 1). These analyzed H₂O contents can be used to derive realistic H₂O contents of the magma with which the kaersutites were in equilibrium by using experimentally established quantitative relationships between H₂O content, redox ratio of Fe, and Ti content of kaersutite as a function of *T*, *P*, and *f*_{O₂}.

The calculation of H₂O content of the parental magma relies on the assumption that the amphiboles did not suffer H₂O-loss subsequent to crystallization. Despite suggestions that post-crystallization processes may affect the crystal chemistry of amphibole in terrestrial magmatic systems, the only study that incorporated data on hydrogen diffusion in minerals (Dyar et al. 1993) concluded that the OH content of kaersutite does not change significantly during transport from the Earth's mantle to the surface. In the Martian environment, the amphiboles may also have been shocked during the impact that led to their transport from Mars to Earth. Amphiboles shocked experimentally to ~30 GPa (Minitti and Rutherford 1998) yield ambiguous results as to whether such exposure may have affected their H content.

METHOD

The initial step in the procedure is to compute, from the experimental calibration data, the fugacity of H₂, *f*_{H₂},

* E-mail: mysen@gl.ciw.edu

TABLE 1. Composition of Vulcan's Throne kaersutite compared with kaersutite from SNC meteorites

	VT*	C†	Z‡
SiO ₂	40.32	39.49	35.15
TiO ₂	3.84	7.00	8.94
Al ₂ O ₃	15.35	14.22	15.13
FeO§	7.70	10.47	16.18
MgO	14.53	11.53	6.40
CaO	10.46	11.80	11.48
Na ₂ O	2.74	2.99	2.38
K ₂ O	1.60	0.33	0.20
H ₂ O	1.27	0.1–0.2	0.1–0.2
F	0.08		0.08

* VT = Vulcan's Throne, Arizona. Fe³⁺/(Fe³⁺ + Fe²⁺) = 0.317 (from Popp et al. 1995a and 1995b).

† C = Chassigny (from Johnson et al. 1991).

‡ Z = Zagami (from Treiman 1985).


§ Total iron as FeO.

|| H₂O concentration range from Watson et al. (1994).

during kaersutite crystallization. This information was obtained from the experimentally derived crystal-chemical model of kaersutite that predicts quantitatively the relationships between iron oxidation and hydrogen deficiency (Popp et al. 1995a, 1995b). This model relies on the premise that there are two exchange equilibria involving H, OH, and O²⁻ on the O3 site in kaersutite as follows:



and,




Whether these two substitution equilibria describe the relationships completely has been evaluated by examining the crystal-chemical relationships in natural and experimentally equilibrated amphiboles (Popp et al. 1995a, 1995b). If this assumption is correct, it follows from Equations 1 and 2 that F + Ti + Fe³⁺ + F = 2, which is exactly what has been observed (Popp et al. 1995a,

1995b). For reaction 1, which relates the redox ratio of Fe to the H₂O content of the amphibole, the equilibrium constant, K, is related to hydrogen fugacity, *f*_{H₂}, and activity product of relevant chemical species in amphibole, *a*, so that

$$\log K = \log f_{\text{H}_2} + \log a. \quad (3)$$

The simple relationship, log *a* = log (Fe³⁺/Fe²⁺) of the amphibole is inadequate (Popp et al. 1995b). Instead, they wrote an oxidation-dehydrogenation reaction of a pair of end-member amphiboles so that,



with the equilibrium constant,

$$K^- = f_{\text{H}_2} \frac{a_{\text{Fe}^{3+}}^{\text{prod}} a_{\text{Fe}^{2+}}^{\text{prod}} a_{[1]}^{\text{prod}}}{a_{\text{Fe}^{2+}}^{\text{react}} a_{\text{OH}}^{\text{react}}} \quad (5)$$

The *a*_[1]^{prod} denotes vacancies resulting from the activity of O3 sites from which H was lost during oxidation. Popp et al. (1995a, 1995b) found that the simplest mixing model, which assumed random mixing of Fe²⁺ and Fe³⁺ in reactant and product amphibole on the five octahedral sites, adequately reproduced the experimental data. Details of experimental data and data reduction can be found in Popp et al. (1995a, 1995b) (see Tables 1 and 2 in Popp et al. 1995a, and Tables 5, 6, and 7 and Fig. 5 in Popp et al. 1995b). The maximum difference between calculated and experimentally observed *f*_{H₂} with this model was found to be about 0.4 log units. K for Equations 3 and 5 was calibrated in the *f*_{H₂} range between magnetite-hematite and quartz-iron-wustite for *T* = 500°–1000 °C and *P* = 1–10 000 bar using a natural kaersutite from Vulcan's Throne, Arizona (Table 1).

The fugacity of H₂O, *f*_{H₂O}, is derived from the calculated *f*_{H₂} and dissociation constant of H₂O provided that the *f*_{O₂} is known. The *f*_{O₂} during equilibration of these SNC samples is reported to be near that of the quartz-fayalite-magnetite (QFM) buffer (Stolper and McSween

TABLE 2. Log *f*_{H₂} values calculated with 0.1 and 0.2 wt% H₂O in Chassigny and Zagami kaersutite

Pressure bar	0.1 wt% H ₂ O			0.2 wt% H ₂ O		
	800 °C	900 °C	1000 °C	800 °C	900 °C	1000 °C
Chassigny						
1	-4.356	-4.009	-3.717	-3.425	-3.078	-2.786
500	-4.301	-3.954	-3.662	-3.370	-3.023	-2.731
1000	-4.246	-3.899	-3.607	-3.315	-2.968	-2.676
2000	-4.136	-3.789	-3.497	-3.205	-2.858	-2.566
2500	-4.081	-3.734	-3.442	-3.150	-2.803	-2.511
4000	-3.916	-3.569	-3.277	-2.985	-2.638	-2.346
5000	-3.806	-3.459	-3.167	-2.875	-2.528	-2.236
6000	-3.696	-3.349	-3.057	-2.765	-2.418	-2.126
Zagami						
1	-3.422	-3.075	-2.783	-2.571	-2.225	-1.933
500	-3.367	-3.020	-2.728	-2.516	-2.170	-1.878
1000	-3.312	-2.965	-2.673	-2.461	-2.115	-1.823
2000	-3.202	-2.855	-2.563	-2.351	-2.005	-1.713
2500	-3.147	-2.800	-2.508	-2.296	-1.950	-1.658
4000	-2.982	-2.635	-2.343	-2.131	-1.785	-1.493
5000	-2.872	-2.525	-2.233	-2.021	-1.675	-1.383
6000	-2.762	-2.415	-2.123	-1.911	-1.565	-1.273

1979; McSween 1985; Johnson et al. 1991). The experimental relationship between f_{O_2} and T for the QFM buffer is from Chou (1987). Using f_{O_2} higher than that of the QFM buffer results in higher $f_{\text{H}_2\text{O}}$, whereas lower f_{O_2} results in lower $f_{\text{H}_2\text{O}}$ [from the H₂O dissociation, the $f_{\text{H}_2\text{O}}$ is proportional to $(f_{\text{O}_2})^{1/2}$].

Experimental studies of the relationships between water activity in silicate melts, $a_{\text{H}_2\text{O}}^m$, and the mole fraction, $X_{\text{H}_2\text{O}}^m$, at H₂O concentrations less than several tens of mole percent is consistent with the relationship (e.g., Burnham and Davis 1974; Burnham 1975; Hamilton et al. 1964; Holtz et al. 1995):

$$a_{\text{H}_2\text{O}}^m = k(X_{\text{H}_2\text{O}}^m)^2. \quad (6)$$

Values of k as a function of T and P for feldspar composition melts have been determined by Burnham (1975, 1994). Normative albite+orthoclase (albite > orthoclase) dominate the norm of the glass inclusions [We used the melt composition for Chassigny estimated by Johnson et al. (1991), so using Equation 6 with the calibration of the k values by Burnham (1975, 1994) is appropriate.] The activity of H₂O in the magma, $a_{\text{H}_2\text{O}}^m$, is simply the fugacity ratio, $f_{\text{H}_2\text{O}}/f_{\text{H}_2\text{O}}^0$. In order to use the experimentally calibrated k values (Eq. 6), the standard state used for $f_{\text{H}_2\text{O}}^0$ was that of Burnham (1975). The water content in ppm (wt) was calculated by using the compositions of glass inclusions coexisting with Chassigny kaersutite (Johnson et al. 1991).

RESULTS AND DISCUSSION

Values of f_{H_2} were calculated from the reported H₂O contents of the kaersutite inclusions (Watson et al. 1994) in the range $T = 800^\circ\text{--}1000^\circ\text{C}$ and $P = 1\text{--}6000$ bar (Table 2). Values of $f_{\text{H}_2\text{O}}$ and $a_{\text{H}_2\text{O}}$ were then computed from Equation 5 and the dissociation constant of H₂O at an f_{O_2} equivalent to the QFM univariant equilibrium. The H₂O concentration is derived with Equation 6.

The H₂O content of the magma in equilibrium with the Chassigny and Zagami kaersutites with 0.1 and 0.2 wt% H₂O is shown in Figure 1. The calculated water concentration in these melts varies with the H₂O content of the kaersutite so that an increase from 0.1 to 0.2 wt% regardless of T or P , results in a 250–300% increase in the water content of the melt (Fig. 1). A positive pressure dependence on H₂O concentration (~8% increase/kbar) is evident. This pressure effect reflects predominantly the influence of P on the k values in Equation 6. The temperature-dependence (15–20% increase/100 °C) reflects primarily the influence of T on the f_{H_2} (Table 2), as the k values in Equation 6 are relatively insensitive to T (Burnham 1975, 1994).

Although precise T and P estimates for crystallization of kaersutite in SNC inclusions are not available, crystallization temperatures near 1000 °C and pressure estimates in the 1–2 kbar range are often suggested (Stolper 1980; Treiman 1985; Johnson et al. 1991; Harvey and McSween 1992). The H₂O content of the melt inclusions at the time of kaersutite crystallization, therefore, was be-




FIGURE 1. H₂O contents of melts coexisting with Chassigny and Zagami kaersutite amphiboles as a function of T and P of kaersutite crystallization in the inclusions. The water contents were calculated with 0.1 and 0.2 wt% H₂O in the kaersutite as marked.

tween 100 and 300 ppm for Chassigny and between 300 and 1000 ppm H₂O for Zagami (Fig. 1). This water concentration range is similar to H₂O concentrations in heat-treated bulk SNC meteorites (Yang and Epstein 1985; Kerridge 1988; Karlsson et al. 1992). A kaersutite crystallization temperature lower than 1000 °C results in a lower H₂O content of the coexisting melt (Fig. 1) and, therefore, in the source region of the parental magma. Although the pressure assumed for the kaersutite crystallization is not critical to the estimate of water content, the H₂O contents for the most likely pressure range, 1–2 kbar, is highlighted in Figure 2.

From the H₂O contents in Figure 1, the H₂O of the Martian mantle that was melted to form the SNC parent magma can be estimated. Derivation of the H₂O content in this manner requires assessment of at least three unknowns: (1) How much crystallization occurred in the melt inclusion before kaersutite crystallized? (2) How much partial melting of the Martian interior occurred to produce the parental magma of the SNC meteorites? (3) The SNC meteorites show cumulate textures (McSween 1985). Therefore, did the melt inclusions get trapped in the cumulate olivine and pyroxene as the cumulate layers grew, or were these inclusions formed during early crystallization in the magma chamber?




FIGURE 2. H₂O of Martian mantle determined on the basis of the water contents in Figure 1 for 1000 °C as a function of pressure of kaersutite crystallization in the inclusions.

From mass-balance calculations and considerations of element partitioning and other phase-equilibrium data, the kaersutites equilibrated with a liquid residue representing 30–40% of the original melt volume (Longhi and Pan 1989; Harvey and McSween 1992). We will use 35% here, which yields H₂O contents in the SNC parent magma at 1000 °C and 1–2 kbar in the range 35–350 ppm prior to any crystallization (35–105 ppm for Chassigny and 105–350 ppm for Zagami).

Whether or not the inclusions were trapped prior to or during accumulation of crystals cannot be ascertained. It will be assumed, in accord with others (Harvey and McSween 1992), that inclusions were trapped prior to crystal accumulation. This implies that the parent magma of the SNC meteorites contained 35–350 ppm H₂O. If this assumption is incorrect, the H₂O content of the SNC parental magma would be lower than this estimate. In any event, even the highest calculated H₂O content is only about 50% of the lowest H₂O contents reported for primitive terrestrial mid-ocean ridge basalt (MORB) (Jambon and Zimmermann 1990) and could be as low as 5% of those values. These results clearly suggest that the source rock of the magma that gave rise to SNC meteorites ≤ 1.3 Ga (e.g., Shih et al. 1982; Jones 1989) could be considerably drier than the terrestrial mantle that melts to yield primitive MORB on Earth today.

The bulk-chemical composition suggested for the parental magma of the SNC meteorites is Al-poor tholeiitic basalt (McSween and Harvey 1993; Treiman 1985; Johnson et al. 1991). Results of melting experiments suggest that such magmas probably were derived from source regions previously depleted in an aluminous phase (Lon-

ghi and Pan 1989; Longhi 1991a). The approximate degree of melting of those source regions is subject to debate, with estimates ranging from <1 to 30–40% (Longhi and Pan 1989; Longhi 1991a, 1991b). The source regions undoubtedly had a complex history so those estimates are uncertain. Nevertheless, we have used 10% melting in our calculations. By assuming that H₂O behaves incompatibly so that all the H₂O in the Martian upper mantle entered the partial melt, the H₂O content of the magma source region is proportional to the degree of melting. The H₂O contents shown in Figure 2 result from 10% melting. For 1% melting, the source regions would contain 10% of the amounts indicated in Figure 2, and for 40% melting, the H₂O abundance in the source region would be four times that indicated in Figure 2. Longhi (1991b) suggested that the range in degree of melting suggested for the SNC parent magmas might reflect different petrogenetic histories. Chassigny might reflect $\sim 1\%$ melting, which would correspond to ~ 1 ppm H₂O in its mantle source. Zagami, on the other hand, might have been contaminated by a crustal component thus increasing its H₂O content (Longhi 1991b), which could explain the higher H₂O content suggested for the source region of its parental magma.

This H₂O content is equal to or less than that suggested on the basis of cosmochemical considerations (Dreibus and Wänke 1987) and is $\leq 5\%$ of that suggested under the assumption that the amphiboles in the SNC inclusions contained stoichiometric proportions of H₂O. In the latter case, 2(OH) pfu, there would be about 4 wt% H₂O in the coexisting magma and 2600 ppm H₂O in a source mantle of such magma. These numbers were obtained by using the same melt fractions as those used to derive Figure 2.

Even though the results of these calculations depend on the exact H₂O content of the kaersutite, it is evident that a Martian upper mantle with 1–35 ppm at the time of SNC parent magma generation is consistent with the existence of kaersutitic amphibole in the inclusions in SNC meteorites. For the first time there is agreement between estimates of H₂O of the Martian interior from cosmochemical considerations and from consideration of hydrous minerals in inclusions in SNC meteorites.

ACKNOWLEDGMENTS

This research was partially supported by PICS-CIW grant 142 and EAR-961443 to B.O.M. Detailed reviews by M. Rutherford, J. Longhi, and K. Lodders are appreciated.

REFERENCES CITED

- Becker, R.H. and Pepin, R.O. (1984) The case for a Martian origin of the shergottites: Nitrogen and noble gases in EETA 79001. *Earth and Planetary Science Letters*, 69, 225–242.
- Bogard, D.D. and Johnson, P. (1983) Martian gases in an Antarctic meteorite. *Science*, 221, 651–654.
- Burnham, C.W. (1975) Thermodynamics of melting in experimental silicate-volatile systems. *Geochimica et Cosmochimica Acta*, 39, 1077–1084.
- (1994) Development of the Burnham model for prediction of H₂O

- solubility in magmas. In *Mineralogical Society of America Reviews in Mineralogy*, 30, 123–130.
- Burnham, C.W. and Davis, N.F. (1974) The role of H₂O in silicate melts: II. Thermodynamic and phase relations in the system NaAl Si₃O₈-H₂O to 10 kilobars, 700 °C to 1100 °C. *American Journal of Science*, 274, 902–940.
- Carr, M.H. and Wänke, H. (1992) Earth and Mars: Water inventories as clues to accretional histories. *Icarus*, 98, 61–71.
- Chou, I.-M. (1987) Oxygen buffer and hydrogen sensor techniques at elevated pressures and temperatures. In G.C. Ulmer and H.L. Barnes, Eds., *Hydrothermal Experimental Techniques*, p. 61–99. Wiley, New York.
- Deer, W.A., Howie, R.A., and Zussman, M.A. (1963) *Rock-Forming Minerals*, 379 p. Longmans, Green and Co., Ltd, London.
- Dreibus, G. and Wänke, H. (1985) Mars: a volatile-rich planet. *Meteoritics*, 20, 367–382.
- (1987) Volatiles on Earth and Mars: A Comparison, 71 p. *Icarus*, 225–240.
- Dyar, M.D., Mackwell, S.J., McGuire, A.V., Cross, L.R., and Robertson, J.D. (1993) Crystal chemistry of Fe³⁺ and H⁺ in mantle kaersutite: Implications for mantle metasomatism. *American Mineralogist*, 78, 968–979.
- Hamilton, D.L., Burnham, C.W., and Osborn, E.F. (1964) The solubility of water and the effects of oxygen fugacity and water content on crystallization of mafic magmas. *Journal of Petrology*, 5, 21–39.
- Harvey, R.P. and McSween, H.Y. (1992) The parent magma of the nakhlite meteorites: Clues from melt inclusions. *Earth and Planetary Science Letters*, 111, 467–482.
- Holtz, F., Behrens, H., Dingwell, D.B., and Johannes, W. (1995) H₂O solubility in haplogranitic melts: Compositional, pressure, and temperature dependence. *American Mineralogist*, 80, 94–108.
- Jambon, A. and Zimmermann, J.L. (1990) Water in oceanic basalts: Evidence for dehydration of recycled crust. *Earth and Planetary Science Letters*, 101, 323–331.
- Jones, J.H. (1989) Isotopic relationships among the shergottites, the nakhlites, and Chassigny. *Proceedings of the 19th Lunar and Planetary Science Conference*, 465–474.
- Johnson, M.C., Rutherford, M.J., and Hess, P.C. (1991) Chassigny petrogenesis: Melt composition, intensive parameters, and water contents of Martian (?) magmas. *Geochimica et Cosmochimica Acta*, 55(1), 349–367.
- Karlsson, H.R., Clayton, R.N., Gibson, E.K., Mayeda, T.K., and Socki, R.A. (1992) Water in SNC meteorites: Evidence for a Martian hydrosphere. *Science*, 255, 1409–1411.
- Kerridge, J.F. (1988) Deuterium in Shergotty and Lafayette (and on Mars?). *Proceedings of the 18th Lunar and Planetary Science Conference*, 14, 599–600.
- Kushiro, I. (1972) Effect of water on the composition of magmas formed at high pressures. *Journal of Petrology*, 13, 311–334.
- (1976) Changes in viscosity and structure of melt of NaAlSi₃O₈ composition at high pressures. *Journal of Geophysical Research*, 81, 6347–6350.
- Lange, R.A. (1994) The effect of H₂O, CO₂, and F on the density and viscosity of silicate melts. In *Mineralogical Society of America Reviews in Mineralogy*, 30, 331–370.
- Longhi, J. (1991a) Comparative liquidus equilibria of hypersthene-normative basalts at low pressure. *American Mineralogist*, 76, 785–800.
- (1991b) Complex magmatic processes on Mars; inferences from the SNC meteorites. *Proceedings of the Lunar and Planetary Science Conference*, 21, 695–709.
- Longhi, J. and Pan, V. (1989) The parent magmas of SNC meteorites. *Proceedings of the 19th Lunar and Planetary Science Conference*, p. 451–464. Lunar & Planetary Institute, Houston TX.
- McSween, H.Y. (1985) SNC Meteorites: Clues to Martian Petrologic Evolution? *Reviews of Geophysics*, 23, 391–416.
- McSween, H.Y. and Harvey, R.P. (1993) Outgassed water on Mars: Constraints from melt inclusions in SNC meteorites. *Science*, 259, 1890–1892.
- Minitti, M.E. and Rutherford, M.J. (1998) Assessment of the shock effects on hornblende: Water contents and isotopic compositions. *Proceedings of the Lunar and Planetary Science Conference*, 29, 1435–1436.
- Pechar, F., Fuess, H., and Joswig, W. (1989) Refinement of the crystal structure of kaersutite (Vlcibra, Bohemia) from neutron diffraction. *Neues Jahrbuch Mineralogie Monatshefte*, 89, 137–143.
- Popp, R.K., Virgo, D., and Phillips, M.W. (1995a) H deficiency in kaersutitic amphiboles: Experimental verification. *American Mineralogist*, 80, 1347–1350.
- Popp, R.K., Virgo, D., Yoder, H.S., Hoering, T.C., and Phillips, M.C. (1995b) An experimental study of phase equilibria and Fe oxy-component in kaersutitic amphibole: Implications for the f_{H_2} and a_{H_2O} in the upper mantle. *American Mineralogist*, 80, 534–548.
- Reid, A.M. and Bunch, T.E. (1975) The nakhlites, II, Where, when, and how? *Meteoritics*, 10, 317–324.
- Schulze, F., Behrens, H., Holtz, F., Roux, J., and Johannes, W. (1996) The influence of H₂O on the viscosity of a haplogranitic melt. *American Mineralogist*, 81, 1155–1165.
- Shih, C.-Y., Nyquist, L.E., Bogard, D.D., McKay, G.G., Wooden, J.L., Bansal, B.M., and Wiesmann, H. (1982) Chronology and petrogenesis of young achondrites, Shergotty, Zagami, and ALHA 77005: Late magmatism on a geologically active planet. *Geochimica et Cosmochimica Acta*, 46, 2323–2344.
- Stolper, E. (1980) Predictions of mineral assemblages in planetary interiors. *Proceedings of the 11th Lunar and Planetary Science Conference*, 235–250.
- Stolper, E.M. and McSween, H.Y. (1979) Petrology and origin of the shergottite meteorites. *Geochimica et Cosmochimica Acta*, 43, 1475–1498.
- Treiman, A.H. (1985) Amphibole and hercynite spinel in Shergotty and Zagami: Magmatic water, depth of crystallization, and metasomatism. *Meteoritics*, 20, 229–243.
- Watson, L.L., Hutcheon, I.D., Epstein, S., and Stolper, E.M. (1994) Water on Mars: Clues from deuterium/hydrogen and water contents of hydrous phases in SNC meteorites. *Science*, 265, 86–90.
- Yang, J. and Epstein, S. (1985) A study of stable isotopes in Shergotty meteorite. *Proceedings of the 15th Lunar and Planetary Science Conference*, 12, 1473–1474.

MANUSCRIPT RECEIVED MARCH 2, 1998

MANUSCRIPT ACCEPTED MAY 4, 1998

PAPER HANDLED BY ROBERT F. DYMEK