

the original formula, although it should be pointed out that there is very little difference in composition for either formula.

The *x*-ray powder diffraction pattern of berryite from British Columbia is identical to that from the Nordmark Mines, Sweden and is, therefore, not given in this paper.

REFERENCES

- BORODAYEV, YU. S. & MOZGOVA, N.N. (1971). New group of the sulfbismuthides of Ag, Pb and Cu. *Soc. Mining Geol. Japan Spec. Issue* **2**, 35-41.
 KARUP-MØLLER, S. (1966). Berryite from Greenland, *Can. Mineral.* **8**, 414-423.
 NUFFIELD, E.W. & HARRIS, D.C. (1966). Studies of mineral sulphosalts: XX Berryite, a new species. *Can. Mineral.* **8**, 407-413.
 RUCKLIDGE, J. (1967). A computer program for processing microprobe data, *Jour. Geol.* **75**, 126.

Manuscript received January 27, 1972.

THE SILVER-ARSENIDE DEPOSITS OF THE COBALT-GOWGANDA REGION, ONTARIO

ERRATA : volume 11, part 1

- p. 72, line 27 : for *forms* read *form* ; for *as* read *an*
 p. 141, Table 13 : for C.O read CaO
 p. 300 : the equations were calculated with 0.08% MnO rather than 0.08% Mn
 p. 320, 4th line from bottom : for *Coldwell* read *Colwell*
 p. 347, caption to Fig. 217 : for *interested* read *intersected*
 p. 407, line 1 : for *Tables 91 and 101* read *Table 94 (p. 357) and 108 (p. 415)*

NOTICE TO AUTHORS

Beginning with Volume 12, Part 1, the Canadian Mineralogist will be published in a two-column format and will be issued four times per annum on the following schedule :

FEBRUARY — MAY — AUGUST — NOVEMBER

Authors submitting papers should consult the instructions published elsewhere in this issue. With the adoption of the new format, the issuance of 50 free separates to authors will cease. Orders for separates will, however, continue to be filled at printer's costs.