

FROM STRUCTURE TOPOLOGY TO CHEMICAL COMPOSITION. III. TITANIUM SILICATES: THE CRYSTAL CHEMISTRY OF BARYTOLAMPHYLLITE

ELENA SOKOLOVA[§] AND FERNANDO CÁMARA

CNR – Istituto di Geoscienze e Georisorse, unità di Pavia, Via Ferrata 1, I-27100 Pavia, Italy

ABSTRACT

The crystal structure of barytolamphyllite, ideally $(\text{BaK})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, a 19.8971(7), b 7.1165(3), c 5.4108(2) Å, β 96.676(2)°, V 760.96(3) Å³, space group $C2/m$, $Z = 2$, D_{calc} 3.521 g·cm⁻³, from Mt. Yuksorr, Khibina alkaline massif, Kola Peninsula, Russia, has been refined to $R_1 = 2.82\%$ on the basis of 1175 unique reflections ($F_o > 4\sigma F$) collected on a Bruker AXS SMART APEX diffractometer with a CCD detector and MoKα radiation. An electron-microprobe analysis gave (wt. %): SiO₂ 29.31, Al₂O₃ 0.16, Nb₂O₅ 0.11, TiO₂ 28.39, Fe₂O₃ 0.49, FeO 2.66, MnO 2.42, MgO 0.26, CaO 0.88, SrO 4.11, BaO 16.37, K₂O 3.52, Na₂O 7.90, F 1.69, H₂O 0.77 (calculated from structure refinement), O=F -0.71, for a total of 98.33. The empirical formula is $(\text{Ba}_{0.88}\text{K}_{0.61}\text{Sr}_{0.33}\text{Ca}_{0.05}\square_{0.13})\Sigma_{2.00}(\text{Na}_{2.09}\text{Fe}^{2+}_{0.30}\text{Mn}^{2+}_{0.28}\text{Ca}_{0.08}\text{Mg}_{0.05}\square_{0.20})\Sigma_{3.00}(\text{Ti}_{2.91}\text{Fe}^{3+}_{0.05}\text{Al}_{0.03}\text{Nb}_{0.01})\Sigma_{3.00}(\text{Si}_2\text{O}_7)_2\text{O}_2$ [F_{0.73}(OH)_{0.70}O_{0.57}] $\Sigma_{2.00}$, $Z = 2$, calculated on the basis of 4 Si *apfu*. The crystal structure of barytolamphyllite can be described as a combination of two blocks, a TS (titanium silicate) block and an I (intermediate) block. The TS block consists of HOH sheets (H heteropolyhedral, O octahedral), which occur in 24 Ti disilicate minerals. In the TS block, there are six cation sites, a [4]-coordinated Si site with $\langle \text{Si}-\text{O} \rangle = 1.631$ Å, two Ti-dominant sites, the [5]-coordinated M^H site and [6]-coordinated M^o(1) site, and two [6]-coordinated Na-dominant sites, M^o(2) and M^o(3). The M^H site (= Ti_{1.97}Al_{0.03} *pfa*) is coordinated by five O atoms, with $\langle \text{M}^{\text{H}}-\text{O} \rangle = 1.914$ Å; the M^o(1) site (= Ti_{0.94}Fe³⁺_{0.05}Nb_{0.01} *pfa*) is coordinated by four O atoms and two OH groups, with $\langle \text{M}^{\circ}(1)-\text{O} \rangle = 1.993$ Å; the M^o(2) and M^o(3) sites (= Na_{1.00} and Na_{1.09}Fe²⁺_{0.30}Mn²⁺_{0.28}Ca_{0.08}Mg_{0.05}□_{0.20} *pfa*, respectively) are coordinated by six O atoms with $\langle \text{M}^{\circ}(2)-\text{O} \rangle = 2.441$ Å, and four O atoms and two OH groups, with $\langle \text{M}^{\circ}(3)-\text{O} \rangle = 2.296$ Å, respectively. The M^H square pyramid and (Si₂O₇) groups constitute the H sheet. The M^o(1–3) octahedra form the close-packed O sheet. Linkage of H and O sheets occurs via common vertices of M^H square pyramids and (Si₂O₇) groups with M^o(1–3) octahedra. The second or intermediate I block includes the [10]-coordinated A^P site (= Ba_{0.88}K_{0.61}Sr_{0.33}Ca_{0.05}□_{0.13} *pfa*) with $\langle \text{A}^P-\text{O} \rangle = 2.843$ Å. The TS block and the I block (which ideally is a layer of Ba and K atoms) alternate along \mathbf{a} . The ideal formula of barytolamphyllite is $(\text{BaK})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, $Z = 2$. The topology of its structure is identical to that of lamphyllite, (SrNa)Na₃Ti₃(Si₂O₇)₂O₂(OH)₂, and nabalamphyllite, Ba Na₃Ti₃(Si₂O₇)₂O₂(OH)₂. We explain why, for lamphyllite, barytolamphyllite and nabalamphyllite, the chemical composition of the I block is of the form (A²⁺A⁺) with a total charge of 3+, and the ideal formula of the minerals is (A²⁺A⁺)Na₃Ti₃(Si₂O₇)₂O₂(OH)₂, where A²⁺ = Sr²⁺, Ba²⁺, Ca²⁺; A⁺ = Na⁺, K⁺.

Keywords: barytolamphyllite, Ti disilicate minerals, structure refinement, electron-microprobe analysis, chemical formula, TS block, lamphyllite, nabalamphyllite.

SOMMAIRE

Nous avons affiné la structure cristalline de la barytolamphyllite, de formule idéale $(\text{BaK})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, a 19.8971(7), b 7.1165(3), c 5.4108(2) Å, β 96.676(2)°, V 760.96(3) Å³, groupe spatial $C2/m$, $Z = 2$, D_{calc} 3.521 g·cm⁻³, provenant du mont Yuksorr, complexe alcalin de Khibina, péninsule de Kola, en Russie, jusqu'à un résidu R_1 de 2.82% en utilisant 1175 réflexions uniques ($F_o > 4\sigma F$) prélevées avec un diffractomètre Bruker AXS SMART APEX muni d'un détecteur CCD et avec un rayonnement MoKα. Une analyse avec une microonde électrique a donné (en %, poids): SiO₂ 29.31, Al₂O₃ 0.16, Nb₂O₅ 0.11, TiO₂ 28.39, Fe₂O₃ 0.49, FeO 2.66, MnO 2.42, MgO 0.26, CaO 0.88, SrO 4.11, BaO 16.37, K₂O 3.52, Na₂O 7.90, F 1.69, H₂O 0.77 (calculé à partir de l'affinement de la structure), O=F -0.71, pour un total de 98.33. La formule empirique est $(\text{Ba}_{0.88}\text{K}_{0.61}\text{Sr}_{0.33}\text{Ca}_{0.05}\square_{0.13})\Sigma_{2.00}(\text{Na}_{2.09}\text{Fe}^{2+}_{0.30}\text{Mn}^{2+}_{0.28}\text{Ca}_{0.08}\text{Mg}_{0.05}\square_{0.20})\Sigma_{3.00}(\text{Ti}_{2.91}\text{Fe}^{3+}_{0.05}\text{Al}_{0.03}\text{Nb}_{0.01})\Sigma_{3.00}(\text{Si}_2\text{O}_7)_2\text{O}_2$ [F_{0.73}(OH)_{0.70}O_{0.57}] $\Sigma_{2.00}$, $Z = 2$, calculé sur une base de 4 Si *apfu*. On peut décrire la structure de la barytolamphyllite comme une combinaison de deux blocs, un bloc TS (silicate de titane) et un bloc I (intermédiaire). Le bloc TS contient des feuillets HOH (H hétéropolyèdres, O octaèdres), qui se retrouvent aussi dans 24 minéraux disilicatés de Ti. Dans le bloc TS, il y a six sites pour cations, un site Si à coordinence [4], avec $\langle \text{Si}-\text{O} \rangle = 1.631$ Å, deux sites où dominent le Ti, un site M^H à coordinence [5], le site M^o(1), à coordinence [6], et deux sites où dominent Na à coordinence [6], M^o(2) et M^o(3). Le site M^H (= Ti_{1.97}Al_{0.03} *pfa*) est lié

[§] Permanent address: Department of Geological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada.
 E-mail address: elena_sokolova@umanitoba.ca

à cinq atomes d'oxygène, avec $\langle M^{\text{H}}-\text{O} \rangle = 1.914 \text{ \AA}$; le site $M^{\circ}(1)$ ($= \text{Ti}_{0.94}\text{Fe}^{3+}_{0.05}\text{Nb}_{0.01}\text{pfu}$) est lié à quatre atomes d'oxygène et deux groupes OH, avec $\langle M^{\circ}(1)-\text{O} \rangle = 1.993 \text{ \AA}$; les sites $M^{\circ}(2)$ et $M^{\circ}(3)$ ($= \text{Na}_{1.00}\text{Na}_{1.09}\text{Fe}^{2+}_{0.30}\text{Mn}^{2+}_{0.28}\text{Ca}_{0.08}\text{Mg}_{0.05}\square_{0.20}\text{pfu}$, respectivement) sont liés à six atomes d'oxygène avec $\langle M^{\circ}(2)-\text{O} \rangle = 2.441 \text{ \AA}$, et quatre atomes d'oxygène et deux groupes OH, avec $\langle M^{\circ}(3)-\text{O} \rangle = 2.296 \text{ \AA}$, respectivement. Une pyramide carrée M^{H} et les groupes (Si_2O_7) constituent le feuillett H. Les octaèdres $M^{\circ}(1-3)$ forment un feuillett O à empilement compact. Les agencements des feuillets H et O sont assurés par les coins communs des pyramides carrées M^{H} et des groupes (Si_2O_7) avec les octaèdres $M^{\circ}(1-3)$. Le second bloc, I ou intermédiaire, inclut le site A^P à coordinence [10] ($= \text{Ba}_{0.88}\text{K}_{0.61}\text{Sr}_{0.33}\text{Ca}_{0.05}\square_{0.13}\text{pfu}$), avec $\langle A^P-\text{O} \rangle = 2.843 \text{ \AA}$. Le bloc TS et le bloc I (qui est idéalement un niveau d'atomes de Ba et K) alternent le long de **a**. La formule idéale de la barytolamprophyllite est $(\text{BaK})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, $Z=2$. La topologie de sa structure est identique à celle de la lamprophyllite, $(\text{SrNa})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, et celle de la nabalamprophyllite, $\text{Ba Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$. Nous expliquons pourquoi, dans le cas de la lamprophyllite, barytolamprophyllite et nabalamprophyllite, la composition chimique du bloc I prend la forme $(\text{A}^{2+}\text{A}^+)$ avec une charge totale de 3+, et la formule idéale de ces minéraux est $(\text{A}^{2+}\text{A}^+)\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, avec $\text{A}^{2+} = \text{Sr}^{2+}, \text{Ba}^{2+}, \text{Ca}^{2+}; \text{A}^+ = \text{Na}^+, \text{K}^+$.

(Traduit par la Rédaction)

Mots-clés: barytolamprophyllite, minéraux disilicatés de Ti, affinement de la structure, analyses à la microsonde électronique, formule chimique, bloc TS, lamprophyllite, nabalamprophyllite.

INTRODUCTION

Currently, the crystal structures of about 160 Ti minerals are known. Half of them are Ti silicates and half of them are Ti oxides. Except for titanite, Ti silicates usually occur as accessory minerals. Nevertheless, Ti silicates are fairly numerous, particularly compared to analogous sulfates, phosphates and arsenates: there are no Ti sulfate minerals, one Ti phosphate mineral, and one Ti arsenate mineral, although several Ti silicate minerals also contain (PO_4) and (SO_4) groups. Sokolova (2006) considered structural hierarchy and stereochemistry for twenty-four titanium disilicate minerals containing the TS block. In the Ti disilicate minerals, the TS block consists of HOH sheets, where H is a heteropolyhedral sheet including (Si_2O_7) groups, and O is a trioctahedral close-packed sheet. Sokolova (2006) established the relation between structure topology and chemical composition for these minerals and divided them into four groups, characterized by different topology and stereochemistry of the TS block. Each group of structures has a different linkage of H and O sheets in the TS block and a different arrangement of Ti (= Ti + Nb) polyhedra. In a structure, the TS block can alternate with another block, called an intermediate (I) block, as it is intercalated between two TS blocks. In Groups I, II III and IV, Ti = 1, 2, 3 and 4 apfu, respectively. Six minerals belong to Group III. Three of these minerals have the same topology: lamprophyllite, $(\text{SrNa})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$ [which has two polytypes, one with orthorhombic symmetry and the other with monoclinic symmetry, designated as lamprophyllite-2O and lamprophyllite-2M by Johnsen (1996)], barytolamprophyllite, $(\text{BaK})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, and nabalamprophyllite, $\text{Ba Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$. Three minerals are of different topology: innelite, $\text{Na}_2\text{Ca Ba}_4\text{Ti}_3(\text{Si}_2\text{O}_7)_2(\text{SO}_4)_2\text{O}_4$ (Chernov *et al.* 1971), epistolite, $\text{Na}_4\text{Ti Nb}_2(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2(\text{H}_2\text{O})_4$ (Sokolova & Hawthorne 2004), and

vuonnemite, $\text{Na}_{11}\text{Ti Nb}_2(\text{Si}_2\text{O}_7)_2(\text{PO}_4)_2\text{O}_3\text{F}$ (Ercit *et al.* 1998). These ideal formulae for the six minerals are written as presented by Sokolova (2006).

There are ambiguities with regard to the ideal formulae of lamprophyllite and barytolamprophyllite that we will address in this paper. There is also need to examine the crystal structure of barytolamprophyllite characterized by a complete chemical analysis. In this paper, the third in this series (Sokolova 2006, Sokolova & Cámara 2007), we present the crystal chemistry of barytolamprophyllite and explain why, for lamprophyllite, barytolamprophyllite and nabalamprophyllite, the chemical compositions of the I block must be presented in the form $(\text{A}^{2+}\text{A}^+)$ with a total charge of 3+, and the ideal formulae, as $(\text{A}^{2+}\text{A}^+)\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$, where $\text{A}^{2+} = \text{Sr}^{2+}, \text{Ba}^{2+}, \text{Ca}^{2+}; \text{A}^+ = \text{Na}^+, \text{K}^+$. In response to a comment by one of the reviewers as to why did we do this work when the minerals have been so well characterized, we offer the following critical analysis.

PREVIOUS WORK

The first descriptions of lamprophyllite, barytolamprophyllite and nabalamprophyllite were published by Ramsay & Hackman (1894), Dudkin (1959) and Chukanov *et al.* (2004), respectively. The crystal structure of lamprophyllite was solved by Woodrow (1964), who gave the formula as $(\text{Ba,Ca,K})\text{Na}(\text{Ti,Fe})\text{Ti Si}_2(\text{O,OH,F})_9$, and the latest refinement of the structure is by Krivovichev *et al.* (2003), who gave the structural formula as $(\text{Sr}_{1.18}\text{Na}_{0.65}\text{Ca}_{0.12})\Sigma_{1.96}\text{Na}(\text{Na}_{1.3}\text{Mn}_{0.36}\text{Fe}_{0.22}\text{Mg}_{0.12})\text{Ti}_3\text{O}_2(\text{Si}_2\text{O}_7)_2(\text{OH})_2$. Rastsvetaeva & Chukanov (1999) reported the crystal structure of nabalamprophyllite (Table 1), and Chukanov *et al.* (2004) gave its ideal formula as $\text{Ba}(\text{Na,Ba})\{\text{Na}_3\text{Ti}[\text{T}_2\text{O}_2\text{Si}_4\text{O}_{14}](\text{OH,F})\}$. The crystal structure of barytolamprophyllite was solved by Peng *et al.* (1984), who gave the *crystallochemical* formula $(\text{Ba,Sr,K})_2\text{NaTi}(\text{Na,Fe}^{2+})_2(\text{Ti,Fe}^{3+})_2[\text{Si}_2\text{O}_7]_2\text{O}_2(\text{O,OH,F})_2$, $Z=$

2. Two later refinements of the structure of minerals given the unapproved names "Ba-lamprophyllite" (Rastsvetaeva & Dorfman 1995) and "K-barytolamprophyllite" (Rastsvetaeva *et al.* 1995) confirmed the lamprophyllite-type structure, space group $C2/m$, $Z = 2$, and gave following structural formulae: $(\text{Sr}_{1.1}\text{Ba}_{0.9})\{\text{Na}(\text{Na}_{0.9}\text{Mn}^{2+})_2(\text{Ti}_{0.7}\text{Mg}_{0.3}(\text{OH})_2)\}\{\text{[Si}_2\text{O}_7]_2 \bullet (\text{Ti}_{0.7}\text{Fe}^{3+})_2\text{O}_2\}$ for "Ba-lamprophyllite" and $\text{KBa}\{\text{Na}[\text{Na}_{1.2}(\text{Fe}^{2+},\text{Mn}^{2+})_{0.8}] \bullet [\text{Ti}(\text{O},\text{OH})_2]\}\{\text{[Si}_2\text{O}_7]_2\text{Ti}_2\text{O}_2\}$ for "K-barytolamprophyllite" (Table 1). We conclude that Rastsvetaeva & Dorfman (1995) and Rastsvetaeva *et al.* (1995) refined the crystal structures of lamprophyllite ($\text{Sr} > \text{Ba}$) and barytolamprophyllite ($\text{Ba} > \text{Sr}$), respectively, according to data presented by them. None of the three papers, Peng *et al.* (1984), Rastsvetaeva & Dorfman (1995), and Rastsvetaeva *et al.* (1995), contains results of a chemical analysis; empirical formulae were reported by Peng *et al.* (1984) and Rastsvetaeva *et al.* (1995): $(\text{Ba}_{1.05}\text{Sr}_{0.12}\text{K}_{0.53}\text{Ca}_{0.25}\text{Na}_{0.10})\Sigma_{2.05}\text{Na Ti}(\text{Na}_{1.37}\text{Fe}^{2+}_{0.29}\text{Mn}_{0.20}\text{Mg}_{0.15})\Sigma_{2.01}(\text{Ti}_{1.68}\text{Fe}^{3+}_{0.28}\text{Mg}_{0.05})\Sigma_{2.01}[(\text{Si}_{3.83}\text{Al}_{0.18})_{2.41}\text{O}_{14}] \text{O}_2(\text{O}_{0.57}\text{OH}_{0.62}\text{F}_{0.68}\text{Cl}_{0.14})\Sigma_{2.01}$ and $(\text{K}_{0.98}\text{Ba}_{0.84}\text{Sr}_{0.21})\Sigma_{2.03}[\text{Na}(\text{Na}_{1.2}\text{Fe}_{0.41}\text{Mn}_{0.42}\text{Ca}_{0.27})_{2.3}(\text{TiO}_2)_{1.26}] \bullet (\text{Si}_2\text{O}_7)_2\text{Ti}_2\text{O}_2$, respectively.

Mandarino & Back (2004) gave the following formulae: lamprophyllite, $(\text{Na,Mn,Ca,Fe})_3(\text{Sr,Ba,K})_2(\text{Ti,Fe})_3\text{O}_2(\text{Si}_2\text{O}_7)_2(\text{O,OH,F})_2$; barytolamprophyllite, $(\text{Na,K})_2(\text{Ba,Ca,Sr})_2(\text{Ti,Fe})_3(\text{SiO}_4)_4(\text{O,OH})_2$. The formula for lamprophyllite has 12 cations, whereas the formula for barytolamprophyllite has 11 cations *p.f.u.* If they are isostructural, this is obviously not correct (as the presence of vacancies should also be recognized in a structural formula). Yakovenchuk *et al.* (2005) wrote the formulae of lamprophyllite and barytolamprophyllite as $(\text{Sr,Ba,Na,Ca})_2\text{Na}(\text{Na,Mn,Fe})_2\text{Ti}_3\text{O}_2[\text{Si}_2\text{O}_7]_2(\text{OH})_2$ and $(\text{Ba,Sr,Na,Ca})_2\text{Na}(\text{Na,Mn,Fe})_2\text{Ti}_3\text{O}_2[\text{Si}_2\text{O}_7]_2(\text{OH})_2$, respectively. Krivovichev *et al.* (2003) and Chukanov *et al.* (2004) stated that lamprophyllite and barytolamprophyllite have formulae $(\text{Sr,Ba})_2\text{Na}_3$

$\text{Ti}_3\text{O}_2(\text{Si}_2\text{O}_7)_2(\text{OH})_2$ and $(\text{Ba,Sr})_2\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$; these formulae are not neutral, as they have an excess charge of +1 [note that Krivovichev *et al.* (2003) wrongly referred to the "Ba-lamprophyllite" of Rastsvetaeva & Dorfman (1995) as barytolamprophyllite- $2M$]. Sokolova (2006) considered the structure work of Rastsvetaeva *et al.* (1995) as the latest structure work on barytolamprophyllite, $(\text{KBa})\text{Na}_3\text{Ti}_3(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$; the latter ideal formula is neutral.

To summarize, (1) the crystal structure of barytolamprophyllite has never been done on a sample for which a chemical composition was given, and hence we do not know if the structure has ever been examined; (2) the chemical formulae of lamprophyllite and barytolamprophyllite quoted in Krivovichev *et al.* (2003) and Chukanov *et al.* (2004), *i.e.*, the latest works on the structure, are not neutral; they have an excess charge of +1, which is obviously wrong; (3) points (1) and (2) make it clear that there is an ambiguous situation with regard to the chemical formulae of the lamprophyllite group of minerals. We resolve this issue here.

EXPERIMENTAL

The crystals of barytolamprophyllite from Mt. Yuksporr, Khibina alkaline massif, Kola Peninsula, Russia, were kindly provided by Adriana and Renato Pagano, Milan, Italy, from their mineral collection (Collezione Mineralogica, sample #6270). These crystals were removed from a sample of delindeite originally obtained from A.P. Khomyakov and described in Khomyakov (1995).

Electron-microprobe analysis

One fragment of the crystal of barytolamprophyllite previously used for the structure refinement was analyzed with a Cameca SX-100 electron microprobe operating in wavelength-dispersion mode with an

TABLE 1. STRUCTURAL FORMULAE* AND UNIT-CELL PARAMETERS FOR SELECTED GROUP-III MINERALS WITH THE TS BLOCK

Mineral	Formula	f block	a (Å)	b (Å)	c (Å)	β (°)	Sp. gr.	Z	Ref.
lamprophyllite- $2M$	$(\text{SrNa})\text{Ti}_2\text{Na}_3\text{Ti}(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$	$(\text{Sr}_{1.18}\text{Na}_{0.65}\text{Ca}_{0.12})$	19.215	7.061	5.3719	96.797	$C2/m$	2	(1)
lamprophyllite- $2O$	$(\text{SrNa})\text{Ti}_2\text{Na}_3\text{Ti}(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$	$(\text{Sr}_{1.16}\text{Na}_{0.65}\text{Ca}_{0.12})$	19.128	7.0799	5.3824		$Pnmm$	2	(1)
nabalamprophyllite	$\text{Ba Na Ti}_2\text{Na}_3\text{Ti}(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$	$(\text{Ba}_{1.30}\text{Na}_{0.50}\text{K}_{0.15}\text{Sr}_{0.05})^{**}$	19.741	7.105	5.408	96.67	$P2/m$	2	(2)
barytolamprophyllite	$(\text{BaK})\text{Ti}_2\text{Na}_3\text{Ti}(\text{Si}_2\text{O}_7)_2\text{O}_2(\text{OH})_2$	$(\text{Ba}_{0.66}\text{K}_{0.61}\text{Sr}_{0.33}\text{Ca}_{0.05})\square_{0.13}$	19.8971	7.1165	5.4108	96.676	$C2/m$	2	(3)

* For each mineral, the structural formula is written in the form $A^P_2\text{M}^H_2\text{M}^O_4(\text{Si}_2\text{O}_7)_4\text{X}^O_4$ following Sokolova (2006).

** This composition is from the description of the new mineral species (Chukanov *et al.* 2004) rather than the structure work of Rastsvetaeva & Chukanov (1999). Nabalamphrophyllite is the only mineral with a lamprophyllite-type structure where there are two independent Ba- and Na-dominant A^P sites.

References (the latest reference on the structure): (1) Krivovichev *et al.* (2003), (2) Rastsvetaeva & Chukanov (1999), (3) this work. The data for "Ba-lamprophyllite" (Rastsvetaeva & Dorfman 1995), a 19.744, b 7.099, c 5.409 Å, β 96.77°, space group $C2/m$ and "K-barytolamprophyllite" (Rastsvetaeva *et al.* 1995), a 20.086, b 7.099, c 5.411 Å, β 96.65°, space group $C2/m$, are not included in the Table as there are no chemical data given in these papers. Sp. gr.: space group.

accelerating voltage of 15 kV, a specimen current of 20 nA, a beam size of 10 μm , and count times on peak and background of 20 and 30 s for major and minor elements (<1 wt.%), respectively. The following standards and crystals were used for *K* or *L* X-ray lines: F: riebeckite; Na: jadeite; Ba: barite; Si, Ca: diopside; Fe: fayalite; Mn: spessartine; K: orthoclase; Ti: titanite; Zn: zircon; Nb: BaNaNbO₄; Mg: forsterite; Sr: strontianite; Al: andalusite. Data were reduced using the $\phi(\rho Z)$ procedure of Pouchou & Pichoir (1985). The chemical composition of barytolamprophyllite is given in Table 2 and is the mean result of 10 point analyses.

The empirical formula is (Ba_{0.88} K_{0.61} Sr_{0.33} Ca_{0.05} □_{0.13})Σ_{2.00} (Na_{2.09} Fe²⁺_{0.30} Mn²⁺_{0.28} Ca_{0.08} Mg_{0.05} □_{0.20})Σ_{3.00} (Ti_{2.91} Fe³⁺_{0.05} Al_{0.03} Nb_{0.01})Σ_{3.00} (Si₂O₇)₂ O₂ [F_{0.73} (OH)_{0.70} O_{0.57}]Σ_{2.00}. *Z* = 2, calculated on the basis of 4 Si *apfu*; the amount of (H₂O) was calculated from the structure refinement, the ideal formula is (BaK) Na₃Ti₃ (Si₂O₇)₂ O₂ (OH)₂, in accord with Sokolova (2006).

Data collection and crystal-structure refinement

A single crystal of barytolamprophyllite was mounted on a Bruker AXS SMART APEX diffractometer equipped with graphite-filtered MoK α X-radiation and a CCD detector. The intensities of 8747 reflections with $-27 < h < 27$, $-9 < k < 9$, $-7 < l < 7$ were collected to 59.86° 20 using 20 s per 0.2° frame, and an empirical absorption-correction (SADABS, Sheldrick 1998) was applied. The refined unit-cell parameters (Table 3) were obtained from 5802 independent reflections. On the basis of 1185 unique observed reflections ($F_o > 4\sigma F$), the crystal structure of barytolamprophyllite was refined in the space group *C*2/m with the Bruker SHELXTL Version 5.1 system of programs (Sheldrick 1997) to R_1 = 2.8% and a GoF of 1.12. Scattering curves for neutral atoms were taken from the International Tables for Crystallography (1992). The *R* indices are

given in Table 3, and are expressed as percentages. Site occupancies for the M^{H} , $M^{\circ}(1)$; $M^{\circ}(2)$, $M^{\circ}(3)$ and A^P sites were refined with the scattering curves of Ti; Na, Fe and Sr, respectively.

Details of the data collection and structure refinement are given in Table 3, final atom-parameters are given in Table 4, selected interatomic distances and angles in Table 5, refined site-scattering values and assigned populations for selected sites in Table 6 and bond-valence values in Table 7. A table of structure factors may be obtained from the Depository of Unpublished Data, MAC web site [document Barytolamprophyllite CM46_403].

DESCRIPTION OF THE STRUCTURE

Cation sites

H sheet: In the crystal structure of barytolamprophyllite, there is one tetrahedrally coordinated site occupied by Si, with a $\langle \text{Si}-\text{O} \rangle$ distance of 1.631 Å. There is one [5]-coordinated Ti-dominant site, M^{H} , coordinated by five O atoms arranged in a square pyramid, with a $\langle M^{\text{H}}-\text{O} \rangle$ distance of 1.914 Å (Table 5); this site is occupied by Ti_{1.97}Al_{0.03} *pfu* (Table 6).

O sheet: The $M^{\circ}(1)$ site, Ti_{0.94}Fe³⁺_{0.05}Nb_{0.01}, is coordinated by four O atoms and two anions (OH,F) (1), with a $\langle M-\text{O},\text{OH} \rangle$ distance of 1.993 Å. There are two [6]-coordinated sites, primarily occupied by alkali cations. The $M^{\circ}(2)$ site is coordinated by O atoms and is fully occupied by Na, with a $\langle M^{\circ}(2)-\text{O} \rangle$ distance of 2.441 Å. The $M^{\circ}(3)$ site, Na_{1.09}Fe²⁺_{0.30}Mn²⁺_{0.28}Ca_{0.08} Mg_{0.05} □_{0.20} *pfu*, is coordinated by four O atoms and two monovalent anions [which occur at the O(1) site; see discussion below], with a $\langle M^{\circ}(3)-\text{O} \rangle$ distance of 2.296 Å.

I block: There is one A^P site, which is occupied by Ba_{0.88}K_{0.61}Sr_{0.33}Ca_{0.05} □_{0.13} *pfu*. There are ten $A^P-\text{O}$ distances less than 3 Å (Table 5), and one longer distance, $A^P-(\text{O})(1) = 3.225$ Å. We consider the A^P site as [10]-coordinated, with a $\langle [10]A^P-\text{O} \rangle$ distance of 2.843 Å.

TABLE 2. CHEMICAL COMPOSITION (wt%) AND UNIT FORMULA (*apfu*) OF BARYTOLAMPROPHYLLITE

SiO ₄	29.31	Si	4.00	Na	2.09
Nb ₂ O ₅	0.11	Σ	4.00	Fe ²⁺	0.30
TiO ₂	28.39			Mn ²⁺	0.28
Al ₂ O ₃	0.16			Ca	0.08
Fe ₂ O ₃	0.49	Ti ⁴⁺	1.97	Mg	0.05
FeO	2.66	Al	0.03	Σ2M ^H	1.80
MnO	2.42			Σ3M ^o	
MgO	0.26			Ba	0.88
CaO	0.88			K	0.61
SrO	4.11	Ti ⁴⁺	0.94	Sr	0.33
BaO	16.37	Fe ³⁺	0.05	Ca	0.05
K ₂ O	3.52	Nb	0.01	Σ2A ^P	1.87
Na ₂ O	7.90	Σ1M ^o	1.00		
F	1.69				
H ₂ O**	0.77			F ⁻	0.73
O=F	-0.71			(OH) ⁻	0.70
Total	98.33			O ²⁻	0.57
				Σ2X ^A	2.00

* formula contents on the basis of 4 Si⁴⁺ (*apfu*).

** calculated from structure refinement.

TABLE 3. MISCELLANEOUS REFINEMENT DATA FOR BARYTOLAMPROPHYLLITE

a (Å)	19.8971(17)	Absorption coefficient (mm ⁻¹)	6.10
b	7.1165(3)	<i>F</i> (000)	763.2
c	5.4108(2)	<i>D</i> _{calc} (g/cm ³)	3.521
β (°)	96.676(2)	Crystal size (mm)	0.03 × 0.10 × 0.16
V (Å ³)	760.96(3)	Radiation/ filter	MoK α /graphite
Space group	<i>C</i> 2/m	2θ range for data collection (°)	59.86
Z	2	R(int) (%)	2.9
Reflections collected	8747		
Independent reflections	1185		
<i>F</i> _o > 4σ <i>F</i>	1175		
Refinement method	Full-matrix least squares on <i>F</i> ² , fixed weights proportional to 1/σ <i>F</i> ²		
Goodness of fit on <i>F</i> ²	1.12		
Final <i>R</i> (obsd) (%) [<i>F</i> _o > 4σ <i>F</i>]	2.82		
<i>R</i> indices (all data) (%)	<i>R</i> ₁ = 2.87, <i>wR</i> ₂ = 7.44, GoF = 1.12		

Anion considerations

There are four anion sites, O(2)–O(5), occupied by O atoms that constitute the tetrahedral coordination of the Si atoms (Tables 5, 7). There is one O site, O(6), which is the common anion for the M^H polyhedron and three octahedra of the O sheet: $M^H(2)$ and two $M^H(3)$ (Tables 5, 7); this is the X_M^o anion in the terminology of Sokolova (2006). The X_M^o anion receives a bond-valence sum of 0.68 vu from $M^H(2)$ and two $M^H(3)$ cations (Table 7, Fig. 1a). To satisfy the bond-valence requirements of the X_M^o atom, additional bond-valence

(1.32 vu) is needed from the M^H cation. In fact, the M^H cation provides a bond valence of 1.42 vu , and the corresponding $M^H-X_M^o$ distance is very short, 1.686 Å (Table 5). This particular distance is in accord with the bond topology of Group III, where the bond length varies from 1.684 (lamprophyllite-2M) to 1.77 Å (epistolite) (Sokolova 2006). There is one site, O(1), which corresponds to 2 $apfu$. This is the X_A^o site in the terminology of Sokolova (2006) (Fig. 1a). This site receives a bond-valence of 1.03 vu (Table 7), and hence is dominated by monovalent anions. The chemical analysis gives 0.73 $apfu$ F (Table 2), and to satisfy the electroneutrality requirements, we assign $F_{0.73}OH_{0.70}O_{0.57}$ to this site. We label it O(1) for better comparison with the crystal structure of barytolamprophyllite (Rastsvetaeva *et al.* 1995) and lamprophyllite (Krivovichev *et al.* 2003). In minerals of Group III, the X_A^o site ideally corresponds to $(OH)_2$ (Sokolova 2006).

Structure topology

In the crystal structure of barytolamprophyllite, the O sheet is of the form $M^oX_2^o$, where M^o are cations and X^o are anions (Fig. 1a). Structures with the TS block have characteristic two-dimensional minimal cell in the plane of the O sheet with $t_1 \approx 5.5$ and $t_2 \approx 7.1$ Å, $t_1 \wedge t_2 \approx 90^\circ$. This minimal cell contains 4 $[M^oX_2^o]$ (shown in red in Fig. 1a). In the H sheet, (SiO_4) tetrahedra link together to form (Si_2O_7) groups. These (Si_2O_7) groups and M^H square pyramids share common vertices to form the H sheet (Fig. 1b). The [10]-coordinated A^P cations are displaced from the plane of the H sheet and are located just above the centers of large voids in the H sheet (Fig. 1b). The H and O sheets link via common vertices of M^o , Si and M^H polyhedra to form the TS block (Fig. 1c). There are two TS blocks per unit cell in the crystal structure of barytolamprophyllite (Fig. 1c). Along the a direction, the TS blocks alternate with layers of A^P atoms, which constitute the I block. The crystal structure of barytolamprophyllite is topologically identical to those of lamprophyllite and nabalamprophyllite.

CHEMICAL FORMULAE FOR LAMPROPHYLLITE, NABALAMPHYLLITE AND BARYTOLAMPHYLLITE

General formula

Sokolova (2006) wrote the following general formula for the TS block within the planar cell: $A_{2+}^{P_2}B_{2-}^{P_2}M_{12}^{H_2}M_{12}^{o_4}(Si_2O_7)_2X_{4+}$, where M^o = cations of the O sheet, M^H = cations of the H sheet and A^P , B^P = cations at the peripheral (P) sites, $X_{4+} = X_{4+}^o + X_{M2}^P + X_{A2}^P$, where X_{4+}^o anions are common vertices of M^o octahedra and two M^H and two A^P polyhedra (they are the X_M^o and X_A^o anions), and anions X_M^P and X_A^P belong to the M^H and A^P polyhedra on the outside of the TS block




FIG. 1. The crystal structure of barytolamprophyllite: the O sheet (a) and the H sheet (b) of the TS block as viewed down [100]; general view of the crystal structure projected onto (001) (c). (SiO_4) tetrahedra are orange, Na octahedra are navy blue; Ti polyhedra, $^{[5]}M^H$ and $^{[6]}M^H(1)$, are yellow, the A^P atoms are shown as raspberry circles, the planar cell with translations $t_1 \approx 5.5$ Å and $t_2 \approx 7$ Å is shown in red.

TABLE 4. FINAL COORDINATES AND DISPLACEMENT PARAMETERS OF ATOMS IN BARYTOLAMPHYLLITE

Atoms	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₂₃	<i>U</i> ₁₃	<i>U</i> ₁₂	<i>U</i> _{eq}	
A ^P	0.28240(2)	0		0.26285(6)	0.0158(2)	0.00744(18)	0.00770(19)	0	0.00154(12)	0	0.01030(13)
M ^H	0.14603(3)	0		0.70520(11)	0.0103(3)	0.0069(3)	0.0047(3)	0	0.0014(2)	0	0.0073(2)
M ^O (1)	0	½	0	0.0301(6)	0.0060(5)	0.0335(7)	0	-0.0225(5)	0	0.0253(3)	
M ^O (2)	0	0	0	0.0235(12)	0.0173(12)	0.0180(12)	0	0.0093(9)	0	0.0191(5)	
M ^O (3)	0	0.26016(15)	½	0.0230(6)	0.0135(5)	0.0141(5)	0	0.0053(4)	0	0.0166(3)	
Si	0.13981(3)	0.28427(10)	0.02040(12)	0.0118(3)	0.0053(3)	0.0068(3)	0.0003(2)	0.0014(2)	-0.0001(2)	0.0079(2)	
O(1)	0.44479(16)	0	0.2738(6)	0.0230(16)	0.0272(18)	0.0268(17)	0	0.0103(12)	0	0.0251(10)	
O(2)	0.17024(10)	0.1883(3)	0.4639(3)	0.0182(9)	0.0101(9)	0.0091(8)	0.0023(7)	0.0012(7)	0.0004(7)	0.0125(4)	
O(3)	0.17064(10)	0.1874(3)	0.9686(3)	0.0178(9)	0.0090(8)	0.0092(8)	-0.0028(6)	0.0033(7)	-0.0005(7)	0.0119(4)	
O(4)	0.17175(14)	½	0.2170(5)	0.0142(12)	0.0060(11)	0.0107(12)	0	0.0015(9)	0	0.0103(5)	
O(5)	0.05813(10)	0.2950(3)	0.1705(4)	0.0116(8)	0.0102(9)	0.0162(9)	0.0002(7)	0.0006(7)	0.0013(7)	0.0128(4)	
O(6)	0.06071(15)	0	0.6672(6)	0.0121(13)	0.0356(18)	0.0133(13)	0	0.0025(10)	0	0.0202(7)	

TABLE 5. INTERATOMIC DISTANCES (Å) AND ANGLES (°) IN BARYTOLAMPHYLLITE

M ^O (1) – O(1)	1.944(3)	×2	M ^O (2) – O(6) _a	2.282(3)	×2
M ^O (1) – O(5)	2.017(2)	×4	M ^O (2) – O(5) _b	2.520(2)	×4
<M ^O (1) – O>	1.993		<M ^O (2) – O>	2.441	
M ^O (3) – O(5)	2.249(2)	×2	M ^H – O(6)	1.686(3)	
M ^O (3) – O(1) _b	2.303(3)	×2	M ^H – O(2)	1.969(2)	×2
M ^O (3) – O(6)	2.335(2)	×2	M ^H – O(3)	1.972(2)	×2
<M ^O (3) – O>	2.296		<M ^H – O>	1.914	
Si – O(5)	1.616(2)		A ^P – O(3) _b	2.768(2)	×2
Si – O(3) _b	1.623(2)		A ^P – O(2) _b	2.770(2)	×2
Si – O(2)	1.624(2)		A ^P – O(4) _d	2.850(3)	
Si – O(4)	1.660(1)		A ^P – O(4) _b	2.856(3)	
<Si – O>	1.631		A ^P – O(3) _b	2.904(2)	×2
Si – O(4) – Si _c	135.3(2)		A ^P – O(2) _b	2.920(2)	×2
			<A ^P – O>	2.843	

a: x, y, z – 1; b: -x + ½, -y + ½, -z + 1; c: x, -y + 1, z; d: -x + ½, -y + ½, -z.

[in the intermediate space (**I**) between two TS blocks]; n is the number of X^P anions: n = 0, 2, 4 depending on the coordination of the M^H site and the position of the A^P polyhedron relative to the adjacent TS block. The core of the TS block, M^H₂ M^O₄ (Si₂O₇)₂ X₄, is shown in bold; the stoichiometry of this part of the TS block is invariant.

Sokolova (2006) wrote the general formula of lamprophyllite, barytolamprophyllite and nabalamprophyllite as A^P₂ M^H₂ M^O₄ (Si₂O₇)₂ X₄ (see Table 1), where A^P₂ represents the **I** block and M^H₂ M^O₄ (Si₂O₇)₂ X₄ represents the TS block. In barytolamprophyllite, the total charge of the TS block [of ideal composition Ti₂ Na₃ Ti (Si₂O₇)₂ O₂ (OH)₂] is -3, and the **I** block has an ideal composition (BaK) with a total charge of +3. We thus write the ideal formula of barytolamprophyllite as (BaK)Ti₂ Na₃ Ti (Si₂O₇)₂ O₂ (OH)₂ or (BaK) Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂. Note that this ideal formula is neutral. The ideal formulae of lamprophyllite, (SrNa) Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂, and nabalamprophyllite, Ba Na Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂, also are neutral.

Why, in several cases, have the ideal formulae for **lamprophyllite**, (Sr,Ba)₂ Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂, and **barytolamprophyllite** (Ba,Sr)₂ Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂, been written with an excess charge of +1? Let us examine the formula (Sr,Ba)₂ Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂. The aggregate charge of the fixed part of the formula, Na₃ Ti₂ (Si₂O₇)₂ O₂ (OH)₂ is 1⁺ × 3 + 4⁺ × 3 + 2⁻ × 2 + 4⁺ × 4 + 2⁻ × 14 + 1⁻ × 2 = 3⁻. A stable structure *must* accord with the electroneutrality principle, and hence the aggregate charge of the alkali and alkaline-earth part of the formula must have a net charge of 3⁺. To write the formula as (Sr,Ba)₂ violates the electroneutrality principle, as (Sr,Ba)₂ has a net charge of 4⁺. Inspection of Table 2 indicates that monovalent cations form a major part of the A^P site: 0.61 K. The properties of an end-member include the following requirements: (1) the end-member formula must be fixed, and (2) the structure must show the maximum degree of order consonant with the electroneutrality principle. For barytolamprophyllite examined here, the resulting end-member formula is (BaK) Na₃ Ti (Si₂O₇)₂ O₂ (OH)₂; the parentheses indicate that Ba and K occupy the same site, but there is no comma, as their relative amounts in the end-member are fixed by the electroneutrality principle.

Reconsideration of the analytical data

Consider the chemical data for lamprophyllite and barytolamprophyllite, taken from literature (Table 8). There are the first analytical data on these minerals, compiled by Anthony *et al.* (1995): Dana (1899), Vlasov (1966) for lamprophyllite, and Dudkin (1959) and Peng & Chang (1965) for barytolamprophyllite. Comparatively recent analyses were done by Johnsen *et al.* (1994) for lamprophyllite and barytolamprophyllite. Three analyses were done on samples used for crystal-structure refinement: lamprophyllite (Krivovichev *et al.* 2003), barytolamprophyllite (this work) and nabalam-

TABLE 6. REFINED SITE-SCATTERING VALUES ($epfu$) AND ASSIGNED SITE-POPULATIONS ($apfu$) FOR BARYTOLAMPROPHYLLITE

Site	Site-scattering	Site population	Calculated site-scattering	$\langle X-\phi \rangle_{\text{calc}}^*$	$\langle X-\phi \rangle_{\text{obs}}$
A^P	77.2(2)	$0.88 \text{ Ba} + 0.61 \text{ K} + 0.33 \text{ Sr} + 0.05 \text{ Ca} + 0.13 \square$	74.4	2.886	2.843
M^H	43.0(2)	$1.97 \text{ Ti} + 0.03 \text{ Al}$	43.7	1.890	1.914
$M^\circ(1)$	22.0(1)	$0.94 \text{ Ti} + 0.05 \text{ Fe}^{3+} + 0.01 \text{ Nb}$	22.4	1.975	1.993
$M^\circ(2)$	11.0(1)	1.00 Na		11.0	2.400
$M^\circ(3)$	30.5(2)	$1.09 \text{ Na} + 0.30 \text{ Fe}^{2+} + 0.28 \text{ Mn}^{2+} + 0.08 \text{ Ca} + 0.05 \text{ Mg} + 0.20 \square$	30.0	2.304	2.441
O(1)	17.0	$0.73 \text{ F} + 0.70 \text{ OH} + 0.57 \text{ O}$	16.7		2.296

* Ionic radii for $^{[4]} \text{O}^{2-}$, $^{[4]} \text{OH}^-$, $^{[4]} \text{F}^-$, Ti, Nb, Al, Mg, Mn^{2+} , Fe^{2+} , Fe^{3+} , Na, Ba, Sr, K are from Shannon (1976); ϕ : unspecified anion.

TABLE 7. BOND-VALENCE* TABLE (vu) FOR BARYTOLAMPROPHYLLITE

Atom	Si	A^P	M^H	$M^\circ(1)$	$M^\circ(2)$	$M^\circ(3)$	Σ
O(1)				0.63 ^{x2} :		0.20 ^{x2} :	1.03
O(2)	1.02	0.20 ^{x2} : 0.14 ^{x2} :	0.64 ^{x2} :				2.00
O(3)	1.02	0.20 ^{x2} : 0.14 ^{x2} :	0.63 ^{x2} :				1.99
O(4)	0.92 ^{x2} -	0.16 ^{x2} :					2.16
O(5)	1.04			0.56 ^{x4} :	0.17 ^{x4} :	0.25 ^{x2} :	2.02
O(6)				1.42	0.26 ^{x2} :	0.21 ^{x2} :-	2.10
Total Aggr. charge	4.00	11.68	3.96	3.50	1.20	1.32	
	4.00	1.59	3.99	3.96	1.00	1.23	

* Bond-valence parameters are from Brown (1981).

prophyllite (Chukanov *et al.* 2004). We have calculated chemical formulae for these nine samples on the basis of 4 Si $apfu$ and have assigned site populations for groups of atoms in accord with the lamprophyllite-type structure. In compositions (1) and (6), the sum of cations is in excess of the number of available sites by 0.5 $apfu$; therefore, these analyses contain significant errors. The sum of the anions (ideally 18) is lower in many cases and is close to 18.7 pfu in composition (6). Although the results in Table 8 are not very accurate, we can still see that: (1) the two A^P sites include A^{2+} and A^+ cations; (2) at the A^P site, divalent cations are slightly dominant over monovalent cations.

We now modify our calculation of the formulae, taking into account the crystal chemistry of the lamprophyllite structure-type (Sokolova 2006): (1) monovalent anions are dominant at the X_A° site, giving 2 $apfu$, and the content of H_2O in the chemical composition can be calculated from the equation, $\text{F} + \text{OH} = 2 \text{ apfu}$; (2) the Ti-dominant $M^\circ(1)$ site is invariably fully occupied and can contain Ti, Nb, Fe^{3+} and Al; (3) Fe^{2+} occurs at

one of the 3 M° sites in the O sheet; (4) all anion sites are fully occupied, and the sum of the anions must be 18 $apfu$. We discard compositions (1) and (6) of Dana (1889) and Peng & Chang (1965), respectively (Σ cations = 12.54 and 12.50 instead of 12.00 $apfu$), neglect any content of Cl (as there is no site suitable for this anion in the crystal structure), and reconsider the $\text{Fe}^{3+}/\text{Fe}^{2+}$ values. Modified compositions are given in Table 9. Consideration of the unit formulae gives us certain information necessary to write ideal formulae and chemical formulae in general.

(1) At the two A^P sites, both divalent (Ba, Sr, Ca) and monovalent (K, Na) cations are present; divalent cations are dominant over monovalent cations; the total charge of two A^P cations is invariably less than +3.5.

(2) The sum of cations at three Ti-dominant sites ($M^H + M^\circ$) is 3 $apfu$, and the total charge is invariably more than +11.5.

(3) At the three M° sites, Na^+ is the dominant cation, $\text{Na} > 2 \text{ apfu}$. The total charge varies from +3.75 to +3.13.

(4) The total charge of cations [4 Si + 2 A^P + Ti-dominant (2 $M^H + M^\circ$) + Na-dominant 3 M°] varies from +34.81 to +34.21. The total charge of anions results from 14 O atoms [of (SiO_4) tetrahedra], 2 O atoms (2 X_M°) and 2 X_A° anions. The 16 O atoms give aggregate charge of -32. The bond-valence sum at the two X_A° anions varies from +2.81 to +2.21 vu , with a mean value of +2.53 vu . To satisfy the electroneutrality of the unit formula, we assign OH, F and O to the two X_A° sites. Our assignments show that monovalent anions are always dominant at the two X_A° sites (Table 9).

We can now more accurately calculate the content of H_2O in each case and make a final calculation of the unit formula. In Table 10, the sums of the cations vary from 12.05 to 11.67 $apfu$ (ideally 12), and the sums of the anions vary from 17.97 to 18.04 $apfu$ (ideally 18). This is a very good convergence and shows that the chemical formulae have been calculated appropriately. Note that in composition (4) (Dudkin 1959), $\text{Sr} > \text{Ba} \text{ apfu}$, and therefore this mineral is Ba-rich lamprophyllite,

TABLE 8. CHEMICAL COMPOSITION (wt %) AND UNIT FORMULA* (*apfu*) OF LAMPROPHYLLITE, BARYTOLAMPROMPHYLITE AND NABALAMPROMPHYLITE[§]

mineral	lamprophyllite				barytolam- prophyllite				nabalamp- prophyllite			
	Sr > Ba				Ba > Sr							
sample	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)			
SiO ₂	30.40	30.93	31.29	32.2	30.95	28.53	29.9	29.31	28.75			
Nb ₂ O ₆	~	0.16	~	0.43	~	~	~	0.11	~			
TiO ₂	27.48	27.14	29.27	30.30	28.15	26.6	28.68	28.39	27.8			
Al ₂ O ₃	~	0.56	0.10	~	0.22	1.12	0.04	0.16	0.44			
Fe ₂ O ₃	3.87	1.36	~	~	0.89	2.72	~	0.49	0.78			
FeO	1.97	2.31	0.66	2.08	4.15	2.63	1.27	2.66	~			
MnO	2.33	3.46	3.54	3.49	1.05	1.75	1.60	2.42	1.10			
MgO	0.25	0.40	0.33	0.66	0.88	1.00	1.03	0.26	0.34			
CaO	1.41	2.48	0.62	0.84	1.64	1.70	0.26	0.88	0.36			
SrO	14.58	14.12	14.89	16.43	7.99	1.47	3.48	4.11	0.65			
BaO	1.25	3.49	5.05	0.49	10.51	17.24	20.06	16.37	24.12			
K ₂ O	2.31	1.26	0.65	0.45	1.82	3.10	0.94	3.52	0.94			
Na ₂ O	12.35	10.64	11.34	12.38	9.8	9.52	11.99	7.90	11.14			
F	1.82	1.10	~	~	1.45	1.60	~	1.69	1.18			
CL	~	~	~	~	0.04	0.51	~	~	~			
H ₂ O	~	0.56	~	~	0.99	0.70	~	0.77**	1.83			
O=F	-0.77	-0.46	~	-0.61	-0.67	~	-0.71	-0.50	~			
O=Cl	~	~	~	~	-0.01	-0.12	~	~	~			
Total	99.25	99.51	97.74	99.75	99.89	99.4	99.25	98.33	98.93			
Si	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00			
Ba	0.06	0.18	0.25	0.02	0.53	0.95	1.05	0.88	1.32			
Sr	1.11	1.06	1.10	1.18	0.60	0.12	0.27	0.33	0.05			
Ca	0.10	0.17	0.05	0.12	0.12	0.13	0.04	0.05	~			
ΣA^{2+}	1.27	1.41	1.40	1.32	1.25	1.20	1.36	1.26	1.37			
K	0.39	0.21	0.11	0.07	0.30	0.55	0.16	0.61	0.17			
Na	0.44	0.47	0.36	0.66	0.13	0.33	0.52	~	0.50			
ΣA^{2+}	0.83	0.68	0.47	0.73	0.43	0.88	0.68	0.61	0.67			
ΣA^P	2.10	2.09	1.87	2.05	1.68	2.07	2.04	1.87	2.04			
Ti	2.72	2.64	2.81	2.83	2.74	2.81	2.89	2.91	2.91			
Nb	~	0.01	~	0.02	~	~	~	0.01	~			
Fe ³⁺	0.38	0.13	~	~	0.09	0.29	~	0.05	0.08			
Al	~	0.09	0.02	~	0.03	0.19	0.01	0.03	0.07			
$\Sigma 3(M^H + M^O)$	3.10	2.87	2.83	2.85	2.86	3.29	2.90	3.00	3.06			
Na	2.71	2.20	2.45	2.32	2.15	2.26	2.59	2.09	2.51			
Fe ²⁺	0.22	0.25	0.07	0.22	0.45	0.31	0.14	0.30	~			
Mn ²⁺	0.26	0.38	0.38	0.37	0.12	0.21	0.18	0.28	0.13			
Ca	0.10	0.17	0.04	~	0.11	0.13	~	0.08	0.05			
Mg	0.05	0.08	0.06	0.12	0.17	0.21	0.21	0.05	0.07			
$\Sigma 3M^O$	3.34	3.08	3.00	3.03	3.00	3.12	3.12	2.80	2.76			
F ⁻	0.76	0.45	~	~	0.59	0.71	~	0.73	0.52			
(OH) ⁻	~	0.48	~	~	0.85	0.66	~	0.70	1.70			
Cl ⁻	~	~	~	~	0.01	0.12	~	~	~			
ΣX^{\pm}	0.76	0.93	~	~	1.45	1.49	~	1.43	2.22			
Cations	12.54	12.03	11.70	11.93	11.70	12.50	12.05	11.67	11.86			
Σ anions	18.06	17.82	17.07	17.27	17.85	18.68	17.30	17.98	18.23			

* formula contents on a basis of 4 Si *apfu*.

** H₂O content calculated from results of structure refinement, $2X^{\pm}_A$ (*apfu*) = 0.73F + 0.70OH + 0.57O.

Sources: (1) Dana (1899), (2) Vlasov (1966), (3) Johnsen *et al.* (1994); sample MM55312, (4) Krivovichev *et al.* 2003, (5) Dudkin (1959); sample 1, (6) Peng & Chang (1965), (7) Johnsen *et al.* (1994); sample NM6004/BB, (8) this work, (9) Chukanov *et al.* (2004). [§] The chemical data reported here are as presented in the original articles.

TABLE 9. CHEMICAL COMPOSITION (wt %) AND UNIT FORMULA* (*apfu*) OF LAMPROPHYLLITE, BARYTOLAMPROMPHYLITE AND NABALAMPROMPHYLITE[§]

mineral	lamprophyllite				barytolam- prophyllite				nabalamp- prophyllite			
	Sr > Ba				Ba > Sr							
sample	(1)	(2)	(3)	(4)	(5)	(6)	(7)					
Fe ₂ O ₃ [†]	2.62	0.73	1.58	2.32	1.00	0.49	0.20					
FeO [†]	1.18	~	0.66	2.86	0.37	2.66	0.52					
H ₂ O ^{**}	1.80	2.35	2.41	1.63	2.24	0.77	1.60					
O=F	-0.46	~	~	-0.61	~	-0.71	-0.50					
Total	100.87	100.16	102.32	100.64	101.59	98.33	98.64					
Si	4.00	4.00	4.00	4.00	4.00	4.00	4.00					
Ba	0.18	0.25	0.02	0.53	1.05	0.88	1.32					
Sr	1.06	1.11	1.18	0.60	0.27	0.33	0.05					
Ca	0.17	0.05	0.06	0.10	0.04	0.05	~					
ΣA^{2+}	1.41	1.41	1.26	1.23	1.36	1.26	1.37					
K	0.21	0.11	0.07	0.30	0.16	0.61	0.17					
Na	0.40	0.39	0.61	0.30	0.54	~	0.50					
ΣA^{2+}	0.61	0.50	0.68	0.60	0.70	0.61	0.67					
ΣA^P	2.02	1.91	1.94	1.83	2.06	1.87	2.04					
ΣA^P charge	+3.40 ^{††}	+3.32	+3.20	+3.06	+3.32	+3.13	+3.34					
Ti	2.66	2.81	2.83	2.74	2.89	2.89	2.89					
Nb	0.01	~	0.02	~	~	~	0.01					
Fe ³⁺	0.26	0.07	0.15	0.23	0.10	0.05	0.02					
Al	0.09	0.02	~	0.03	0.01	0.03	0.01					
$\Sigma 3(M^H + M^O)$	3.02	3.00	3.00	3.00	3.00	3.00	3.00					
charge	+11.66	+11.51	+11.87	+11.74	+11.89	+11.93	+11.91					
Na	2.27	2.42	2.37	2.16	2.57	2.09	2.51					
Fe ²⁺	0.13	~	0.07	0.31	0.04	0.30	0.06					
Mn ²⁺	0.38	0.38	0.37	0.12	0.18	0.28	0.13					
Ca	0.17	0.04	0.06	0.13	~	0.08	0.05					
Mg	0.08	0.06	0.12	0.17	0.21	0.05	0.07					
$\Sigma 3M^O$	3.03	2.90	2.99	2.89	3.00	2.80	2.82					
3M ⁰ charge	+3.75 ^{††}	+3.38	+3.61	+3.62	+3.43	+3.51	+3.13					
Total charge of cations	+34.81	+34.21	+34.68	+34.43	+34.64	+34.57	+34.38					
BVS [†] at $2X^{\pm}_A$	+2.81	+2.21	+2.68	+2.43	+2.64	+2.57	+2.38					
F ⁻	0.45	~	~	0.59	~	0.73	0.52					
(OH) ⁻	0.74	1.79	1.32	0.99	1.33	0.70	1.10					
O ²⁻	0.81	0.21	0.68	0.41	0.67	0.57	0.38					
$\Sigma 2X^{\pm}_A$	2.00	2.00	2.00	2.00	2.00	2.00	2.00					
cations	12.03	11.70	11.93	11.70	12.05	11.67	11.86					

* As is made clear in these footnotes, the chemical data from original papers have been modified.

[†] wt % of SiO₂, Nb₂O₆, TiO₂, Al₂O₃, MnO, MgO, CaO, SrO, BaO, K₂O, Na₂O and F are the same as in Table 8, formula contents on a basis of 4 Si *apfu*.

^{**} H₂O content is calculated in accord with F + OH = 2 *apfu*, except for this work (6).

^{††} The ratio Fe^{3+*/Fe²⁺}

was recalculated from the crystal-topology basis (see text).

[‡] Where sum of cations exceeds the multiplicity of a site (or a group of sites), the charge is normalized to the multiplicity of a site (or a group of sites).

[†] bond-valence sum. Sources: (1) Vlasov (1966), (2) Johnsen *et al.* (1994); sample MM55312, (3) Krivovichev *et al.* 2003, (4) Dudkin (1959); sample 1, (5) Johnsen *et al.* (1994); sample NM6004/BB, (6) this work, (7) Chukanov *et al.* (2004).

not barytolamprophyllite [note that Krivovichev *et al.* (2003) erroneously referred to Dudkin (1959) as the author of the first description of a *Ba-analogue of lamprophyllite, barytolamprophyllite*]. Note also that in composition (7) (Table 8) [= composition (5) (Tables 9, 10)] (Johnsen *et al.* 1994), K < Na *apfu* at the AP site,

and therefore this mineral is actually nabalamprophyllite, not barytolamprophyllite.

On the ideal formulae of lamprophyllite, nabalamprophyllite and barytolamprophyllite

We showed above (Tables 9, 10) that the total charge at three Ti-dominant sites, $[2M^H + M^O(1)]$, is invariably more than +11.5 (ideally +12), and that the ideal composition of three Ti-dominant sites is Ti_3 . In every case, Na is dominant at three M^O sites $[M^O(2) + 2 M^O(3)]$,

TABLE 10. CHEMICAL COMPOSITION (wt.%) AND FINAL UNIT-FORMULA^a
(*apfu*) FOR LAMPROPHYLLITE, BARYTOLAMPHYLLITE AND
NABALAMPHYLLITE

mineral	lamprophyllite Sr>Ba				barytolampro- phyllite, Ba>Sr			nebalam- phyllite
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	
H ₂ O ^{**}	0.86	2.1	1.59	1.14	1.5	0.77	1.19	
Si	4.00	4.00	4.00	4.00	4.00	4.00	4.00	
Nb	0.01	—	0.02	—	—	0.01	—	
Ti	2.64	2.81	2.83	2.74	2.89	2.91	2.91	
Al ³⁺	0.09	0.02	0	0.03	0.01	0.03	0.07	
Fe ^{3+†}	0.26	0.07	0.15	0.23	0.10	0.05	0.02	
Fe ^{2+*}	0.13	—	0.07	0.31	0.04	0.30	0.06	
Mn ^{2+*}	0.38	0.38	0.37	0.12	0.18	0.28	0.13	
Mg	0.08	0.06	0.12	0.17	0.21	0.05	0.07	
Ca	0.34	0.09	0.11	0.23	0.04	0.13	0.05	
Sr	1.06	1.10	1.18	0.60	0.27	0.33	0.05	
Ba	0.18	0.25	0.02	0.53	1.05	0.88	1.32	
K	0.21	0.11	0.07	0.30	0.16	0.61	0.17	
Na	2.67	2.81	2.98	2.46	3.11	2.09	3.01	
F	0.45	—	—	—	—	0.73	0.52	
H ⁺	0.74	1.79	1.32	0.98	1.34	0.70	1.10	
Σ cations	12.03	11.70	11.93	11.70	12.05	11.67	11.86	
Σ anions	18.01	18.00	18.01	17.97	18.02	17.98	18.04	

Note that the chemical data from original papers have been modified, as explained here. * The wt.% values are the same as in Table 9; formula contents are calculated on a basis of 4 Si⁴⁺ *apfu*; ** H₂O content was calculated on the basis of electroneutrality requirements. † Fe^{3+†}/Fe^{2+*} was calculated as in Table 9.
Columns: (1) Vlasov (1966), (2) Johnsen *et al.* (1994), sample MM56312, (3) Krivovichev *et al.* 2003, (4) Dukin (1959), sample 1, (5) Johnsen *et al.* (1994), sample NM6004/8B, (6) This work, (7) Chukanov *et al.* (2004).

Na > 2 *apfu* (Tables 9, 10), and we can write the ideal composition of the three M^o sites as Na₃. Monovalent anions, OH + F, are invariably dominant at the two X^o_A sites, and we can write the ideal composition of the two X^o_A anions as (OH)₂. So far, we can write the ideal formula as A^P₂ Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂, which leaves us with A^P₂ to assign with a total charge of +3. Note that nabalamphyllite is the only mineral with lamprophyllite-type structure in which there are two independent Ba- and Na-dominant A^P sites. The observed total charge of the two A^P cations is in every case less than +3.5 (Tables 9, 10) and ideally is +3; therefore, we write the ideal composition of the two A^P sites as (A²⁺ A⁺). This gives us the ideal formula (A²⁺ A⁺) Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂ for

$$\begin{array}{ll} \text{lamprophyllite} & A^{2+} A^+ = (\text{SrNa}), \\ \text{barytolamprophyllite} & A^{2+} A^+ = (\text{BaK}), \\ \text{nabalamprophyllite} & A^{2+} A^+ = (\text{BaNa}), \end{array}$$

and suggests another possible composition: (SrK) Na₃ Ti₃ (Si₂O₇)₂ O₂ (OH)₂.

ACKNOWLEDGEMENTS

We are very grateful to Adriana and Renato Pagano for the sample of barytolamprophyllite from their collection of minerals. We thank the Editor, Robert F.

Martin, and Ole Johnsen and an anonymous referee, for useful comments and suggestions. ES thanks Frank C. Hawthorne for critical review of the manuscript. Fernando Cámaras was supported by funding from CNR-IGG and Italian MIUR-PRIN 2005 project "From minerals to materials: crystal-chemistry, microstructures, modularity, modulations".

REFERENCES

- ANTHONY, J.W., BIDEAUX, R.A., BLADH, K.W. & NICHOLS, M.C. (1995): *Handbook of Mineralogy*. II. *Silica, Silicates*. Mineral Data Publishing, Tucson, Arizona.
- BACK, M.E. & MANDARINO, J.A. (2008): *Fleischer's Glossary of Mineral Species 2008*. The Mineralogical Record Inc., Tucson, Arizona.
- BROWN, I.D. (1981): The bond-valence method: an empirical approach to chemical structure and bonding. In *Structure and Bonding in Crystals II* (M. O'Keeffe & A. Navrotsky, eds.). Academic Press, New York, N.Y. (1-30).
- CHERNOV, A.N., ILYUKHIN, V.V., MAKSYMOW, B.A. & BELOV, N.V. (1971): Crystal structure of innelite, Na₂Ba₃(Ba,K,Mn)(Ca,Ba)Ti(TiO₂)₂(Si₂O₇)₂(SO₄)₂. Sov. Phys. Crystallogr. **16**, 65-69.
- CHUKANOV, N.V., MOISEEV, M.M., PEKOV, I.V., LAZEBNIK, K.A., RASTSVETAeva, R.K., ZAYAKINA, N.V., FERRARIS, G. & IVALDI, G. (2004): Nabalamphyllite Ba(Na,Ba){[Na₃Ti₃(TiO₂O₂Si₄O₁₄)₂] (OH,F)₂} – a new layer titanosilicate of the lamprophyllite group from Inagli and Kovdor alkaline-ultrabasic massifs, Russia. Zap. Vser. Mineral. Obshchest. **133**(1), 59-72 (in Russ.).
- DANA, E.S. (1899): *Dana's System of Mineralogy*, (6th ed.). First Appendix. John Wiley & Sons, New York, N.Y. (40).
- DUKIN, O.B. (1959): On barium lamprophyllite. Zap. Vses. Mineral. Obshchest. **88**, 713-715 (in Russ.).
- ERCIĆ, T.S., COOPER, M.A. & HAWTHORNE, F.C. (1998): The crystal structure of vuonnemite, Na₁₁Ti⁴⁺Nb₂(Si₂O₇)₂(P₂O₇)₂O₃(F,OH), a phosphate-bearing sorosilicate of the lomonosovite group. Can. Mineral. **37**, 1311-1320.
- INTERNATIONAL TABLES FOR X-RAY CRYSTALLOGRAPHY (1992): Vol. C. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- JOHNSEN, O. (1996): TEM observations and X-ray powder data on lamprophyllite polytypes from Gardiner Complex, East Greenland. Neues Jahrb. Mineral., Monatsh., 407-417.
- JOHNSEN, O., NIELSEN, T.F.D. & RØNSBO, J.G. (1994): Lamprophyllite and barytolamprophyllite from the Tertiary Gardiner complex, East Greenland. Neues Jahrb. Mineral., Monatsh., 328-336.
- KHOMYAKOV, A.P. (1995): *Mineralogy of Hyperagpaitic Alkaline Rocks*. Clarendon Press, Oxford, U.K.

- KRIVOVICHEV, S.V., ARMBRUSTER, T., YAKOVENCHUK, V.N., PAKHOMOVSKY, YA.A. & MEN'SHIKOV, YU.P. (2003): Crystal structures of lamprophyllite-*2M* and lamprophyllite-*2O* from the Lovozero alkaline massif, Kola peninsula, Russia. *Eur. J. Mineral.* **15**, 711-718.
- PENG, TZE-CHUNG & CHANG, CHIEN-HUNG (1965): New varieties of lamprophyllite–barytolamprophyllite and orthorhombic lamprophyllite. *Scientia Sinica* **14**, 1827-1840.
- PENG, ZHIZHONG, ZHANG, JIANHONG & SHU, JINFU (1984): The crystal structure of barytolamprophyllite. *Kexue Tongbao* **29**, 237-241.
- POUCHOU, J.L. & PICHOIR, F. (1985): "PAP" $\phi(\rho)$ procedure for improved quantitative microanalysis. In *Microbeam Analysis*. San Francisco Press, San Francisco, California (104-106).
- RAMSAY, W. & HACKMANN, V. (1894): Das Nephelinsyenitgebiet auf der Halbinsel Kola. *Bull. Soc. Géographie Finlande* **11**, N2, 1-225.
- RASTSVETAeva, R.K. & CHUKANOV, N.V. (1999): Crystal structure of a new high-barium analogue of lamprophyllite with a primitive unit cell. *Dokl. Chem.* **368**(4-6), 228-231.
- RASTSVETAeva, R.K. & DORFMAN, M.D. (1995): Crystal structure of Ba-lamprophyllite in the isomorphous lamprophyllite–barytolamprophyllite series. *Crystallogr. Rep.* **40**, 951-954.
- RASTSVETAeva, R.K., EVSYUNIN, V.G. & KONEV, A.A. (1995): Crystal structure of K-barytolamprophyllite. *Crystallogr. Rep.* **40**, 472-474.
- SHANNON, R.D. (1976): Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallogr.* **A32**, 751-767.
- SHELDICK, G.M. (1997): *SHELX-97: Program for the Solution and Refinement of Crystal Structures*. Siemens Energy and Automation, Madison, Wisconsin.
- SHELDICK, G.M. (1998): *SADABS User Guide*. University of Göttingen, Göttingen, Germany.
- SOKOLOVA, E. (2006): From structure topology to chemical composition. I. Structural hierarchy and stereochemistry in titanium disilicate minerals. *Can. Mineral.* **44**, 1273-1330.
- SOKOLOVA, E. & CÁMARA, F. (2007): From structure topology to chemical composition. II. Titanium silicates: revision of the crystal structure and chemical formula of delindeite. *Can. Mineral.* **45**, 1247-1261.
- SOKOLOVA, E. & HAWTHORNE, F.C. (2004): The crystal chemistry of epistolite. *Can. Mineral.* **42**, 797-806.
- VLASOV, K.A., ed. (1966): *Geochemistry and Mineralogy of Rare Elements and Genetic Types of their Deposits*. Israel Program for Scientific Translations, Jerusalem, Israel (II, 207-209).
- WOODROW, P.J. (1964): Crystal structure of lamprophyllite. *Nature* **204**, 375.
- YAKOVENCHUK, V.N., IVANYUK, G.YU., PAKHOMOVSKY, YA.A. & MEN'SHIKOV, YU.P. (2005): *Khibiny*. Laplandia Minerals, Apatity, Russia.

Received May 17, 2007, revised manuscript accepted January 15, 2008.