

Clintonite

Ca(Mg, Al)₃(Al₃Si)O₁₀(OH)₂

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Monoclinic. *Point Group:* 2/m. In tabular pseudo-hexagonal crystals, complexly twinned, to 2.5 cm; foliated or lamellar radiated; massive. *Twinning:* On {001}, twin axis [310] or [3 $\bar{1}$ 0]; spiral polysynthetic twinning.

Physical Properties: *Cleavage:* {001}, perfect. *Tenacity:* Brittle. Hardness = 3.5 on {001}; 6 \perp {001}. D(meas.) = 3.0–3.1 D(calc.) = 3.096

Optical Properties: Transparent to translucent. *Color:* Colorless, yellow, orange, red-brown, brown, green. *Streak:* White, slightly yellow-gray. *Luster:* Vitreous, pearly, submetallic. *Optical Class:* Biaxial (-). *Pleochroism:* X = colorless, pale orange, red-brown; Y = Z = pale brownish yellow, pale green. *Orientation:* Y \simeq a or Y = b; Z = b or \simeq a; X \wedge c = 5°–10°. *Dispersion:* r < v, weak. *Absorption:* Y \simeq Z > X. α = 1.643–1.648 β = 1.655–1.662 γ = 1.655–1.663 2V(meas.) = 2°–40°

Cell Data: *Space Group:* C2/m. a = 5.204 b = 9.026 c = 9.812 β = 100°20' Z = 2

X-ray Powder Pattern: Zlatoust, Russia.

2.56 (100), 3.21 (70), 2.11 (70), 1.505 (60), 9.68 (50), 2.45 (50), 1.485 (50)

Chemistry:	(1)	(2)	(1)	(2)
SiO ₂	18.78	16.74	MgO	20.56
TiO ₂	0.54		CaO	12.90
Al ₂ O ₃	40.00	42.70	F	2.01
Fe ₂ O ₃		2.85	H ₂ O ⁺	[4.20]
FeO	1.86	0.41	–O = F ₂	[0.85]
MnO	0.00		Total	[100.00]
				100.31

(1) Amity, New York, USA; by electron microprobe, H₂O by difference; corresponds to Ca_{0.97}(Mg_{2.15}Al_{0.70}Fe_{0.11}²⁺Ti_{0.03}) Σ =2.99(Al_{2.68}Si_{1.32}) Σ =4.00O_{9.45}[(OH)_{1.55}F_{0.45}] Σ =2.00.

(2) Crestmore, California, USA; corresponds to Ca_{0.98}(Mg_{2.09}Al_{0.70}Fe_{0.15}³⁺Fe_{0.02}²⁺) Σ =2.96(Al_{2.83}Si_{1.17}) Σ =4.00O_{9.92}(OH)_{2.10}.

Polymorphism & Series: 1M polytype; 2M₁ and 3A polytypes rare.

Mineral Group: Mica group.

Occurrence: In chlorite schists; in metasomatically altered limestones; in siliceous skarns near contact metamorphic zones.

Association: Talc, spinel, grossular, vesuvianite, clinopyroxene, monticellite, chondrodite, phlogopite, chlorite, quartz, calcite, dolomite.

Distribution: In the USA, around Amity, Edenville, and Warwick, Orange Co., New York; from Crestmore, Riverside Co., California; near Ludwig, Lyon Co., Nevada; at Sulzer, Prince of Wales Island, Alaska. On Mt. Monzoni and at Mts. Castone and Adamello, Trentino-Alto Adige, Italy. In the Pargas district, Finland. From the Akhmatovsk mine, near Zlatoust, Ural Mountains, Russia. In the Chichibu mine, Saitama Prefecture, Japan. On the Ertzberg, Irian Jaya.

Name: For De Witt Clinton (1769–1828), American statesman.

References: (1) Dana, E.S. (1892) Dana's system of mineralogy, (6th edition), 638–640 [seybertite, xanthophyllite]. (2) Deer, W.A., R.A. Howie, and J. Zussman (1963) Rock-forming minerals, v. 3, sheet silicates, 99–102. (3) Forman, S.A., H. Kodama, and S. Abbey (1967) A re-examination of xanthophyllite [clintonite] from the type locality. Can. Mineral., 9, 25–30. (4) MacKinney, J.A., C.I. Mora, and S.W. Bailey (1988) Structure and crystal chemistry of clintonite. Amer. Mineral., 73, 365–375.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.