

Itoigawaite, a new mineral, the Sr analogue of lawsonite, in jadeitite from the Itoigawa-Ohmi district, central Japan

H. MIYAJIMA¹, S. MATSUBARA², R. MIYAWAKI² AND K. ITO³

¹ Fossa Magna Museum, Itoigawa, Niigata 941-0056, Japan

² Department of Geology, National Science Museum, Tokyo 169-0073, Japan

³ Membership of Fossa Magna Museum, Itoigawa, Niigata 941-0056, Japan

ABSTRACT

Itoigawaite, $\text{SrAl}_2\text{Si}_2\text{O}_7(\text{OH})_2 \cdot \text{H}_2\text{O}$, is a new member of the lawsonite group found in the Itoigawa-Ohmi district, central Japan. It is orthorhombic, $Cmcm$, $a = 6.031(2)$, $b = 8.945(2)$, $c = 13.219(4)$ Å, $Z = 4$. The strongest lines in the X-ray powder diffraction pattern are 4.68 (s) (111), 4.26 (vs) (021), 3.31 (vs) (113), 2.75 (vs) (114, 202), 2.68 (vvs) (130), 2.63 (s) (131), 2.50 (s) (220, 203, 132), 2.45 (s) (221). Electron microprobe analysis gave SiO_2 32.98, TiO_2 0.87, Al_2O_3 27.67, Fe_2O_3 0.39, MgO 0.27, CaO 0.45, SrO 27.71, H_2O (diff.) 9.66, total 100.00 wt.%, corresponding to $(\text{Sr}_{0.97}\text{Ca}_{0.03}\text{Mg}_{0.02})_{\Sigma 1.02}(\text{Al}_{1.97}\text{Ti}_{0.04}\text{Fe}_{0.02})_{\Sigma 2.03}\text{Si}_{1.99}\text{H}_{3.89}\text{O}_{10}$ on the basis of $\text{O} = 10$. It is transparent, blue in colour with vitreous lustre. The streak is white and the cleavage good in one direction. The hardness is 5–5.5. The calculated density is 3.3 g cm^{-3} . It is colourless in thin section and optically biaxial positive with $\alpha = 1.664$ (2), $\beta = 1.674$ (2), $\gamma = 1.688$ (2) and $2V(\text{calc.}) = 81^\circ$. It occurs in a thin veinlet cutting a lavender-coloured Ti-bearing jadeitite. The veinlet includes irregularly-shaped aggregates composed of minute tabular crystals of itoigawaite up to $50 \mu\text{m}$ across and minute jadeite crystals with interstitial natrolite aggregates. Itoigawaite is considered to be crystallized from fluid during later stage activity of high- P/T metamorphism. The name is for the locality.

KEYWORDS: itoigawaite, new mineral, strontium, jadeite, jadeitite, lawsonite, Renge metamorphic belt, Itoigawa-Ohmi district, Japan.

Introduction

THE Itoigawa-Ohmi district in the most eastern part of the Renge belt (Nishimura, 1998) is where the first jadeitite in Japan was found (Kawano, 1939; Ohmori, 1939).

In 1996 a bluish mineral on a lavender-coloured jadeitite (Fig. 1) was collected from the seashore of the Itoigawa-Ohmi district, Niigata Prefecture, Japan. It was submitted to the first author for identification by the fourth author. A preliminary microprobe analysis showed it to be a Sr-Al silicate possibly including elements with atomic numbers < 9 . Subsequent X-ray powder study proved the diffraction pattern of the mineral to be very close to that of the lawsonite group of minerals. The mineral data and the name have been approved by the Commission on New Mineral and Mineral Names of the

International Mineralogical Association. The type specimens of itoigawaite are deposited at the National Science Museum, Tokyo, under the registered number NSM M-27872 and at Fossa Magna Museum, Itoigawa, Niigata, under the catalogue no. FMM00662.

Occurrence

The Itoigawa-Ohmi district of the Renge belt is characterized by the serpentinite melange with high- P/T type schists and various fragments of Palaeozoic accretionary complexes composed of greenstone, limestone, chert and mudstone, etc. (Nakajima *et al.*, 1992). Although no outcrops of jadeitite had been found to date, jadeitite is considered to be closely associated with serpentinite melange. Boulders and pebbles of jadeitite have been recognized in several riverbeds,




FIG. 1. Itoigawaite on jadeitite from the Itoigawa-Ohmi district. I = itoigawaite, J = jadeite.

including the River Himekawa running through the serpentine melange zone, and on the seashore of the Itoigawa-Ohmi district where they spread towards the west from the mouth of the River Himekawa, Itoigawa City. The itoigawaite-bearing jadeitite was collected on the seashore of Oyashirazu, situated ~15 km WSW of Itoigawa Station (Fig. 2).

Three specimens of itoigawaite are recognized. Type itoigawaite occurs as a thin veinlet, <0.8 mm wide, cutting the jadeitite composed essentially of lavender-coloured and colourless jadeites with minor constituents of Sr-bearing pectolite, rutile, titanite and an undetermined Sr-Ti silicate. The veinlet includes outer fillings of long prismatic jadeite crystals which are elongated perpendicular to the wall of the veinlet, and inner fillings of irregularly-shaped aggregates composed of minute tabular crystals of itoigawaite, up to ~50 μm across, with interstitial natrolite (Fig. 3). Jadeite is absent from the inner part of the veinlet. The specimen is characterized by the existence of Sr-rich minerals and euhedral jadeite crystals in the natrolite matrix (Fig. 4).

Physical and optical properties

Itoigawaite is blue with vitreous lustre, and is colourless in thin section. The cleavage is good in one direction. Fluorescence is not observed under

short wavelength ultraviolet light. The Vickers microhardness is 550–680 kg mm^{-2} (100 g load). It corresponds to 5–5.5 on Mohs' hardness scale. The density could not be measured because of the close association with natrolite. The calculated density is 3.37 g cm^{-3} . It is optically biaxial positive with refractive indices $\alpha = 1.664(2)$, $\beta = 1.593(2)$, $\gamma = 1.688(2)$, and $2V(\text{calc.}) = 81^\circ$.

Chemistry

Chemical analyses were carried out using an energy dispersive X-ray spectrometer, Link QX 2000 (Table 1). The operating conditions included a beam diameter of 2–3 μm , 15 kV and 10 nA probe current, and standard ZAF correction. The standards used were albite (for Na), Mg_2SiO_4 (for Mg), sillimanite (for Al), wollastonite (for Si and Ca), orthoclase (adularia) (for K), TiO_2 (for Ti), Cr_2O_3 (for Cr), tephroite (for Mn), Fe_2SiO_4 (for Fe) and celestine (for Sr). Chemical analyses and BEI observations show no compositional variation within single grains or between adjacent grains. X-ray diffraction patterns of the other lawsonite group minerals indicate that the difference between them and itoigawaite is water content. The water content of itoigawaite could not be measured because of the small quantity of sample available for analysis. The close association with natrolite adds to this

ITOIGAWAITE, A NEW LAWSONITE GROUP MINERAL




FIG. 2. Index map of the Itoigawa-Ohmi district, Niigata Prefecture, central Japan.

TABLE 1. Chemical analyses of itoigawaite

	1	2	3	4	5	6	7	Average
SiO ₂	33.13	33.08	33.15	33.00	33.19	32.55	32.74	32.98
TiO ₂	n.d.*	n.d.	n.d.	n.d.	0.89	0.86	n.d.	0.25
Al ₂ O ₃	28.17	27.83	27.88	27.73	27.23	26.84	28.00	27.67
FeO	n.d.	n.d.	n.d.	n.d.	0.37	0.33	n.d.	0.10
MgO	n.d.	n.d.	n.d.	0.27	0.27	0.26	n.d.	0.11
CaO	0.26	0.46	0.60	0.26	0.43	0.55	0.65	0.46
SrO	28.32	27.77	27.29	28.04	27.54	27.61	27.41	27.71
H ₂ O**	10.12	10.86	11.08	10.70	10.08	11.00	11.20	10.72
Total	100	100	100	100	100	100	100	100
Formulae on the basis of O = 8								
Si	1.997	2.006	2.008	2.001	1.996	1.986	1.991	1.998
Ti					0.040	0.039		0.011
Al	2.000	1.989	1.990	1.981	1.931	1.930	2.006	1.975
Fe					0.019	0.017		0.005
Mg				0.024	0.025	0.024		0.010
Ca	0.016	0.030	0.039	0.017	0.028	0.036	0.042	0.300
Sr	0.990	0.976	0.959	0.986	0.961	0.977	0.966	0.973
Total	5.003	5.000	4.996	5.009	4.998	5.009	5.006	5.003

* not detected

** calculated by difference




FIG. 3. (a) Photomicrograph (PPL) and (b) back-scattered electron image of the veinlet including itoigawaite. I = itoigawaite, N = natrolite, J = jadeite, P = Sr-bearing pectolite.

difficulty. Thus, the H₂O content of itoigawaite was estimated by difference. The average of seven analyses lead to the empirical formula, (Sr_{0.97}Ca_{0.03}Mg_{0.02}) Σ 1.02(Al_{1.97}Ti_{0.04}Fe_{0.02}) Σ 2.03Si_{1.99}H_{3.89}O₁₀ which gives the ideal formula SrAl₂Si₂O₇(OH)₂·H₂O.

X-ray study

Because of the minute grainsize of the material studied, the X-ray powder diffraction pattern for itoigawaite was obtained using a Gandolfi camera, 114.6 mm in diameter, employing Ni-filtered Cu-K α ₁ radiation. Single crystal studies could not be carried out due to the small crystal size. As shown in Table 2, the notable coincidence of the patterns suggests that itoigawaite is isostructural with lawsonite and hennomartinite. The refined unit cell parameters are; $a = 6.031(2)$, $b = 8.945(2)$,

$c = 13.219(4)$ Å. Therefore, the proposed space group is *Cmcm*, given the comparison with the lawsonite group.

Discussion

Itoigawaite is the Sr-Al member of the lawsonite group of sorosilicates of general formula A²⁺B³⁺Si₂O₇(OH)₂·H₂O. Features of the four known members of the group are summarized in Table 3. The cell parameters and density of itoigawaite are larger than those of lawsonite and are smaller than those of hennomartinite. These are attributed to the difference in the ionic radius and atomic weight between Sr and Ca, and Al and Mn, respectively.

The chemical composition for slawsonite, SrAl₂Si₂O₈, is close to that for itoigawaite. It is monoclinic, space group *P2*₁/*a*, in the specimen

ITOIGAWAITE, A NEW LAWSONITE GROUP MINERAL



FIG. 4. Euhedral crystals of jadeite in natrolite (PPL). J = jadeite, N = natrolite.

from Oregon, USA, corresponding to the Sr analogue of paracelsian (Griffen *et al.*, 1977), and is triclinic, space group $P1$, in that from Sarusaka, Japan (Tagai *et al.*, 1995). Although the ideal formula of itoigawaite is equivalent to that of slawsonite with two additional water molecules, the density of itoigawaite, 3.37 g cm^{-3} , is greater than that of slawsonite, 3.10 g cm^{-3} . The same relationship can be observed between lawsonite, 3.09 g cm^{-3} , and anorthite, 2.76 g cm^{-3} . The coexistence with jadeite, as well as the difference in density between itoigawaite and slawsonite, indicates that itoigawaite has a higher pressure paragenesis than slawsonite.

Itoigawaite occurs in a veinlet, cutting jadeitite (Fig. 2). The vein-filling constituents are prismatic jadeite crystals along the vein wall, with

aggregates of itoigawaite and natrolite in the inner part. Thus, it is concluded that these minerals were crystallized from fluid intruding into a fracture in the jadeitite. The concentration of Sr in the fluid increased gradually, because the ionic radius of Sr is too large to be accepted in the $M2$ site of jadeite. In the latter stage, Sr was saturated, then itoigawaite and natrolite crystallized from the residual liquid. The absence of jadeite in the inner part of veinlet indicates that jadeite could not have crystallized at the latter stage. This may be attributed to the lowering of pressure at the latter stage and/or increasing activities of silica and water. Natrolite in association with itoigawaite is considered to have crystallized from the fluid instead of jadeite.

Kobayashi *et al.* (1987) described the occurrence of stronalsite in jadeitite from Mt Ohsa, Okayama, the western range of the Renge belt (Nishimura, 1998). This is the only description of Sr-rich minerals in jadeitite. During mineralogical and petrological research of jadeitites and omphacitites from the Itoigawa-Ohmi district, however, many species of Sr-rich minerals such as tausonite, Sr-apatite, itoigawaite, stronalsite, lamprophyllite, slawsonite, and the Sr-Zr analogue of perrierite etc. have been found (Miyajima *et al.*, 1998*a,b,c,d*). The discovery of these Sr-rich minerals is important in understanding the formation mechanism of jadeitite.

Acknowledgements

The authors thank Dr K. Yokoyama of the Department of Geology, National Science Museum, Tokyo for his with the microprobe measurements. The first author is grateful to emeritus Professor K. Aoki of Tohoku University, Dr K. Kunugiza of Toyama University, Dr T. Oba of Joetsu University of Education, and to Mr K. Takenouchi of the Fossa Magna Museum, for helpful discussions and encouragement. The first author is also grateful to Mr K. Hirokawa and Mr T. Magishi for information about itoigawaite-bearing jadeitites. He would also like to thank Mr and Mrs Masuoka for their hospitality during the time spent in Tokyo.

References

- Armbruster, T., Oberhänsli, R., Bermanec, V. and Dixon, R. (1993) Hennomartinite and kornite, two new Mn^{3+} rich silicates from the Wessels Mine,

TABLE 2. X-ray powder diffraction data for itoigawaite, lawsonite and hennomartinite

<i>h k l</i>	Itoigawaite*			Lawsonite**		Hennomartinite***	
	<i>d</i> _{obs.}	<i>d</i> _{calc.}	<i>I</i>	<i>d</i>	<i>I</i>	<i>d</i>	<i>I</i>
0 0 2	6.60	6.61	vw	6.58	35		
1 1 0				4.84	60	5.149	30
1 1 1	4.68	4.68	s	4.57	30	4.804	86
0 2 1	4.26	4.24	vs	4.18	40	4.287	56
1 1 2	4.00	3.99	w	3.91	5		
0 2 2	3.70	3.70	m	3.66	60	3.725	26
1 1 3	3.31	3.31	vs	3.26	35	3.373	66
0 0 4	3.30	3.30	m			3.347	22
0 2 3	3.15	3.14	w				
2 0 0				2.925	45		
1 1 4		{ 2.76		2.726	70	2.807	82
2 0 2	2.75	{ 2.74	vs	2.680	50	2.833	100
1 3 0	2.68	2.67	vvs	2.624	100	2.715	58
0 2 4						2.695	98
1 3 1	2.63	2.62	s	2.573	5	2.666	24
2 2 0		{ 2.50					
2 0 3	2.50	{ 2.49	s				
1 3 2		{ 2.48		2.433	40	2.518	48
2 2 1	2.45	2.46	s	2.395	30		
2 2 2		{ 2.34		2.284	20	2.401	68
1 1 5	2.33	{ 2.34	m	2.311	15	2.379	26
0 2 5		{ 2.28		2.254	35		
0 4 0	2.27	{ 2.24	m	2.190	15		
0 0 6		{ 2.20					
0 4 1	2.21	{ 2.20	m				
2 2 3	2.17	2.17	s	2.129	60	2.231	54
0 4 2				2.089	5		
1 3 4	2.09	2.08	vw	2.053	5	2.113	20
1 1 6				1.990	5		
0 4 3		{ 1.994		1.965	30	2.018	20
2 2 4	1.993	{ 1.994	s				
3 1 1				1.884	15	2.009	20
1 3 5				1.852	5		
2 2 5				1.786	5		
2 0 6		{ 1.779		1.752	35	1.819	40
1 1 7	1.767	{ 1.767	s			1.795	14
0 2 7	1.743	1.740	m	1.726	20		
2 4 2				1.697	15		
0 4 5				1.685	5		
3 1 4	1.687	1.687	w	1.657	5		
2 4 3		{ 1.663					
1 5 2	1.662	{ 1.660	s	1.631	30		
2 2 6		{ 1.653		1.626	40	1.687	38
3 3 2				1.575	10		
0 4 6	1.568	1.569	s	1.550	80		
1 5 4				1.501	10		
0 6 0	1.490	1.491	w	1.464	40		

* Itoigawaite: $a = 6.031(2)$, $b = 8.945(2)$, $c = 13.219(4)$ Å (present study)** Lawsonite: $a = 5.846 \pm 0.006$, $b = 8.788 \pm 0.008$, $c = 13.129 \pm 0.010$ Å (Davis and Pabst, 1960)*** Hennomartinite: $a = 6.255(1)$, $b = 9.034(1)$, $c = 12.397(2)$ Å (Armbruster *et al.*, 1993)

ITOIGAWAITE, A NEW LAWSONITE GROUP MINERAL

 TABLE 3. The comparison of crystallographic, physical and optical data for the lawsonite group minerals, $A^{2+}B_2^{3+}Si_2O_7(OH)_2 \cdot H_2O$. Standard deviations given in parentheses

A,B in formula	Itoigawaite*	Lawsonite**	Hennomartinite [†]	Noélbensonite ^{††}
Crystal system	Sr, Al orthorhombic	Ca, Al orthorhombic	Sr, Mn orthorhombic	Ba, Mn orthorhombic
Space group	(<i>Cmcm</i>)	(<i>Cmcm</i>)	(<i>Cmcm</i>)	(<i>Cmcm</i>)
Cell parameters				
<i>a</i> , Å	6.031(2)	5.846	6.255(1)	6.325(1)
<i>b</i> , Å	8.945(2)	8.788	9.034(1)	9.120(1)
<i>c</i> , Å	13.219(4)	13.129	13.397(2)	13.618(1)
<i>V</i> , Å ³	713.1	674.5	757.03	785.5
Z	4	4	4	4
Colour	blue	colourless–pale blue	yellow–brown	dark brown
Hardness (Mohs)	5–5.5	6	~4	~4
<i>D</i> _{calc.} , g cm ⁻³	3.37	3.08–3.09	3.68	3.87
Colour in thin section	colourless	colourless	yellow–brown	orange yellow– brownish orange
Refractive indices				
α	1.664(2)	1.663–1.655	n.d.	1.82
β	1.674(2)	1.672–1.675	n.d. (n ³ , 1.82)	1.835
γ	1.668(2)	1.682–1.686	n.d.	1.85
2 <i>V</i> , °	81 (calc.)	76–87	63	46

* Present study

**Davis and Pabst (1960)

† Armbruster *et al.* (1993)

†† Kawachi *et al.* (1996)

- Kalahari, South Africa. *Schweiz. Mineral. Petrogr. Mitt.*, **73**, 349–55.
- Davis, G.A. and Pabst, A. (1960) Lawsonite and pumpellyite in glaucophane schist, North Berkeley Hills, California, with notes on the X-ray crystallography of lawsonite. *Amer. J. Sci.*, **258**, 689–704.
- Griffen, D.T., Ribbe, P.H. and Gibbs, G.V. (1977) The structure of lawsonite, a strontium analogue of paracelsian. *Amer. Mineral.*, **62**, 31–5.
- Kawachi, Y., Coombs, D.S. and Miura, H. (1996) Noélbensonite, a new BaMn silicate of the lawsonite structure type, from Woods mine, New South Wales, Australia. *Mineral. Mag.*, **60**, 369–74.
- Kawano, Y. (1939) A new occurrence of jade (jadeite) in Japan and its chemical properties. *J. Japan. Assoc. Mineral. Petrol. Econ. Geol.*, **22**, 195–201. (in Japanese).
- Kobayashi, S., Miyake, H. and Shoji, T. (1987) A jadeite rock from Oosa-cho, Okayama Prefecture, Southwestern Japan. *Mineral. J.*, **13**, 314–27.
- Miyajima, H., Matsubara, S., Miyawaki, R. and Hirokawa, K. (1998a) Strontiochevkinite and tausonite in blue jade from the Itoigawa district, the Hida marginal belt, central Japan. *Abstracts of 1998 Japan Earth and Planetary Science Joint Meeting*, p. 247. (in Japanese).
- Miyajima, H., Matsubara, S., Miyawaki, R. and Ito, K. (1998b) Sr analogue of lawsonite in jade from the Itoigawa district. *Annual Meeting Abstracts of the Mineral. Soc. Japan*, p. 18. (in Japanese).
- Miyajima, H., Matsubara, S., Miyawaki, R. and Ito, K. (1998c) Lamprophyllite in jade from the Itoigawa district. *Annual Meeting Abstracts of the Mineral. Soc. Japan*, p. 83. (in Japanese).
- Miyajima, H., Matsubara, S., Miyawaki, R., Hirokawa, K. and Ito, K. (1998d) Strontium minerals in jade from Itoigawa-Ohmi district. *Annual Meeting Abstracts of the Japan. Assoc. Mineral. Petrol. Econ. Geol.*, p. 175. (in Japanese).
- Nakajima, T., Ishiwatari, A., Sano, S., Kunugiza, K., Okamura, M., Kano, T., Soma, T. and Hayasaka, Y. (1992), Geotraverse across the Southwest Japan Arc: an overview of tectonic setting of Southwest Japan. In *29th IGC Field Trip Guide Book, vol.5: Metamorphic belts and related plutonism in the Japanese Islands* (H. Kato and H. Noro, eds). Geol. Surv. Japan, **A25**, pp. 1–83.
- Nishimura, Y. (1998) Geotectonic subdivision and areal

H. MIYAJIMA ETAL.

- extent of the Sangun belt, Inner Zone of Southwest Japan. *J. metam. Geol.*, **16**, 129–40.
- Ohmori, K. (1939) Optical properties of jade (jadeite) newly occurred in Japan. *J. Japan. Assoc. Mineral. Petrol. Econ. Geol.*, **22**, 201–12. (in Japanese).
- Tagai, T., Hoshi, T., Suzuki, M., Kato, A. and Matsubara, S. (1995) A new modification of slawsonite, $\text{SrAl}_2\text{Si}_2\text{O}_8$: Its structure and phase transition. *Zeits. Kristallogr.*, **210**, 741–5.
- [Manuscript received 4 January 1999:
revised 19 February 1999]