

## Stability of genthelvite, $Zn_4(BeSiO_4)_3S$ : An exercise in chalcophilicity using exchange operators

DONALD M. BURT

Department of Geology, Arizona State University, Tempe, Arizona 85287-1404, U.S.A.

### ABSTRACT

Despite differences in ionic radii and chemical affinities, Be and Zn have somewhat similar crystal chemical tendencies, although they rarely occur together in minerals. The only silicate in which they do so is genthelvite, a rare accessory in alkaline to peralkaline granites, syenites, and related pegmatites and greisens. The formula of genthelvite,  $Zn_4(BeSiO_4)_3S$ , can be expressed as a mixture of phenakite, willemite, and sphalerite. This fact, plus the fact that Zn is much more chalcophile than Be, allows the prediction of a series of seven sulfide-silicate reactions that involve  $SO_{-1}$  (the "acid anhydride" of  $H_2S$ ) and the seven phases sphalerite, bromellite, zincite, phenakite, willemite, genthelvite, and quartz. When  $Al_2O_3$  and  $F_2O_{-1}$  (the "acid anhydride" of HF) are added to the model system (in the additional phases corundum, gahnite, chrysoberyl, beryl, andalusite, and topaz), the resulting quartz-saturated  $\mu_{F_2O_{-1}} - \mu_{SO_{-1}}$  diagram shows that genthelvite might be incompatible with beryl (or, at best, shares a narrow stability field with it). In addition, the rare assemblage genthelvite-topaz is stable only at low  $H_2S$  and very high HF fugacities, whereas the pegmatite assemblage gahnite-quartz sulfidizes and fluorinates to the greisen assemblage sphalerite-topaz with increases in the fugacity of both  $H_2S$  and HF. Because the order of chalcophilicity—i.e., the tendency to chalcophile behavior—is  $Zn > Fe > Mn$ , genthelvite is stable only at lower  $H_2S$  fugacities than either danalite or helvite, and it is therefore unlikely to occur in solid solution with helvite.

### INTRODUCTION

Genthelvite,  $Zn_4(BeSiO_4)_3S$ , is the rarest of the sulfosilicates in the helvite group, which also contains helvite,  $Mn_4(BeSiO_4)_3S$ , and danalite,  $Fe_4(BeSiO_4)_3S$ . Genthelvite and helvite were synthesized by Mel'nikov et al. (1971) and all three (including danalite) by Fursenko and Klyakhin (1981) and Fursenko (1982); no further experimental studies on genthelvite appear to have been published. This paper represents a first attempt to delineate some phase relations involving genthelvite using physical-chemical principles (particularly the concept of exchange operators) and natural occurrences.

Natural occurrences of genthelvite, reviewed by Haapala and Ojanpera (1972), Dunn (1976), and Zubkov et al. (1976), are mainly restricted to highly fractionated alkaline to peralkaline granites and syenites or to their associated pegmatites, although genthelvite also occurs in greisens, or rarely, skarns. Genthelvite is commonly found in solid solution with danalite, but not with helvite (Dunn, 1976). Interesting questions regarding genthelvite thus are why it is so rare compared to beryl or to other members of the helvite group, why its occurrences are essentially restricted to one rock type (alkaline granite), and what its stability has to say about the comparative crystal chemistry of Be and Zn, on the one hand, and of Zn, Fe, and Mn, on the other.

Even though Zn is less abundant in the crust than Fe

or Mn, it is relatively common in sphalerite (or gahnite, in some environments); also danalite is rarer than helvite, despite the fact that Fe is more abundant than Mn. Furthermore, Zn, Fe, and Mn are all concentrated relative to Mg by fractional crystallization. In other words, the geochemical rarity of Zn alone is insufficient to explain the rarity of genthelvite.

Although Zn (Shannon, 1976) has a divalent octahedral radius (0.74 Å) that is comparable to that of Mg (0.72), Fe (0.78), or Mn (0.83), it commonly occurs in fourfold, rather than sixfold coordination with oxygen, and in that respect (and some others) seems to have as much in common with the much smaller Be (tetrahedral radius 0.27 Å; Shannon, 1976) as with Mg, Fe, or Mn (Newmann, 1949; Plakhov, 1982; cf. Ross, 1964; Hormann, 1974; Brehler, 1969; Wedepohl, 1972; Robert and Gasperin, 1985). That is—although Zn is in sixfold coordination in its fluoride, carbonates (smithsonite and minrecordite: Garavelli et al., 1982), sulfate (zinkosite), and tungstate (sanmartinite: Dunn, 1978) and when in solid solution in most rock-forming silicates (e.g., mica, pyroxene, and amphibole)—it is in fourfold coordination (with average radius 0.60 Å; Shannon, 1976) in staurolite (e.g., Griffen, 1981) and in most of its simple compounds. Familiar examples include its sulfides (sphalerite and wurtzite), oxide (zincite), normal spinel (gahnite), and silicates (willemite, hemimorphite, hardystonite, and larsenite). Many

**TABLE 1.** Some phases in the system  $ZnO$ - $BeO$ - $SiO_2$ - $SO_{-1}$ 

Mineral	Symbol	Formula
Sphalerite	Sl	$ZnS$
Beryllium sulfide		$BeS$
Silicon disulfide		$SiS_2$
Zincite	Znc	$ZnO$
Bromellite	Bro	$BeO$
Quartz	Qtz	$SiO_2$
Willemite	Wil	$Zn_2SiO_4$
Phenakite	Phe	$Be_2SiO_4$
Genthelvite	Gen	$Zn_4(BeSiO_4)_3S$

of these have Be analogues (cf. Tables 1 and 3, also guaiate for hardystonite: Grice and Robinson, 1984).


Genthelvite itself (Ross, 1964; Hassan and Grundy, 1985) is isostructural with sodalite,  $Na_4(AlSiO_4)_3Cl$ , with Na proxied by Zn, Al by Be, and Cl by S.  $BeO_4$  tetrahedra are linked at all four corners to  $SiO_4$  tetrahedra, and Zn is coordinated to 3 O and 1 S (i.e., Zn, is in fourfold coordination here as well).

Notwithstanding their crystal-chemical similarities and tendency to be concentrated by fractional crystallization, Zn and Be rarely occur together in minerals (other than minor solid solution of, e.g., Be in willemite: Gurvich, 1965; Hahn and Eysel, 1970; Chatterjee and Ganguli, 1975). The only mineral other than genthelvite that contains both as essential constituents, according to listings in Wedepohl (1972) and Hormann (1974), is karpinskyite, which is discredited (Micheelsen and Petersen, 1971). More recently, Robinson et al. (1985) described the new calcium-beryllium-zinc phosphate hydrate ehreleite, the exception that proves the rule. In this phase, as in genthelvite, Be and Zn are both four-coordinated (Hawthorne and Grice, 1987).

The point of this discussion is that the anomalous tendency of Zn to seek four-coordination should, in anything, make genthelvite more common than helvite or danalite, in which the larger divalent Mn and Fe are "forced" into four-coordination. The correct explanation for the comparative rarity of genthelvite must be sought elsewhere, in the realm of comparative mineral stabilities.

#### STABILITY DIAGRAMS AND EXCHANGE OPERATORS

The composition of genthelvite is equivalent to a mixture of sphalerite, phenakite, and willemite, according to the following relation (cf. Burt, 1980):



This compositional relation is also shown in Figure 1, a depiction of mineral compatibilities in the reciprocal ternary system  $/Zn,Be//SiO_4,S/$ , which could also be expressed as  $Zn_2SiO_4$ - $2BeZn_{-1}$ - $S_2(SiO_4)_{-1}$  in the exchange operator notation of Burt (1974). (That is, 2Be can be




Fig. 1. Assumed mineral compatibilities in the reciprocal ternary system  $/Zn,Be//SiO_4,S/$ . For abbreviations, see Table 1. (A) The composition triangle  $Zn_2SiO_4$ - $2BeZn_{-1}$ - $S_2(SiO_4)_{-1}$ . (B) The square defined by the exchange operators  $Be_2Zn_{-2}$  (or  $2BeZn_{-1}$ ) and  $S_2(SiO_4)_{-1}$  as orthogonal vectors.

exchanged for each 2Zn and 2S for each  $SiO_4$ .) Figure 1A is a composition triangle using exchange operators as points; Figure 1B a reciprocal square based on the exchange operators as orthogonal vectors (cf. Thompson, 1982; Černý and Burt, 1984). Points outside of the reciprocal quadrilateral  $Zn_2SiO_4$ - $Be_2SiO_4$ - $BeS$ - $ZnS$  in Figure 1A are physically unattainable, although their compositions can be represented algebraically. The two representations of the quadrilateral are seen to be equivalent. (Note that in Fig. 1A, genthelvite is straightforwardly plotted from the relation  $2\text{Gen} = 7Zn_2SiO_4 + 3Be_2Zn_{-2} + S_2(SiO_4)_{-1}$ ; in Fig. 1B from the ratios  $Be:Zn = 3:4$ ;  $S_2:(SiO_4) = 1:6$ , that is, genthelvite plots  $3/7$  of the way across and  $1/7$  of the way up the square.)

This system is a subsystem of the four-component system  $ZnO$ - $BeO$ - $SiO_2$ - $SO_{-1}$ , where  $SO_{-1}$ , another exchange operator, can be thought of as the "oxide of (reduced) sulfur" or the "acid anhydride" of  $H_2S$ . The operator  $SO_{-1}$  is useful for defining the relative chalcophile-lithophile tendencies of the elements (Burt, 1974, 1979a).

Table 2 lists nine compounds in this system. As mentioned, Be and Zn have analogues (and isostructural) sul-




Fig. 2. Presumed low-pressure mineral compatibilities in the reciprocal system  $/Zn,Be//O,S/$ , showing that Zn is more chalcophile than Be.

fides (although BeS, not a mineral, is isostructural with wurtzite, not sphalerite), oxides, and silicates. (The non-mineral  $SiSi_2$  is included only for completeness; it is not isostructural with quartz.) These analogies are further emphasized in the presumed low-pressure mineral compatibilities shown on Figures 2 and 3. Figure 2 is for the system  $/Zn,Be//S,O/$ , and Figure 3, for the system  $/Zn,Be//O,SiO_4/$ . Only triangular representations are shown. Note that Figure 3 could also be described as the system ZnO-BeO- $SiO_2$  (studied experimentally by Chatterjee and Ganguli, 1975), inasmuch as the operator  $(SiO_4)O_{-2}$  can also be written as  $SiO_2$  (silica or quartz; i.e., this component has a physical counterpart). Figure 2 shows that Zn, relative to Be, prefers S to oxygen (i.e., is more chalcophile); Figure 3 shows that Zn, relative to Be, prefers  $SiO_4$  to oxygen (i.e., has a greater affinity for silica). Somewhat surprisingly, according to the data in Pankratz et al. (1984), Be has even less of an affinity for S than does Si (that is, the assemblage BeO- $SiSi_2$  is more stable than BeS- $SiO_2$ ).

For all three figures, the most basic phase is generally placed at the lower left, and the exchange operators are generally acidic (in the electronic or Lewis sense: Lewis, 1938; Burt, 1974, 1979a).  $BeZn_{-1}$  is clearly acidic, owing to the much smaller size of Be (its greater attraction for electrons), as is  $(SiO_4)O_{-2}$  (it is silica, after all).  $SO_{-1}$  is likewise acidic (it is the "acid anhydride" of  $H_2S$ ,  $S^{2-}$  is a weaker base than  $O^{2-}$ , and sulfides are more polarizable and covalent than oxides). The exception is  $S_2(SiO_4)_{-1}$  in Figure 1, which can be written as  $2SO_{-1}$  minus  $(SiO_4)O_{-2}$ ; that is, as the difference between two acidic operators; its acidity is therefore difficult to predict. It may be considered as generally basic because reactions such as the following go to the right:  $2CaS + Zn_2SiO_4 = 2ZnS + Ca_2SiO_4$ , and  $2MnS + Zn_2SiO_4 = 2ZnS + Mn_2SiO_4$ ; the exchange operators  $ZnCa_{-1}$  and  $ZnMn_{-1}$ , on the basis of ionic radii, should be acidic.

When the above configuration is chosen, a diagonal relation between mineral stabilities is usually seen, as in Figures 2 and 3, or in cases involving pure oxides (cf. Korzhinskii, 1959b; Burt, 1979a). The most basic phases




Fig. 3. Presumed low-pressure mineral compatibilities in the reciprocal system  $/Zn,Be//O,SiO_4/$ , which could also be described as the system ZnO-BeO- $SiO_2$ . The figure shows that Zn has a greater affinity for silica than Be does.

will be at the lower left, the most acidic at the top (for composition triangles) or upper right (for reciprocal systems), and they will tend to react with each other to yield diagonal tie lines (from upper left to lower right) between the stable assemblages.

Figures 1, 2, and 3 merely represent three planar sections through the system ZnO-BeO- $SiO_2$ - $SO_{-1}$  (or  $/Zn,Be,Si//O,S/$ ), which is depicted as a triangular prism on Figure 4. In order to understand the phase compatibilities, one could draw tie lines for the entire system, or else draw some planar views of it that are projected through the compositions of pure phases. The latter procedure is much easier, particularly inasmuch as there is little interest in and almost nothing is known about the parts of the prism outside the tetrahedron ZnO-BeO- $SiO_2$ -ZnS (to the right of the ruled plane BeO- $SiO_2$ -ZnS in Fig. 4). In particular, from natural occurrences, nothing can be known about sulfides in the triangular face ZnS-BeS- $SiSi_2$  on the back of the diagram.

Figures 5 and 6 show two such views, respectively projected through quartz and bromellite (and onto the planes of Figs. 2 and 3). All seven of the presumably stable mineral assemblages of interest can thus be depicted; they are the numbered triangles on Figures 5 and 6. Assemblage 7, for example, can be seen from Figure 5 to be quartz + phenakite + sphalerite + genthelvite.

#### CHEMICAL POTENTIAL DIAGRAMS

The stability of genthelvite can also be represented on schematic diagrams involving the chemical potential ( $\mu$ ) of  $SO_{-1}$  and  $SiO_2$ , that is, as a function of the reduced S and silica activities of the magma or postmagmatic fluid. Silica activity or chemical potential is of interest inasmuch as genthelvite occurs in syenites and other silica-undersaturated rocks. The diagram (Fig. 7) must remain schematic, given the lack of thermochemical data on genthelvite, although it is drawn roughly consistent with available data on the sulfides, oxides, and silicates of Be and Zn (Robie et al., 1979; Barton, 1986). The method of drawing such diagrams is given in Korzhinskii (1959a).

The main topological (and arbitrary) assumption made





Fig. 4. System ZnO-BeO-SiO<sub>2</sub>-SO<sub>-1</sub> (or /Zn,Be,Si/O,S/, written to show it is a reciprocal system) depicted as a triangular prism. The subsystem of interest, ZnO-BeO-SiO<sub>2</sub>-ZnS, is that to the left of the ruled plane.

in drawing Figure 7 is that genthelvite is not stable with zincite, that is, that the genthelvite stability field is relatively restricted. This is equivalent to assuming that the following reaction goes to the right:



Either assemblage would only occur in extremely S- and silica-depleted rocks; relevant natural occurrences appear to be lacking at Franklin and Sterling Hill, New Jersey (Pete Dunn, pers. comm., 1987) or elsewhere (although genthelvite inclusions in the Zn end-member clinopyroxene petedunnite have recently been reported by Essene and Peacor, 1987). The dashed line shows the appearance of the diagram for the case of genthelvite stable with zincite; only the invariant point numbered 4 is affected. For more silica-rich rocks, those containing phenakite or quartz, the qualitative phase relations are unaffected.

The three lines sloping gently (+1/2) to the upper right are lines of equal μ<sub>S<sub>2</sub>(SiO<sub>4</sub>)-1</sub> (sulfidation, desilication). The

TABLE 2. Mineral assemblages in the system ZnO-BeO-SiO<sub>2</sub>-SO<sub>-1</sub> that indicate μ<sub>SO<sub>-1</sub></sub>

1.	BeO + Be <sub>2</sub> SiO <sub>4</sub> + 2Zn <sub>2</sub> SiO <sub>4</sub> + SO <sub>-1</sub> = Zn <sub>4</sub> (BeSiO <sub>4</sub> ) <sub>3</sub> S
	Bro Phe Wil Gen
2.	ZnO + SO <sub>-1</sub> = ZnS
	Znc Si
3.	3Be <sub>2</sub> SiO <sub>4</sub> + 4Zn <sub>2</sub> SiO <sub>4</sub> + 2SO <sub>-1</sub> = 2Zn <sub>4</sub> (BeSiO <sub>4</sub> ) <sub>3</sub> S + SiO <sub>2</sub>
	Phe Wil Gen Qtz
4.	3BeO + 3Zn <sub>2</sub> SiO <sub>4</sub> + 3SO <sub>-1</sub> = Zn <sub>4</sub> (BeSiO <sub>4</sub> ) <sub>3</sub> S + 2ZnS
	Bro Wil Gen Si
5.	3BeO + Zn <sub>4</sub> (BeSiO <sub>4</sub> ) <sub>3</sub> S + 3SO <sub>-1</sub> = 3Be <sub>2</sub> SiO <sub>4</sub> + 4ZnS
	Bro Gen Phe Si
6.	Zn <sub>2</sub> SiO <sub>4</sub> + 2SO <sub>-1</sub> = 2ZnS + SiO <sub>2</sub>
	Wil Si Qtz
7.	2Zn <sub>4</sub> (BeSiO <sub>4</sub> ) <sub>3</sub> S + 6SO <sub>-1</sub> = 3 Be <sub>2</sub> SiO <sub>4</sub> + 8ZnS + 3SiO <sub>2</sub>
	Gen Phe Si Qtz

Note: The assemblages are listed as reactions in approximate order of increasing μ<sub>SO<sub>-1</sub></sub>, that is, in order of their appearance as the numbered points in Fig. 7.




Fig. 5. Presumed low-pressure mineral compatibilities in the system ZnO-BeO-SiO<sub>2</sub>-SO<sub>-1</sub>, projected through SiO<sub>2</sub> onto the plane of Fig. 2. Numbers identify stable four-phase assemblages that plot as invariant points in Fig. 7 and are reactions (SO<sub>-1</sub> indicators) in Table 2. For abbreviations, see Table 1.

lowest marks the formation of genthelvite from willemite + phenakite; the intermediate one, the alteration of willemite to sphalerite; and the third, the breakdown of genthelvite to sphalerite + phenakite (or at low temperatures in nature, bertrandite). Each of these intersects the vertical line of silica saturation in an invariant point involving quartz; the points are respectively numbered 3, 6, and 7.

These and the other invariant points are numbered in approximate order of increasing μ<sub>SO<sub>-1</sub></sub>. They can also be considered as reactions involving only SO<sub>-1</sub>, as in Table 2. The reactions of major interest—those occurring with increasing reduced S content in quartz-bearing alkaline granites—are again 3, 6, and 7, with 3 being the first appearance of genthelvite with quartz, 6 being the disappearance of willemite and first appearance of sphalerite, and 7 being the disappearance of genthelvite. Natural occurrences of these phases are reviewed below.

### EXTENDED SYSTEM (ADDED Al AND F)

Natural occurrences of these minerals mainly involve Na, Al, and F in addition to the components used above.




Fig. 6. Presumed low-pressure mineral compatibilities in the system ZnO-BeO-SiO<sub>2</sub>-SO<sub>-1</sub>, projected through BeO (bromellite) onto the plane of Fig. 3. Numbers identify stable four-phase assemblages that plot as invariant points in Fig. 7 and as reactions (SO<sub>-1</sub> indicators) in Table 2. For abbreviations, see Table 1.




Fig. 7. Schematic isobaric, isothermal  $\mu_{SO_{-1}}-\mu_{SiO_2}$  diagram for the system  $ZnO-BeO-SiO_2-SO_{-1}$ , drawn at moderate temperatures and pressures, showing the stability of genthelvite in silica-saturated (vertical line to right) and undersaturated environments. For abbreviations, see Table 1.

To keep things relatively simple Na is omitted from phases such as nepheline, albite, villiaumite, cryolite, chiolite, chkalovite, and possibly its zinc analogue (cf. Burt, 1979b; Burt and London, 1982), and genthelvite is simply noted as being stable approximately in the field of phenakite (i.e., conditions that are alkaline or F-rich or both so that Al is tied up in feldspar or topaz: Burt, 1981) on acidity-salinity or other diagrams involving Na (or K) and F.  $H_2O$  is also omitted in low-temperature phases such as bertrandite, euclase, eudidymite, epidymite, and many others. This approach leaves Al and F as variables.

Some additional phases introduced with the addition of  $Al_2O_3$  and  $F_2O_{-1}$  (again, the "oxide" of F or the "acid





Fig. 8. Assumed mineral compatibilities in the system  $BeO-Al_2O_3-SiO_2$  at moderate temperatures and pressures. For abbreviations, see Tables 1 and 3.

anhydride" of HF) are listed on Table 3. Of these,  $AlF_3$  is not a mineral, although ralstonite in nature is not too far away from it in composition. Note that although  $ZnAl_2O_4$  and  $BeAl_2O_4$  have analogous formulas, the former is a normal spinel with Zn in four-coordination and the latter has the olivine ( $Mg_2SiO_4$ ) structure with Al in the Mg sites and Be in the Si site (cf. Ross, 1964).

Mineral compatibilities in the system  $BeO-Al_2O_3-SiO_2$  (Fig. 8) were studied on the basis of natural assemblages by Burt (1978) and experimentally by Barton (1986). The main area of disagreement involved chrysoberyl, which very rarely occurs in quartz-bearing pegmatites (commonly those that also contain sillimanite), and, according to the experimental work, is stable with quartz only above 600 °C. This time (unlike in 1978) I have therefore shown beryl + andalusite as stable (with a dashed line), although in nature the Al is usually present in feldspars, topaz, or mica rather than andalusite.

The system  $ZnO-Al_2O_3-SiO_2$  (Fig. 9) is much simpler; the common occurrence of gahnite in quartz-bearing pegmatites rules out any ambiguity (cf. Bunting, 1932), omitting, of course, Zn pyroxenes (Morimoto et al., 1975; Olesch et al., 1982) or other phases that might appear at high pressure.

Projecting through quartz, these data can be combined as shown in Figure 10 for the system  $ZnO-BeO-Al_2O_3-SiO_2$ . Chrysoberyl would appear on this figure only at high temperatures (Barton, 1986). Gahnite plus quartz is assumed stable with all of the Be minerals; the assemblage gahnite + beryl is commonly seen in pegmatites (Němec, 1973; Černý and Hawthorne, 1982), whereas the assemblage gahnite + phenakite is based on the assumption that the following reaction goes to the right:



This assumption seems reasonable based on the rareness of willemite, commonness of gahnite, and the discussion that accompanies Figure 11.

TABLE 3. Phases added with  $Al_2O_3$  and  $F_2O_{-1}$

Mineral	Symbol	Formula
Aluminum fluoride		$AlF_3$
Corundum	Cor	$Al_2O_3$
Gahnite	Gah	$ZnAl_2O_4$
Chrysoberyl	Chr	$BeAl_2O_4$
Beryl	Brl	$Be_3Al_2Si_6O_{18}$
Andalusite, etc.	Als	$Al_2SiO_5$
Topaz	Tpz	$Al_2SiO_4F_2$





Fig. 9. Assumed mineral compatibilities in the system ZnO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub> at moderate temperatures and pressures. For abbreviations, see Tables 1 and 3.

It remains to sulfidize and fluorinate the system of Figure 10 to create assemblages containing genthelvite, sphalerite, and topaz. This is done on the  $\mu_{\text{SO}_-1} - \mu_{\text{F}_2\text{O}_-1}$  diagram (Fig. 11). It was assumed in drawing the figure that the following reaction goes to the right:



This assumption is consistent with keeping the genthelvite field relatively restricted, and it leads to a simpler configuration of the diagram. An "association" (not necessarily assemblage) between impure genthelvite and beryl is reported only by Vasil'ev (1961) and implied by Chistyakova and Moleva (1966; cf. Chistyakova, 1968) and Haapala (1977, p. 68; in a 1988 personal communication he reports never finding the two phases touching); no other associations are reported. However, assemblages between the much more widespread helvite-danalite series minerals and beryl are reported fairly commonly (e.g., DeMark, 1981; Gallegher and Hawkes, 1966; Orlov et al., 1961; Kalenov, 1959), and so my assumption may be wrong.

If so, the alternative configuration of the multisystem (see Korzhinskii, 1959a, for a definition of multisystems) would be stable, as shown in configuration B to the lower left of Figure 12. The multisystem, as represented by Reaction 4 above, is made up of the phases Gen, Brl; Sl, Gah, Phe, Qtz. By the stable-to-metastable topological relation of Burt (1971, 1978), the invariant points that are stable are labeled by the phase assemblage that is metastable in each of the two alternative configurations of the multisystem. Inasmuch as Figure 11 assumes that the assemblage Gen + Brl is metastable, these two "absent" (or nonparticipating) phases can be chosen to label the two stable invariant points 3 and 4. This is done to the upper right (A) of Figure 12, with the invariant-point labels placed in brackets, [Gen] (above) and [Brl] (below). In the alternative configuration (B) to the lower left, the assemblage Sl + Gah + Phe + Qtz is metastable, and





Fig. 10. Assumed mineral compatibilities in the system ZnO-BeO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>, projected through quartz, at moderate temperatures and pressures. For abbreviations, see Tables 1 and 3.

the points bearing those labels become stable. I am projecting through Qtz, and therefore its point cannot be seen, but the other three points are seen: [Sl] at the bottom, [Phe] nearly at the top, and [Gah] at the top. The two points [Brl] and [Gen] that were stable in configuration A are now metastable, and they have switched places, as shown by their labels and the dashed, now metastable, reaction lines. The assemblage genthelvite + beryl is now stable over a limited area of the diagram (in the half-open rectangle closed to the right by the invariant point labeled [Gah] above and [Sl] below).

The SO<sub>-1</sub> indicators to the left of invariant points 1, 2, and 3 of Figure 11 correspond to 3, 6, and 7 on Table 2, respectively. Two new SO<sub>-1</sub> indicators appear, to the left of points 4 and 5. The higher is gahnite + quartz going to sphalerite + andalusite, the lower is gahnite + phenakite going to sphalerite + beryl. The two F<sub>2</sub>O<sub>-1</sub> indicators that are the vertical lines beneath points 5 and 4 have been reported elsewhere (Burt, 1977, 1981), the first is the conversion of andalusite to topaz, the second the breakdown of beryl to topaz, phenakite, and quartz. A still higher F<sub>2</sub>O<sub>-1</sub> indicator occurs to the right, beneath point 2 and passing through 1; it is



No natural examples of the right-hand assemblage are reported, although the association willemite-topaz is implied for alkaline metasomatites and greisens in Table 23 of Zubkov et al. (1976, p. 92-93). One reason might be that willemite-bearing rocks are commonly peralkaline (see below), in which case F occurs mainly as sodium fluorides, rather than in topaz (Burt and London, 1982). Thus Zubkov et al. (1976, Table 23, p. 92-93) imply a strong aluminofluoride-willemite association in mica-quartz-feldspar metasomatites and greisens.

A second reason for the lack of topaz + willemite assemblages might be that they are unstable with respect to end-member AlF<sub>3</sub> (instead of just Na-Al fluorides). This





Fig. 11. Schematic isobaric, isothermal  $\mu_{SO_2}$ - $\mu_{F_2O_3}$  diagram for the system  $ZnO$ - $BeO$ - $Al_2O_3$ - $SiO_2$ - $SO_2$ - $F_2O_3$ , in the presence of quartz, at moderate temperatures and pressures. Genthelvite is assumed to be unstable with beryl. The five invariant points are numbered in order of increasing  $\mu_{SO_2}$  and decreasing  $\mu_{F_2O_3}$ , or  $\log a_{HF}^2/a_{H_2S}$ . Inset shows chemography of phases (see Tables 1 and 3 for abbreviations).


phase is shown as forming from topaz along the vertical line to the far right of Figure 11. If the  $AlF_3$ -forming reaction is moved sufficiently far to the left, Reaction 5 above becomes metastable, owing to the multisystem relation



Inasmuch as neither the left- nor right-hand side of this relation is definitely known ( $AlF_3$  is not reported as a mineral), perhaps this would be a good point to stop drawing more phase diagrams. However, the wide gap between the breakdown of albite to cryolite + topaz + quartz and the breakdown of topaz to  $AlF_3$  + quartz, as a function of increasing  $\mu_{F_2O_3}$  (Anovitz et al., 1987, Fig. 5, p. 3098), especially with falling temperature, suggests that this reaction goes to the right, as assumed in drawing Figure 11. The albite breakdown to cryolite, topaz, and quartz (Burt, 1979b) would then be a vertical line (not

shown) between the breakdown of beryl (passing through point 4) and that of gahnite plus quartz (passing through points 1 and 2).

A third and more probable reason is that HF-rich natural environments such as greisens are normally also enriched in  $H_2S$ , so that the typical reaction terminal to gahnite is the diagonal one (with a  $-1$  slope) that joins points 2 and 5, namely



The left-hand assemblage is typical of pegmatites; the right-hand one of greisens. The rare greisen assemblage genthelvite + topaz (Haapala and Ojanpera, 1972; Haapala, 1977) is stable to the right of the line joining point 1 to point 3, again, very F rich, low S conditions.

Subtracting  $SO_{-1}$  from  $F_2O_{-1}$  yields  $F_2S_{-1}$ . The chemical potential of this component (or the  $\log a_{HF}^2/a_{H_2S}$ , for an activity ratio in the presence of  $H_2O$ ) increases to the lower right of Figures 11 and 12, as indicated by the vectors to the upper left of Figure 12. If specific mineralogic indicators of this tendency (toward F-rich, S-poor) are desired, the five numbered invariant points of Figure 11 could serve as guides.

**NATURAL ASSOCIATIONS**

As mentioned above, genthelvite is the rarest mineral of the helvite group, and genthelvite occurrences are mainly restricted to alkaline granites, pegmatites, syenites, and related metasomatic rocks (Gurvich et al., 1965; Galetskii et al., 1970; Haapala and Ojanpera, 1972; Dunn, 1976). Common associates are quartz, feldspars, micas, other Zn-bearing phases such as sphalerite (Haapala, 1977; Bowden and Kinnaird, 1978), willemite (Gurvich, 1965; Galetskii, 1966, 1971), and gahnite (Gurvich, 1965; Haapala, 1977), as well as other Be-bearing phases, mainly phenakite (Gurvich, 1965; Gurov and Marchenko, 1970), but also bertrandite (Chistyakova and Moleva, 1966).

The common occurrence of willemite in these associations, particularly those from peralkaline rocks, is one that was not known until relatively recently (e.g., Pough, 1941, did not mention them). More recently, Metcalf-Johnson (1977) reported willemite from the Ilimaussaq complex with chkalovite as the only accompanying Be mineral; genthelvite occurs elsewhere in the complex (Bollinberg and Petersen, 1967). Similarly, Bank (1975) reported willemite from Mont St. Hilaire in Canada; Dunn (1976) reported genthelvite. In such peralalcaline willemite-bearing associations it would be unlikely to find topaz; rather sodium fluorides are typical. For example Kudrin (1978) reported peralkaline metasomatic rocks from the USSR with willemite, phenakite, and genthelvite; fluorine occurs in sodium fluorides such as pachtolite,




Fig. 12. Two alternative configurations for the multisystem derived from Reaction 4. Configuration A, to the upper right, is that of Fig. 11, in which the assemblage Gen + Brl is unstable. Configuration B, to the lower left, assumes that the assemblage Si + Gah + Phe + Qtz is unstable. See text for discussion; for abbreviations, see Tables 1 and 3.

thompsenolite, ralstonite, weberite, neighborite, and gargarinite.

Nigerite (Jacobson and Webb, 1947) is another accessory Zn mineral of granites and pegmatites, but no occurrences with genthelvite are reported; common associates are sillimanite, chrysoberyl, and gahnite (cf. Kloosterman, 1974; Burke et al., 1977; Cech et al., 1978). These data suggest that nigerite is restricted to peraluminous, high-temperature environments. As mentioned and discussed above, occurrences of genthelvite with aluminous minerals such as beryl and topaz seem to be quite restricted.


**SOLID SOLUTION RELATIONS**

Despite their somewhat similar crystal chemistry, Zn is very much more chalcophile than Be. In fact, Be is not




chalcophile at all (probably less so than Si), but Zn is more chalcophile than Fe, and much more so than Mn (e.g., Marakushev and Bezmen, 1974; Maurel, 1969). This fact presumably explains (Burt, 1977) the otherwise inexplicable lack of Zn-Mn solid solutions in the Zn-Fe-Mn helvite-group minerals noted by Dunn (1976). That is, under environments rich enough in  $SO_{-1}$  [i.e., under sufficiently high  $H_2S$  fugacities; actually, the operator involved is  $S_2(SiO_4)_{-1}$ ] that the FeS or especially MnS component of helvite would be stable, the  $Zn_2SiO_4$  component would have been sulfidized to ZnS. Conversely, under the low  $SO_{-1}$  conditions needed to stabilize genthelvite, the FeS and particularly MnS components of danalite and helvite would not yet be stable, and Fe and particularly Mn, if present, would occur in coexisting silicates or oxides.

In terms of specific reactions, data given by Robie et al. (1979) as well as common natural assemblages confirm that the following reaction goes to the right among the end members:



Presumably also the following end-member reaction would go to the right:



or, if not tephroite, some other manganese silicate.

### SUMMARY

The extremely high chalcophilicity of Zn explains, in part, why genthelvite is such a rare mineral. It is only somewhat more stable than willemite and is therefore restricted to the extremely low-S environments provided by alkaline granites and syenites (where it typically occurs with other Zn minerals such as willemite and gahnite). These alkali-rich environments also tie up all available Al as feldspars or feldspathoids; this tendency destabilizes beryl and allows the occurrence of phenakite and of members of the helvite group (Burt, 1981). It is unclear (Reaction 4 above) whether beryl is stable with genthelvite at all, although beryl is reported with other minerals of the helvite group. Under strongly peralalaline (agpaitic) conditions, genthelvite or phenakite are replaced by sodium-beryllium silicates such as chkalovite and possibly a zinc analogue (synthesized by Litvin et al., 1974) or by hydrous phases such as eudidymite and epididymite.

Figure 7 shows that genthelvite becomes stable at a lowered  $SO_{-1}$  potential under conditions of silica activity less than one; conversely, a raised  $SO_{-1}$  potential destroys genthelvite more readily under these conditions. That is, genthelvite will be the most resistant to sulfidation ("burning off") in a silica-saturated environment; with rising  $H_2S$  it will break down to a mixture of sphalerite,

phenakite, and quartz (an assemblage that could coexist with a danalite-enriched member of the helvite group).

Assemblages of sphalerite with aluminum silicates (or pyrophyllite) indicate the instability of gahnite under relatively high  $H_2S$ , low HF conditions (as in metamorphosed sulfide deposits). Assemblages of topaz with ZnO-rich minerals such as gahnite, genthelvite, and willemite (the latter not reported with topaz, to my knowledge) indicate an environment unusually rich in HF and poor in  $H_2S$ , as indicated by the operator  $F_2S_{-1}$  or  $\log a_{HF}^2/aH_2S$ . Details are given in Figure 11. The more normal assemblage is topaz with sphalerite, especially in greisens; this assemblage indicates an environment enriched in both HF and  $H_2S$ . The topaz + sphalerite assemblage is compositionally equivalent (Reaction 7 above) to the pegmatite assemblage gahnite + quartz, which indicates an environment depleted in both HF and  $H_2S$ .

### ACKNOWLEDGMENTS

I am grateful to the Interlibrary Loan Service, Arizona State University, for helping me to obtain many obscure references in Russian, to the late D. S. Korzhinskii for inspiring me to learn Russian, and to J. B. Thompson, Jr., for suggesting that I use the exchange operator  $F_2O_{-1}$  in doing my Ph.D. thesis. Special thanks are due I. J. Haapala and P. J. Dunn for their careful studies of and for discussions about genthelvite. I also thank them and L. C. Hsu, David London, Jo Laird, and an anonymous reviewer for careful reviews of earlier versions of this paper and Deborah Barron for drafting the figures.

The work discussed above was mainly completed in 1977 (Burt, 1977); the paper was finally written and the work brought up to date while I was Visiting Scientist at the Lunar and Planetary Institute, Houston, which is operated by the Universities Space Research Association under Contract No. NASW-4066 with the National Aeronautics and Space Administration. This paper is Lunar and Planetary Institute Contribution no. 671.

### REFERENCES CITED

- Anovitz, L.M., Hemingway, B.S., Westrum, E.F., Jr., Metz, G.W., and Essene, E.J. (1987) Heat capacity measurements for cryolite ( $Na_3AlF_6$ ) and reactions in the system Na-Fe-Al-Si-O-F. *Geochimica et Cosmochimica Acta*, 51, 3087-3103.
- Bank, H. (1975) Durchsichtiger schleifwuerdiger blauer Willemit vom Mt. St. Hilaire in Kanada. *Zeitschrift der Deutsche Gemmologische Gesellschaft (Ikar-Oberstein)*, 24, 250.
- Barton, M.D. (1986) Phase equilibria and thermodynamic properties of minerals in the  $BeO-Al_2O_3-SiO_2-H_2O$  (BASH) system, with petrologic implications. *American Mineralogist*, 71, 277-300.
- Bollinberg, H., and Petersen, O.V. (1967) Genthelvite from the Ilimausaq alkaline intrusion, South Greenland. *Meddelelser om Grønland*, 181, no. 4, 1-9.
- Bowden, P., and Kinnaird, J.A. (1978) Younger granites of Nigeria—A zinc-rich tin province. *Transactions of the Institution of Mining and Metallurgy, Section B*, 87 (May), 66-69.
- Brehler, B. (1969) Crystal chemistry [of zinc]. In K.H. Wedepohl, Ed., *Handbook of geochemistry*, II/3, Chapter 30, Zinc, p. 30-A-1-30-A-10. Springer-Verlag, New York.
- Bunting, E.N. (1932) Phase equilibria in the system  $SiO_2-ZnO-Al_2O_3$ . *National Bureau of Standards Journal of Research*, 8, 279-287.
- Burke, E.A.J., Lof, P., and Hazebroek, H.P. (1977) Nigerrite from the Rosendal pegmatite and aplites, Kemio Island, southwestern Finland. *Bulletin of the Geological Society of Finland*, 49, 151-157.
- Burt, D.M. (1971) Multisystems analysis of the relative stabilities of babingtonite and ilvaite. *Carnegie Institution of Washington Year Book* 70, 189-197.
- (1974) Concepts of acidity and basicity in petrology—The exchange operator approach. *Geological Society of America Abstracts with Programs*, 6, 674-676.

- (1977) Chalcophile-lithophile tendencies in the helvite group. Genthelvitite stability in the system  $ZnO-BeO-Al_2O_3-SiO_2-SO_{-1}-F_2O_{-1}$  (abs.). EOS, 58, 1242.
- (1978) Multisystems analysis of beryllium mineral stabilities: The system  $BeO-Al_2O_3-SiO_2-H_2O$ . American Mineralogist, 63, 664–676.
- (1979a) Exchange operators, acids, and bases. In V.A. Zharikov, V.I. Fonarev, and S.P. Korikovskii, Eds., Problemy fiziko-khimicheskoi petrologii (Problems of physico-chemical petrology), vol. 2, p. 3–15. Nauka Press, Moscow (in Russian).
- (1979b) Geometrical analysis of cryolite and related mineral stabilities in the system  $Na_2O-Al_2O_3-SiO_2-F_2O_{-1}$  (abs.). EOS, 60, 965.
- (1980) The stability of danalite,  $Fe_4Be_3(SiO_4)_3S$ . American Mineralogist, 65, 355–360.
- (1981) Acidity-salinity diagrams—Application to greisen and porphyry deposits. Economic Geology, 76, 832–843.
- Burt, D.M., and London, D. (1982) Subsolidus equilibria. Mineralogical Association of Canada Short Course Handbook 8, p. 329–346.
- Cech, F., Rieder, M., Novak, F., and Novotny, J. (1978) Accessory nigerite in a granite from central Bohemia, Czechoslovakia. Neues Jahrbuch für Mineralogie Monatshefte, 331–347.
- Černý, P., and Burt, D.M. (1984) Paragenesis, crystallochemical characteristics, and geochemical evolution of micas in granitic pegmatites. Mineralogical Society of America Reviews in Mineralogy, 13, 257–297.
- Černý, P., and Hawthorne, F.C. (1982) Selected peraluminous minerals. Mineralogical Association of Canada Short Course Handbook 8, 163–186.
- Chatterjee, M., and Ganguli, D. (1975) Phase relationships in the system  $BeO-ZnO-SiO_2$ . Neues Jahrbuch für Mineralogie Monatshefte, 518–526.
- Chistyakova, M.B. (1968) Beryl and bazzite from rock crystal voids in granitic pegmatites of Kazakhstan. Trudy Mineralogicheskogo Muzeya, Akademia Nauk SSSR, 18, 140–153 (in Russian).
- Chistyakova, M.B., and Moleva, V.A. (1966) Genthelvitite from rock crystal voids in the Kentsk granitic massif (Central Kazakhstan). Trudy Mineralogicheskogo Muzeya, Akademia Nauk SSSR, 17, 240–245 (in Russian).
- DeMark, R.S. (1981) Beryl at Iron Mountain, Sierra County, New Mexico (abs.). New Mexico Geology, 3, no. 1, 15.
- Dunn, P.J. (1976) Genthelvitite and the helvite group. Mineralogical Magazine, 40, 627–636.
- (1978) Sanmartinite: New data. Mineralogical Magazine, 42, 281.
- Essene, E.J., and Peacor, D.R. (1987) Petedunnite ( $CaZnSi_2O_6$ ), a new zinc clinopyroxene from Franklin, New Jersey, and phase equilibria for zincian pyroxenes. American Mineralogist, 72, 157–166.
- Fursenko, D.A. (1982) Dependence of the composition of helvite on the acidity-basicity of mineralizing solutions. In A.A. Godovikov, Ed., Fiz.-khim. issled. mineralobrazuyushch. sistemakh, p. 104–107. Akademia Nauk SSSR, Sib. Otd., Inst. Geol. Geofiz., Novosibirsk, USSR (in Russian, not seen; abstract in Chemical Abstracts, v. 98, no. 219084n).
- Fursenko, D.A., and Klyakhin, K.A. (1981) Synthesis of minerals of the helvite group. In Yu.P. Melnik, Ed., Vses. sovesch. po eksperim. i tekhn. mineral. i petrogr. [Mater.], 10th, 1978, p. 119–126. Naukova Dumka, Kiev, Ukrainian SSR, USSR (in Russian, not seen; abstract in Chemical Abstracts, v. 98, no. 201508u).
- Galetsii, L.S. (1966) Unusual paragenesis of genthelvitite and cassiterite. Mineralogicheskii Sbornik (L'vov), 20, 46–52 (in Russian).
- (1971) Genthelvitite mineralization as a new high-grade type of beryllium raw material. Geologia Rudnykh Mestorozhdenii, 13, 21–30 (in Russian).
- Galetsii, L.S., Mel'nik, Yu.M., and Razumeeva, N.N. (1970) Formation conditions of genthelvitite. Mineralogicheskii Sbornik (L'vov), 24, 176–184 (in Russian).
- Gallegher, M.J., and Hawkes, J.R. (1966) Beryllium minerals from Rhodesia and Uganda. Bulletin of the Geological Survey of Great Britain 25, 59–75.
- Garavelli, C.G., Vurro, F., and Fioravanti, G.C. (1982) Minrecordite, a new mineral from Tsumeb. Mineralogical Record, 13, 131–136.
- Grice, J.D., and Robinson, G.W. (1984) Jeffreyite,  $(Ca,Na)_2(Be,Al)Si_2(O,OH)_7$ , a new mineral species and its relation to the melilitite group. Canadian Mineralogist, 22, 443–446.
- Griffen, D.T. (1981) Synthetic Fe/Zn staurolite and the ionic radius of  $^{IV}Zn^{2+}$ . American Mineralogist, 66, 932–935.
- Gurov, E.P., and Marchenko, E.Ya. (1970) Relation of genthelvitite and phenakite in beryllium-bearing metamatites. Geologia Rudnykh Mestorozhdenii, 12, 107–110 (in Russian).
- Gurvich, S.I. (1965) Discovery of beryllium-bearing willemite in the USSR. Doklady of the Academy of Sciences of the USSR, Earth Science Sections, 153, 136–138 (translated from Doklady Akademii Nauk SSSR, 153, 681–683, 1963).
- Gurvich, S.I., Zubkov, L.B., and Galetskiy, L.S. (1965) Geological and mineralogical characteristics of beryllium mineralization related to genthelvitite. Sovetskaya Geologiya, 8, no. 2, 29–44 (in Russian).
- Haapala, I. (1977) Petrography and geochemistry of the Eurajoki stock, a rapakivi-granite complex with greisen-type mineralization in southwestern Finland. Geological Survey of Finland Bulletin, 286, 128 p.
- Haapala, I., and Ojanpera, P. (1972) Genthelvitite-bearing greisens in southern Finland. Geological Survey of Finland Bulletin 259, 22 p.
- Hahn, Th., and Eysel, W. (1970) Solid solubility in the system  $Zn_2SiO_4-Zn_2GeO_4-Be_2SiO_4-Be_2GeO_4$ . Neues Jahrbuch für Mineralogie Monatshefte, 263–275.
- Hassan, I., and Grundy, H.D. (1985) The crystal structures of helvite group minerals,  $(Mn,Fe,Zn)_4(Be_3Si_6O_{24})S_2$ . American Mineralogist, 70, 186–192.
- Hawthorne, F.C., and Grice, J.D. (1987) The crystal structure of ehrleite, a tetrahedral sheet structure. Canadian Mineralogist, 25, 767–774.
- Hormann, P.K. (1974) Beryllium. In K.H. Wedepohl, Ed., Handbook of geochemistry, vol. II/1, Chapter 4. Springer-Verlag, New York.
- Jacobson, R.J., and Webb, J.S. (1947) The occurrence of nigerite, a new tin mineral, in quartz-sillimanite rocks from Nigeria. Mineralogical Magazine, 28, 118–128.
- Kalenov, A.D. (1959) Greisen type of helvite mineralization. Zapiski Vsesoyuznogo Mineralogicheskogo Obshchestva, 88, 481–485 (in Russian).
- Kloosterman, J.B. (1974) Nigerite in the tin-tantalum pegmatites of Amapá, Brazil. Mineralogical Magazine, 39, 837–846.
- Korzhinskii, D.S. (1959a) Physicochemical basis of the analysis of the paragenesis of minerals, 142 p. Consultants Bureau, New York (translated 1959 from 1957 book published by Nauka Press, Moscow).
- (1959b) Acid-basic interaction of components in silicate melts and the direction of the cotectic lines. Doklady of the Academy of Sciences of the USSR, Earth Science Sections (in translation), 128, 821–823.
- Kudrin, V.S. (1978) Rare metal alkaline quartz-albite-microcline metamorphites (qualmites) of zones of regional metamorphism. In D.S. Korzhinskii, Ed., Metasomatism and ore deposition, p. 183–194. Nauka Press, Moscow (in Russian).
- Lewis, G.N. (1938) Acids and bases. Journal of the Franklin Institute, 226, 293–313.
- Litvin, B.N., Tarasenkova, O.S., and Belov, N.V. (1974) Synthesis and crystallochemical properties of compounds in the system  $Na_2O-ZnO-SiO_2-H_2O$  formed at 400° to 500 °C. In V.A. Zharikov and V.N. Zyrvanov, Eds., Ocherki fiziko-khimicheskoi petrologii, 4, 89–101. Nauka Press, Moscow.
- Marakushev, A.A., and Bezmen, N.I. (1974) Chemical affinity of metals for oxygen and sulfur. International Geology Review, 13, 1781–1794 (translated from Geologia Rudnykh Mestorozhdenii, 11, no. 4, 8–23, 1969).
- Maurel, C. (1969) Étude expérimentale de la distribution de Mn, Fe, Co, Ni, Zn et Cd entre olivine et sulfure. Bulletin de la Société Française de Minéralogie et de Cristallographie, 92, 453–461.
- Mel'nikov, O.K., Litvin, B.N., and Fedosova, S.P. (1971) Production of helvite-group compounds. In A.N. Lobachev, Ed., Hydrothermal synthesis of crystals, p. 119–125. Consultants Bureau, New York (translated from Gidrottermalnykh Sintesis Kristallov, p. 167–174. Nauka Press, Moscow, 1968).
- Metcalf-Johnson, J. (1977) Willemite from the Ilmaussaq alkaline intrusion. Mineralogical Magazine, 41, 71–75.
- Micheelsen, H., and Petersen, O.V. (1971) Leifite, revised and karpinskyite, discredited (abs.). Mineralogical Society of Japan Special Paper 1, 264–265. (Proceedings of the International Mineralogical Association, 7th General Meeting, Tokyo, 1970.)
- Morimoto, N., Nakajima, Y., Syono, Y., Akimoto, S., and Matsui, Y.

- (1975) Crystal-structure of pyroxene-type  $ZnSiO_3$  and  $ZnMgSi_2O_6$ . *Acta Crystallographica*, B31, 1041-1049.
- Němec, D. (1973) Das vorkommen der Zn-spinelle in der Boehmischen Masse. *Tschermaks Mineralogische und Petrographische Mitteilungen*, 19, 95-109.
- Newman, H. (1949) Notes on the mineralogy and geochemistry of zinc. *Mineralogical Magazine*, 28, 575-581.
- Olesch, M., Doroshev, A.M., and Nechaev, P.Ju (1982) Low pressure stability of zinc clinopyroxene  $ZnSiO_3$  and the possible stable occurrence of zinc orthopyroxene. *Neues Jahrbuch für Mineralogie Monatshefte*, 312-320.
- Orlov, Yu.L., Ginzburg, A.I., and Pinevich, G.N. (1961) Paragenetic interrelation of beryllium minerals in some pegmatite veins. *Trudy Mineralogicheskogo Muzeya, Akademia Nauk SSSR*, no. 11, 103-113 (in Russian).
- Pankratz, L.B., Stuve, J.M., and Gokcen, N.A. (1984) Thermodynamic data for mineral technology. U.S. Bureau of Mines Bulletin 677, 355 p.
- Plakhov, G.F. (1982) Comparative geocrystallochemistry of beryllium and zinc. *Probl. Kristall., Moscow*, no. 3, 257-263 (in Russian) (not seen, from Chemical Abstracts 98-19858f, in turn from Referatny Zhurnal, *Geologiya*, 1982, Abs. No. 10V186).
- Pough, F.H. (1941) Occurrence of willemite. *American Mineralogist*, 26, 92-102.
- Robert, J.-L., and Gasparin, M. (1985) Crystal structure refinement of hendricksite, a Zn- and Mn-rich trioctahedral potassium mica: A contribution to the crystal chemistry of zinc-bearing minerals. *Tschermaks Mineralogische und Petrographische Mitteilungen*, 34, 1-14.
- Robie, R.A., Hemingway, B.S., and Fisher, J.R. (1979) Thermodynamic properties of minerals and related substances at 298.1 K and 1 bar ( $10^5$  pascals) pressure and at higher temperatures. U.S. Geological Survey Bulletin 1452, 456 p.
- Robinson, G.W., Grice, J.D., and Van Velthuizen, J. (1985) Ehrleite, a new calcium beryllium zinc phosphate hydrate from the Tip Top pegmatite, Custer, South Dakota. *Canadian Mineralogist*, 23, 507-510.
- Ross, M. (1964) Crystal chemistry of beryllium. U.S. Geological Survey Professional Paper 468, 30 p.
- Shannon, R.D. (1976) Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallographica*, A32, 751-767.
- Thompson, J.B., Jr. (1982) Composition space: An algebraic and geometric approach. *Mineralogical Society of America Reviews in Mineralogy*, 10, 1-31.
- Vasil'ev, V.A. (1961) On genthelvite. *Zapiski Vsesoyuznogo Mineralogicheskogo Obshchestva*, 90, 571-578 (in Russian).
- Wedepohl, K.H. (1972) Zinc. In K.H. Wedepohl, Ed., *Handbook of Geochemistry*, vol. II/3, Chapter 30-B to 30-O. Springer-Verlag, New York.
- Zubkov, L.B., Galetskii, L.S., and Metalidi, S.V. (1976) Helvite-group minerals and their deposits, 214 p. *Naukova Dumka*, Kiev, Ukrainian SSR, USSR (in Russian).

MANUSCRIPT RECEIVED FEBRUARY 3, 1988

MANUSCRIPT ACCEPTED AUGUST 1, 1988