

Chemical variation in garnets from aplites and pegmatites, peninsular Thailand—a correction

I AM grateful to Dr M. Pichavant for promptly bringing to my attention an error in Table I of Manning (1983). The averaged analyses for the garnets are erroneous, and a corrected version is presented here (Table I). The atomic proportions, 2σ and end-member proportions previously given are correct. This error, which arose during the preparation of the manuscript, in no way affects the observations or conclusions made in the paper.

REFERENCE

Manning, D. A. C. (1983) *Mineral. Mag.* **47**, 353–8.

[Manuscript received 27 October 1983]

© Copyright the Mineralogical Society

TABLE I. Average electron microprobe analyses (wt. % oxides) of garnets from aplites and pegmatites, peninsular Thailand. Values below the detection limit are denoted by —

	4012		4036		4041	4059		4063		4069		4070		4071
SiO ₂	36.00	35.19	37.07	35.61	36.68	35.92	35.66	35.95	37.07	35.98	37.26	35.83	36.01	36.30
Al ₂ O ₃	20.78	20.60	21.39	20.75	21.10	20.55	20.60	20.79	21.29	20.53	21.29	20.64	20.71	20.91
TiO ₂	—	—	0.10*	0.22*	0.11*	—	—	—	0.07*	—	0.04*	—	—	0.15*
FeO	11.40	10.55	11.09	21.62	18.34	21.11	19.83	25.34	25.06	23.09	22.17	20.00	20.89	19.07
MnO	31.55	33.27	31.97	20.94	23.40	21.95	23.13	16.65	15.56	19.79	20.52	22.71	21.73	22.66
MgO	—	—	0.04†	0.22†	0.08†	—	—	0.23*	0.47	0.46	0.26	0.22	0.27*	0.12†
CaO	0.36	0.38	0.38	0.75	0.80	0.67	0.52	0.66	0.69	0.29	0.34	0.25	0.24	0.33
Total	100.09	99.99	102.04	100.11	100.51	100.20	99.97	99.86	100.20	99.94	101.84	99.70	99.58	99.54
n	20	12	24	32	56	22	27	20	26	18	26	9	20	16
	*n = 13	*n = 9	*n = 44				*n = 9		*n = 14		*n = 11	*n = 3		*n = 12
	†n = 12	†n = 10	†n = 47											†n = 15

Department of Geology, The University,
Newcastle upon Tyne NE1 7RU

D. A. C. MANNING