

## Thermochemistry and phase equilibria in calcium zeolites

IRINA KISELEVA,<sup>1</sup> ALEXANDRA NAVROTSKY,<sup>1</sup> IGOR A. BELITSKY,<sup>2</sup> AND BORIS A. FURSENKO<sup>2</sup>

<sup>1</sup>Department of Geological and Geophysical Sciences, Princeton University, Princeton, New Jersey 08544, U.S.A.

<sup>2</sup>Institute of Mineralogy and Petrography, Academy of Sciences, Novosibirsk, 630090, Russia

### ABSTRACT

Thermodynamic properties of the natural calcium zeolites laumontite, leonhardite, dehydrated leonhardite (metaleonhardite), wairakite, and yugawaralite were studied by calorimetry in lead borate solvent at 975 K. Enthalpies of formation from the elements at 298 K are as follows:  $-7251.0 \pm 8.5$  kJ/mol for laumontite,  $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ ;  $-7107.3 \pm 5.6$  kJ/mol for leonhardite,  $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ ;  $-5964.3 \pm 5.1$  kJ/mol for metaleonhardite,  $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ ;  $-6646.7 \pm 6.3$  kJ/mol for wairakite,  $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ ; and  $-9051.3 \pm 10.4$  kJ/mol for yugawaralite,  $\text{CaAl}_2\text{Si}_6\text{O}_{16} \cdot 4\text{H}_2\text{O}$ . The value for leonhardite is in good agreement with early values from acid calorimetry (Barany 1961) but not with revised values from Hemingway and Robie (1977). The enthalpy of dehydration of leonhardite is  $140.2 \pm 6.7$  kJ/mol, and the loss of one mole of  $\text{H}_2\text{O}$  is associated with an endothermic effect of about 40 kJ. Standard entropies,  $S_{298}^\circ$ , of wairakite [400.7 J/(mol·K)] and yugawaralite [609.8 J/(mol·K)] were derived from our new enthalpy data combined with reversed  $P$ - $T$  phase equilibria (Liou 1971; Zeng and Liou 1982). The upper limit of wairakite stability, the univariant curve for equilibrium of wairakite with anorthite, quartz, and fluid, was calculated from these values of enthalpy and entropy. Good agreement between thermodynamic calculations and reversed phase equilibria supports the reliability of the new thermodynamic data.

### INTRODUCTION

Calcium zeolites are common in diagenetic, sedimentary, hydrothermally altered, and low-grade metamorphic environments. The major zeolites encountered are laumontite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ ), leonhardite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ ), which is the partially dehydrated form of laumontite, wairakite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ ), which is the Ca analog of analcime, and yugawaralite ( $\text{CaAl}_2\text{Si}_6\text{O}_{16}$ ). Laumontite phase equilibria are especially important for  $P$ - $T$  boundaries of zeolite metamorphic facies (laumontite facies). Laumontite and wairakite are the index minerals of the zeolite facies.

$P$ - $T$  stability relations in the system  $\text{CaAl}_2\text{Si}_2\text{O}_8$ - $\text{SiO}_2$ - $\text{H}_2\text{O}$  have been investigated both experimentally and by thermodynamic calculations (Coombs et al. 1959; Koizumi and Roy 1960; Liou 1970, 1971; Thompson 1970; Zeng and Liou 1982; Senderov 1980, 1988; Ivanov and Gurevich 1975). Despite extensive experimental work, some of the zeolite phase relations have not been satisfactorily determined because of low rates of reactions under low-temperature conditions and because of the formation of metastable phases in experiments of short duration. Many experiments report only synthesis and not reversals; hence, estimations of free energies are uncertain. Thermodynamic data obtained from experimental equilibria, estimated from  $P$ - $T$  data in contemporary geothermal systems, and calculated from various thermodynamic data sets show wide scatter. Thus, direct cal-

orimetric measurement of enthalpies of formation of calcium zeolites and related minerals is desirable.

No calorimetric data exist for the enthalpies and entropies of laumontite, wairakite, and yugawaralite. The enthalpy of formation of leonhardite was obtained by Barany (1961) using hydrofluoric-acid calorimetry and later revised by Hemingway and Robie (1977). Low-temperature heat capacity and entropy of leonhardite were measured by King and Weller (1961). All published thermodynamic data for wairakite, laumontite, and yugawaralite were estimated or calculated from phase equilibrium studies and are not consistent. For instance, the standard entropy values for wairakite given by Helgeson et al. (1978) and by Glushko (1981) differ by about 80 J/(mol·K).

Several studies (Kiseleva and Ogorodova 1983; Circone and Navrotsky 1992; Smelik et al. 1994; Navrotsky et al. 1994) have shown that oxide melt solution calorimetry can determine the enthalpies of formation of hydrous minerals. The present work uses drop-solution calorimetry in molten  $2\text{PbO} \cdot \text{B}_2\text{O}_3$  to measure the enthalpies of formation of natural laumontite, metaleonhardite (dehydrated laumontite), leonhardite, wairakite, and yugawaralite. The enthalpy of dehydration in the laumontite-leonhardite-metaleonhardite series was studied by transposed temperature-drop calorimetry. Stability relations of wairakite, yugawaralite, and leonhardite and the upper temperature limit for wairakite stability were calculated on the basis of these thermochemical data.

TABLE 1. Chemical analyses and formulas of zeolites studied

	Leonhardite	Wairakite	Yugawaralite
<b>Chemical analysis (wt%)</b>			
SiO <sub>2</sub>	51.72	55.20	61.78
TiO <sub>2</sub>	0.01	0.00	0.01
Al <sub>2</sub> O <sub>3</sub>	21.93	23.08	17.01
MnO	0.09	0.00	0.00
MgO	0.04	0.01	0.01
CaO	12.14	12.48	9.35
Na <sub>2</sub> O	0.05	0.01	0.02
K <sub>2</sub> O	0.13	0.02	0.01
P <sub>2</sub> O <sub>5</sub>	0.11	—	—
H <sub>2</sub> O*	13.67	8.44	11.98
Total	99.98	99.23	100.25
<b>Cations and H<sub>2</sub>O molecules per unit cell</b>			
Si	15.96 (16)**	32.17 (32)	12.08 (12)
Ti	0.00	0.00	0.00
Al	7.98 (8)	15.86 (16)	3.92 (4)
Mn	0.02	0.00	0.00
Mg	0.02	0.01	0.00
Ca	4.02 (4)	7.79 (8)	1.96 (2)
Na	0.03	0.06	0.00
K	0.05	0.01	0.00
P	0.03	—	—
H <sub>2</sub> O	14.07 (14)	16.41 (16)	7.80 (8)

\* H<sub>2</sub>O contents also verified by thermogravimetric analysis.  
 \*\* Values in parentheses represent ideal molar ratios.

### ZEOLITE CHARACTERIZATION

We chose well-crystallized natural samples of laumontite, wairakite, and yugawaralite. The laumontite is from Nidym River (left tributary of Nizhnyaya Tunguska River), Siberia. White to pinkish nontransparent crystals up to 10 mm in length were separated from hydrothermal veins in Triassic basalts.

Wairakite from Bondai-Atami, Koriyama City, Japan (Mineralogical Museum of the Russian Academy of Sciences, catalog no. 87622), was provided by A.A. Godovikov. It had been separated under a microscope from an aggregate of white semitransparent crystals with some admixture of quartz and calcite. Clean, transparent, colorless tabular crystals were used.

Yugawaralite from Obora Toi-cho, Shizuoko Prefecture, Japan (Mineralogical Museum of the Russian Academy of Sciences, catalog no. 87616), was provided by A.A. Godovikov. Transparent and semitransparent colorless platy crystals up to 1–2 mm in dimension from the central parts of the small veins (10–15 mm thick) in hydrothermally altered brecciated volcanic rock were chosen.

Leonhardite was chemically analyzed by X-ray fluorescence using a Carl Zeiss VRA 20R instrument. Wairakite and yugawaralite were analyzed by electron microprobe (Camebax MICRO). Table 1 shows the analytical data.

High-temperature thermal behavior was studied using a TA 3000 (Mettler) thermoanalyzer consisting of a thermogravimeter (293–1273 K), a scanning calorimeter (103–873 K), and a differential dilatometer (173–1273 K). The thermograms are similar to those published by Gottardi and Galli (1985) and are discussed further below.

TABLE 2. Lattice parameters of zeolites

<i>a</i> (Å)	<i>b</i> (Å)	<i>c</i> (Å)	$\beta$ (°)	<i>V</i> (Å <sup>3</sup> )	Ref.
<b>Leonhardite</b>					
14.747 (5)	13.067 (3)	7.532 (4)	111.9 (3)	1346 (1)	1
14.770	13.056	7.595	112.8		2
14.77 (2)	13.09 (2)	7.58 (2)	112.0 (1)		3
<b>Wairakite</b>					
13.700 (15)	13.666 (9)	13.559 (10)	90.44 (8)	882.9 (8)	1
13.692 (3)	13.643 (3)	13.560 (3)	90.50 (10)		4
<b>Yugawaralite</b>					
6.728 (5)	14.007 (6)	10.049 (9)	111.20 (4)	882.9 (8)	1
6.729	14.008	10.050	111.18		5

Note: For unit cells on the basis of 48 (leonhardite), 96 (wairakite), and 32 (yugawaralite) O atoms in the framework. Values in parentheses represent confidence interval (for 95% probability). References are as follows: 1 = this work, 2 = Pipping (1966), 3 = Yamazaki et al. (1991), 4 = Takéuchi et al. (1979), and 5 = Eberlein et al. (1971).

Lattice parameters of samples studied are shown in Table 2. They agree well with previous studies (Gottardi and Galli 1985).

### CALORIMETRY

The enthalpies of formation and dehydration were determined using a Tian-Calvet high-temperature heat-flux microcalorimeter described in detail by Navrotsky (1977). Drop-solution calorimetric methods (Chai and Navrotsky 1993; Navrotsky et al. 1994) were chosen to avoid decomposition of zeolites at the calorimeter temperature prior to dissolution. The samples were dropped from room temperature into molten 2PbO·B<sub>2</sub>O<sub>3</sub> at 975 K. Most calorimetric experiments were performed using pressed pellets about 3 mm in diameter, 0.5–1 mm in height, and 10–25 mg in weight. A few experiments used pieces of single crystals. The heat of drop solution was a sum of the heat of solution in the melt plus the heat content ( $H_{975}^0 - H_{298}^0$ ).

Heats of dehydration and heat contents at 975 K were obtained using transposed temperature-drop calorimetry (sample dropped into an empty platinum crucible equilibrated in the calorimeter). The heat effect contained two contributions, the enthalpy of dehydration at 975 K and the heat content of the mineral ( $H_{975}^0 - H_{298}^0$ ). When the solid decomposition products were dropped into the calorimeter, the heat effect contained only the heat content ( $H_{975}^0 - H_{298}^0$ ) of the dehydrated zeolite.

We were concerned that when the small samples used in this study were dropped into the calorimeter, the heat picked up during the drop, before the sample reached the region of the thermopile, might be a different fraction of the total heat effect than seen for more massive platinum-encapsulated samples, for 30–50 mg pellets, or for platinum calibration pieces weighing about 200 mg. Rather than using such large platinum pieces for calibration in the usual procedure (Navrotsky 1977), we chose to calibrate using corundum pellets of weight similar to that of the samples. The corundum (Johnson Matthey 99.999%)

TABLE 3. Results of calorimetric experiments at 975 K

Mineral	Formula	F.W.*	Observed enthalpy			
			Drop**		Dropso†	
			(J/g)	(kJ/mol)	(J/g)	(kJ/mol)‡
Laumontite	CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·4H <sub>2</sub> O	470.44	1502.8 ± 13.2(9)	707.0 ± 6.2(9)		
Leonhardite	CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·3.5H <sub>2</sub> O	461.44	1450.6 ± 13.1(6)	669.4 ± 6.0(6)	1390.8 ± 6.4(9)	641.8 ± 3.0(9)§
Dehydrated leonhardite (metaleonhardite)	CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub>	398.39	722.3 ± 7.0(7)	287.8 ± 2.8(7)	646.9 ± 4.9(6)	257.7 ± 2.0(6)§
Wairakite	CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·2H <sub>2</sub> O	434.41			1171.3 ± 9.7(10)	(508.8 ± 4.2(10)§) 506.4 ± 4.2(10)
Yugawaralite	CaAl <sub>2</sub> Si <sub>6</sub> O <sub>16</sub> ·4H <sub>2</sub> O	590.62			1235.1 ± 13.5(7)	(729.5 ± 8.0(7)§) 734.2 ± 8.0(7)
Corundum	Al <sub>2</sub> O <sub>3</sub>	101.96			1058.6 ± 9.6(8)	107.9 ± 1.0(8)§
Quartz	SiO <sub>2</sub>	60.08	715.0 ± 5.3(12)	43.0 ± 0.3(12)	551.3 ± 5.2(9)	39.1 ± 0.3(9)§
Calcite	CaCO <sub>3</sub>	100.09			1931.8 ± 6.8(10)	193.4 ± 0.7(10)§

\* F.W. = gram formula weight.

\*\* Drop = transposed temperature-drop calorimetry, no solvent present.

† Dropso = drop solution into 2PbO·B<sub>2</sub>O<sub>3</sub> solvent.

‡ Uncertainty is two standard deviations of mean; number in parentheses is number of experiments.

§ Assuming ideal stoichiometry.

|| Corrected for nonideal H<sub>2</sub>O content; this is the preferred value.

was heated at 1773 K for 15 h to ensure dryness, crystallinity, and complete conversion to  $\alpha$ -Al<sub>2</sub>O<sub>3</sub>. The calibration factor obtained (using  $H_T^0 - H_{298}^0$  from the National Bureau of Standards Certificate, 1982) for corundum was indeed about 2% higher than that obtained for the same calorimeter using large platinum pieces. We believe that this modified calibration method better reflects the conditions used in the present experiments. For the calibrations, an uncertainty of about  $\pm 0.5\%$  (two standard deviations of the mean) was obtained.

Because laumontite is not stable at room temperature in air, it was obtained from leonhardite by keeping the sample in water-saturated conditions for 10–16 h. Laumontite with a little excess water (not more than  $6 \times 10^{-5}$  mole H<sub>2</sub>O per mole laumontite) was wrapped in a small platinum capsule and dropped from room temperature into an empty platinum crucible in the calorimeter. The measured heat effect contained four contributions: the enthalpy of dehydration at 975 K, the heat content of the mineral, the platinum capsule heat content, and the excess water heat content. Using platinum capsules introduced the need for additional calibration by dropping empty platinum capsules. The platinum heat content was 10–12% of the total measured heat effect, and the excess water heat effect was about 2%.

All experiments were performed under a dry-air atmosphere flowing through the calorimeter at 30–40 cm<sup>3</sup>/min. Navrotsky et al. (1994) showed that calorimetry under flowing gas results in a negligible enthalpy of interaction between the evolved H<sub>2</sub>O and the solvent. Under flowing gas conditions we have been able to obtain reproducible results with excellent baseline stability.

CaCO<sub>3</sub>, rather than CaO, was used as the Ca reference material because pellets of CaO did not dissolve reproducibly and showed evidence of partial hydration. Chai and Navrotsky (1993) showed that CaCO<sub>3</sub> can be used reliably for calorimetric purposes under flowing atmo-

sphere at 975 K. The calcite (Aldrich, 99.995%) and quartz (Fluka, >99.9%) were commercial products, dried at 383 K prior to calorimetry.

### CALORIMETRIC RESULTS

Calorimetric data are summarized in Table 3. Six to 12 enthalpy measurements were made for each substance. The calibration factor did not depend on the mass of the alumina dropped, which ranged from 10 to 25 mg, and neither the heat of drop solution nor the heat of transposed temperature drop (in joules per gram or kilojoules per mole) varied with sample mass. This consistency, see Figure 1, indicates proper equipment performance and reproducible sample dissolution. The heats of drop solution did not depend on the amount of sample previously dissolved (up to about 150 mg in 30 g lead borate), and the calorimetric curves returned cleanly to the original baseline in almost all experiments. The statistical errors reported (two standard deviations of the mean) are  $\pm 1\%$  or better, which also indicates problem-free dissolution.

Table 4 compares the enthalpies of drop solution, solution, and transposed temperature drop obtained for corundum, quartz, and calcite in this work with values from previous studies at the same temperature. In general the agreement is good. Our value for the enthalpy of drop solution of calcite is slightly higher than that of Chai and Navrotsky (1993); this may reflect the difference between using calibration factors based on small alumina pellets and large platinum pieces. However, similar differences are not seen for quartz and corundum. We note that both solution calorimetry and drop-solution calorimetry on quartz and corundum are frequently performed in our laboratory to train new users and check calorimeter operation; this accounts for the large amount of unpublished data in Table 4. Because the enthalpy of solution of quartz





FIGURE 1. Observed enthalpy of drop solution vs. mass of sample used for leonhardite, wairakite, and yugawaralite.


appears to depend significantly on temperature, only data at 973–978 K are included in Table 4. Lastly, we note considerable scatter in the tabulated heat contents of quartz; this largely reflects the different ways of treating the energetics of the  $\alpha$ - $\beta$  transition.

#### ENTHALPIES OF DEHYDRATION OF LEONHARDITE AND LAUMONTITE


The enthalpy of dehydration of leonhardite may be calculated (see thermodynamic cycle A in the Appendix Table). For the reaction



The enthalpy of dehydration of laumontite (see cycle B) is given by



The enthalpy of partial dehydration of laumontite to leonhardite is represented by



Its enthalpy (see cycle C) is  $3.2 \pm 4.6$  kJ/mol. This value is essentially zero.

Because this reversible transition takes place in air at near room temperature, Crawford and Fyfe (1965) suggested that the free-energy change of this reaction is extremely small, not exceeding a few hundred joules. If we assume that  $\Delta G^0$  for the laumontite-leonhardite reaction is zero, we can calculate its enthalpy change using entropy values from Helgeson et al. (1978). The estimated en-

TABLE 4. Comparison of enthalpies of drop solution of quartz, corundum, and calcite obtained at various times in this laboratory

Drop solution	Solution	Transposed temperature drop (heat content)
	<b>Corundum, <math>\alpha</math>-<math>\text{Al}_2\text{O}_3</math></b>	
$107.9 \pm 1.0^a$	$33.5 \pm 0.8^b$	$74.3 \pm 0.4^a$
$107.4 \pm 2.1^c$	$33.0 \pm 0.8^d$	$74.2 \pm 0.8^a$
$107.9 \pm 0.7^f$	$32.8 \pm 0.3^g$	$75.3^a$
	$(33.6 \pm 1.0)^j$	
	<b>Quartz, <math>\text{SiO}_2</math></b>	
$39.1 \pm 0.3^a$	$-4.3 \pm 0.2^e$	$43.0 \pm 0.3^a$
$40.0 \pm 0.2^i$	$-4.3 \pm 0.2^k$	$42.3^a$
$39.3 \pm 0.4^a$	$-3.9 \pm 0.4^e$	$44.0^a$
		$45.1^m$
	<b>Calcite, <math>\text{CaCO}_3</math></b>	
$193.4 \pm 0.7^a$		
$189.9 \pm 0.7^l$		

Note: a = this work, alumina calibration; b = Brown and Navrotsky (1989, unpublished), Pt calibration; c = Tinker and Navrotsky (1993, unpublished), Pt calibration; d = Hovis and Navrotsky (1994, unpublished), Pt calibration; e = Petrovic and Navrotsky (1994, unpublished), Pt calibration; f = Gerardin et al. (1994), Pt calibration; g = calculated as enthalpy of drop solution minus enthalpy of transposed temperature drop, this work; h = Robie et al. (1978), values for corundum similar in other tabulations; i = Ellison and Navrotsky (1992), Pt calibration; j = Chai and Navrotsky (1993), Pt calibration; k = Berman (1988), Holland and Powell (1990); l = Richet et al. (1982); and m = Hemingway (1987).

thalpy change is 3.1 kJ/mol, in excellent agreement with experimental data. Thus, our experiments show that the loss of  $\text{H}_2\text{O}$  from laumontite at room temperature is indeed associated with a very small endothermic effect of about 6 kJ/mol  $\text{H}_2\text{O}$  lost. In contrast, the dehydration of leonhardite is associated with more significant endothermic effects (about 40 kJ/mol  $\text{H}_2\text{O}$ ).

The enthalpy of dehydration of the sodium zeolite analcime was measured by acid calorimetry by Barany (1961), who obtained about 30 kJ/mol  $\text{H}_2\text{O}$ , similar to our value for leonhardite. However, such comparisons can be misleading. Dehydration is a very complicated process that can occur continuously or stepwise and may be accompanied by changes in the aluminosilicate framework. For example, as shown by Belitsky et al. (1992, 1993), dehydration can be followed by compression of the crystal structure owing to framework polyhedral tilting. Different zeolite frameworks deform differently and have different positions of the  $\text{H}_2\text{O}$  molecules within the channels. Thus, the dehydration process related to such structural changes may be accompanied by different energy effects.

Yakubovich and Simonov (1985) refined the crystal structure of leonhardite, giving it the structural formula  $\text{Ca}(\text{H}_2\text{O})_{2.8}(\text{Al}_2\text{Si}_4\text{O}_{12}) \cdot 0.5\text{H}_2\text{O}$  to represent better the different roles of  $\text{H}_2\text{O}$  in the crystal structure. Four positions were occupied by  $\text{H}_2\text{O}$  molecules. Our differential scanning calorimetry (DSC) and differential thermogravimetry curves show three peaks, but the third peak is asym-

metric and its high-temperature shoulder may be related to removal of the most tightly bonded H<sub>2</sub>O molecules.


Using DSC we measured the total heat effects associated with dehydration of leonhardite, CaAl<sub>2</sub>Si<sub>4</sub>O<sub>12</sub> · 3.5H<sub>2</sub>O, from 298 to 863 K to be about 139 kJ/mol, which agrees with the value measured by transposed temperature calorimetry (143 kJ/mol). DSC measurements show the disproportionate distribution of energy into several peaks. The first peak reflects the loss of weakly bonded zeolitic water without changes in the Al-Si framework and has a small heat effect corresponding to about 6–8% of the total heat of dehydration. The second peak represents removal of H<sub>2</sub>O molecules bonded to both framework O atoms and Ca and comprises about 20% of the total heat effect. The third peak is related to loss of water from the Ca coordination sphere and the formation of a metastable dehydrated phase, accompanied by changes in the framework structure. It accounts for 70–80% of the total energy. The loss of one mole of H<sub>2</sub>O during the first and the second peaks has an endothermic effect about 10–20 kJ/mol, but removing H<sub>2</sub>O molecules from the Ca coordination sphere requires about 70–80 kJ/mol.

The enthalpy of dehydration of yugawaralite was evaluated only by DSC to be about 150 kJ/mol CaAl<sub>2</sub>Si<sub>6</sub>O<sub>16</sub> · 4H<sub>2</sub>O, or about 37.5 kJ/mol H<sub>2</sub>O. Kerr and Williams (1969) found four water sites, each with 75% occupancy and each linked to a Ca site. All hydrogen bonds were described as long or very long. The DSC curve shows three main peaks, but the third peak, as for leonhardite, has an asymmetric high-temperature shoulder. The first peak (about 10–15% of the total heat effect) presumably corresponds to loss of weakly bonded H<sub>2</sub>O molecules with the longest hydrogen bonds. The loss of one mole of such H<sub>2</sub>O requires 5–20 kJ. The second peak corresponds to 15–20% of the total heat effect with an endothermic effect of about 10–20 kJ/mol H<sub>2</sub>O. The third peak (about 70% of the total heat effect) probably corresponds to the removal of H<sub>2</sub>O molecules with shorter hydrogen bonds. This is associated with a larger endothermic effect of about 70 kJ/mol.

## ENTHALPIES OF FORMATION OF CALCIUM ZEOLITES

### Leonhardite and metaleonhardite

The enthalpy of formation of leonhardite from the oxides at 298 K was calculated using two thermochemical cycles for the reaction



One uses the enthalpy of drop solution of leonhardite and of constituent oxides (see cycle D). The second cycle (E) is used to derive the enthalpy of formation of leonhardite and metaleonhardite. It involves dehydration and subsequent dissolution of fully dehydrated samples.


The enthalpy of formation of leonhardite from the oxides at 298 K is  $-153.3 \pm 3.5$  kJ/mol using the first cycle and  $-150.9 \pm 7.2$  kJ/mol using the second cycle. Because

the first cycle requires fewer steps, it yields a value with smaller uncertainty and we adopt this as our preferred value. The agreement between the first and second cycles indicates the overall consistency of the data.


Per mole H<sub>2</sub>O, the difference between the enthalpies of formation obtained from the first and second cycles is  $0.7 \pm 2.3$  kJ/mol. Because the second cycle does not use any reactions that dissolve a hydrous phase in lead borate, whereas the first cycle does, their good agreement once more validates the methodology of drop-solution calorimetry under flowing atmosphere for volatile-bearing phases (Navrotsky et al. 1994).

The standard enthalpy of formation of leonhardite from oxides obtained in this work is  $-153.3 \pm 3.5$  kJ/mol. This agrees within experimental error with the value of  $-154.2 \pm 2.7$  kJ/mol (Barany 1961) previously obtained from solution calorimetry in aqueous hydrofluoric acid for a natural sample with slightly higher (by 0.33%) water content. Later, Hemingway and Robie (1977) recalculated Barany's data. They considered that the enthalpy of formation of gibbsite, Al(OH)<sub>3</sub>, obtained by Barany and Kelley (1961) was incorrect and that the heat of solution value used for  $\alpha$ -quartz refers to very small particles. They revised Barany's data and recommended a value of  $-169.2 \pm 3.5$  kJ/mol for the enthalpy of formation. Our new determination supports the original rather than the revised HF value.

For the formation of metaleonhardite (fully dehydrated leonhardite),



(see cycle F). Thus, the dehydrated zeolite has only a rather modest energetic stability with respect to the oxides and is metastable with respect to other anhydrous phase assemblages. For the reaction (see cycle G)



On the basis of two moles of O (Ca<sub>w</sub>Al<sub>y</sub>Si<sub>z</sub>O<sub>2</sub>), metaleonhardite is only 13.1 kJ/mol higher energetically with respect to its stable phase assemblage of anorthite + quartz. Petrovic et al. (1993) found that a series of synthetic zeolites of SiO<sub>2</sub> composition were 10–15 kJ/mol higher in energy than quartz, and Hu et al. (1995) found that several aluminophosphate zeolites (Al<sub>0.5</sub>P<sub>0.5</sub>O<sub>2</sub>) were only 7–10 kJ/mol higher in energy than berlinite. Thus, the present data support the observation that anhydrous open zeolite frameworks are only slightly destabilized with respect to their dense anhydrous stable assemblages.

The enthalpies of formation from the elements were calculated with the use of the thermodynamic values of the constituent oxides taken from Robie et al. (1978) and the enthalpies of formation from the oxides obtained in this study:  $\Delta H_f$  at 298 K =  $-7107.3 \pm 5.6$  kJ/mol for leonhardite (CaAl<sub>2</sub>Si<sub>4</sub>O<sub>12</sub> · 3.5H<sub>2</sub>O), and  $-5964.3 \pm 5.1$  kJ/mol for metaleonhardite (CaAl<sub>2</sub>Si<sub>4</sub>O<sub>12</sub>).

**TABLE 5.** Thermodynamic properties at 298 K of some Ca-zeolites obtained from this study and from the literature

Mineral and formula	$\Delta H_f^0$ (kJ/mol)	Ref. (methods)	$S^0$ [J/(mol·K)]	Ref. (methods)
Laumontite (CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·4H <sub>2</sub> O)	-7251.0 ± 8.5*	1, C	485.3 ± 20.9	2, E
	-7268.9 ± 6.3	2, PE	485.76	3, E
	-7231.3 ± 10.1	4, PE	499.2 ± 19.2	5, E
	-7233.6	3, PE		
	-7265.6	6, E		
		1, C	461.1 ± 5.45	7, C
Leonhardite (CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·3.5H <sub>2</sub> O)	-7107.3 ± 5.6	8, PE		
	-7123.2 ± 4.8	9, C		
	-7108.3 ± 5.2	1, C	255.0	1, E
Metaleonhardite (CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> )	-5964.3 ± 5.1	1, C	400.7	1, PE, C
Wairakite (CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> ·2H <sub>2</sub> O)	-6646.7 ± 6.3	2, PE	376.6 ± 20.9	2, PE
	-6678.9 ± 5.0	4, PE	439.7	3, PE
	-6606.9 ± 8.1	3, PE	463.6 ± 4.2	5, E
	-6608.8	1, C	609.8	1, C, PE
Yugawaralite (CaAl <sub>2</sub> Si <sub>6</sub> O <sub>16</sub> ·4H <sub>2</sub> O)	-9051.3 ± 10.4	10, PE	609.9	10, PE
	-9036.19	6, E		
	-9087.6			

Note: References are as follows: 1 = this study, 2 = Glushko (1981), 3 = Helgeson et al. (1978), 4 = Senderov (1980), 5 = Zen (1972), 6 = Chermak and Rimstidt (1989), 7 = King and Weller (1961), 8 = Hemingway and Robie (1977), 9 = Barany (1961), and 10 = Zeng and Liou (1982). Methods are as follows: C = calorimetry, PE = phase equilibrium, and E = estimation.


\* Enthalpy of formation from elements.

### Laumontite

Calorimetric study of laumontite under excess water allows us to calculate the enthalpy of formation of laumontite using two cycles. First (cycle H), we can use the enthalpy of the reaction leonhardite = laumontite, the enthalpy of formation of leonhardite obtained in this work, and the data for H<sub>2</sub>O from Robie et al. (1978). The enthalpy of formation from the elements of laumontite (CaAl<sub>2</sub>Si<sub>4</sub>O<sub>12</sub>·4H<sub>2</sub>O) is -7253.4 ± 10.3 kJ/mol. The second cycle (I) uses the complete dehydration of laumontite to metaleonhardite, the enthalpy of formation of metaleonhardite, and data for H<sub>2</sub>O. The calculated enthalpy of formation from the elements is -7251.0 ± 8.5 kJ/mol. This value is close to that obtained using the first cycle, but the second cycle yields a value with smaller uncertainty and we adopt this as our preferred value (Table 4). Our value lies between previously reported data based on phase equilibria, -7233.6 kJ/mol (Helgeson et al. 1978) and -7268.9 ± 6.3 kJ/mol (Glushko 1981).

### Wairakite


The wairakite used has a nearly stoichiometric composition, except for a slightly higher water content (0.15%), as is very common for natural samples of wairakite (Gottardi and Galli 1985). The very small amounts of Na, K, and Mg have a negligible effect on the enthalpy, but we made a correction to the enthalpy of drop solution for 0.05 mole of excess H<sub>2</sub>O. We subtracted the heat content of 0.41 moles of H<sub>2</sub>O from the measured heat effect and obtained the heat of drop solution of stoichiometric wairakite, 506.4 ± 4.2 kJ/mol (Table 3). This preferred value was used to calculate the enthalpy of formation. For the reaction



we obtained  $\Delta H_{f,ox}^0 = -123.5 \pm 4.7$  kJ/mol at 298 K using thermodynamic cycle J. The enthalpy of formation from the elements,  $\Delta H_{f,el}^0$ , is -6646.7 ± 6.3 kJ/mol. Without correction for water,  $\Delta H_{f,el}^0$  would be -6649.2 ± 6.3 kJ/mol. Previous data for  $\Delta H_{f,el}^0$  of wairakite calculated from phase equilibria or otherwise estimated differ significantly from each other, ranging from -6606.9 ± 5.0 kJ/mol (Senderov 1980) to -6678 ± 5.0 kJ/mol (Glushko 1981). Our value, the first direct calorimetric measurement, is close to the middle of this range (see Table 5). It is worth noting that solid solution exists between wairakite and its Na analog, analcime, and that this substitution influences crystal structure, thermodynamic properties, and phase equilibria (Aoki and Minato 1980).

### Yugawaralite

The yugawaralite studied is nearly stoichiometric (Eberlein et al. 1971; Gottardi and Galli 1985), but the H<sub>2</sub>O content is about 0.1 mole less than theoretical. The experimental value for the enthalpy of drop solution was corrected by adding the heat content of 0.1 moles of H<sub>2</sub>O (Table 3). The thermodynamic cycle K was used to calculate the enthalpy of formation from the oxides at 298 K. For the reaction



$\Delta H_{f,ox}^0 = -133.0 \pm 8.4$  kJ/mol. The enthalpy of formation from the elements is -9051.3 ± 10.4 kJ/mol. Without correction for the missing H<sub>2</sub>O,  $\Delta H_{f,el}^0$  would be -9046.5 ± 10.4 kJ/mol. Table 5 shows that only two estimates of  $\Delta H_f^0$  are available in the literature: -9036.19 kJ/mol (Zeng and Liou 1982) calculated from phase equilibria involving somewhat uncertain data for wairakite




FIGURE 2.  $P$ - $T$  diagram, showing the relations among zeolites in the presence of excess quartz and  $H_2O$  fluid. The curves marked a, b, and c represent the laumontite + yugawaralite, yugawaralite + wairakite, and laumontite + wairakite equilibria, respectively, calculated from thermochemical data and in agreement (to  $\pm 5$  K) with curves calculated by Zeng and Liou (1982). The metastable extension of the yugawaralite + wairakite reaction, reversed by Zeng and Liou (1982), is shown as a dashed curve. Curve d represents the breakdown of wairakite to anorthite as calculated from thermochemical data. The four solid circles represent the reversals of that equilibrium by Liou (1970). The error in these points is approximately  $\pm 5$  K and  $\pm 20$  bars.

from Helgeson et al. (1978), and a significantly more exothermic value estimated by Chermak and Rimstidt (1989). Our value is closer to the value calculated from phase equilibria.

#### CALCULATION OF THE ENTROPIES OF WAIRAKITE AND YUGAWARALITE

Low-temperature heat capacity and entropy at 298 K have been measured only for leonhardite (King and Weller 1961). There are many estimates of the entropies of the other minerals in the various datasets (Table 5).

Because leonhardite is related to laumontite, the standard entropy of laumontite has been estimated rather consistently by different authors using the leonhardite data (see Table 5). In contrast, estimates of the entropy of wairakite from phase equilibria scatter from  $376.6 \pm 20.9$  J/(mol·K) (Glushko 1981) to  $463.6 \pm 4.2$  J/(mol·K) (Zen 1972). The standard entropy of yugawaralite was calculated in only one study (Zeng and Liou 1982) on the basis of a value for the entropy of wairakite from Helgeson et al. (1978).

The standard entropies of wairakite and yugawaralite were derived from our new, independently determined enthalpy data, in combination with experimentally reversed  $P$ - $T$  phase equilibria. The equilibrium decompo-

sition of laumontite to wairakite is the most suitable reaction for calculating the entropy of wairakite:




Because the breakdown product of laumontite may be either ordered or disordered wairakite depending on the length of the experiments, we did not consider studies that had no proof of reaction reversal and for which the products may have represented various intermediate stages of this reaction. The equilibria demonstrated by Liou (1971) appear reliable. From these reversals, we calculated the entropy of wairakite using

$$\Delta G_{P,T} = \Delta H_r^0 - T\Delta S_r^0 + \Delta V_s^0 P + RT \ln f_{H_2O} = 0 \quad (10)$$

where  $\Delta G_{P,T}$ ,  $\Delta H_r^0$ , and  $\Delta S_r^0$  are the total free energy, enthalpy, and entropy change of the reaction, respectively,  $\Delta V_s^0$  is the volume change among the solid phases of the reaction, and  $f_{H_2O}$  is the fugacity of  $H_2O$  (Burnham et al. 1969). Because no experimental heat-capacity data are available for laumontite and wairakite we assume as a first approximation that  $\Delta C_p$  for the reaction is 0 and that  $\Delta V$  of the solid phases is constant. These assumptions are reasonable at the low- $P$  and low- $T$  conditions in these studies. Using the new calorimetric data for the enthalpies of formation of laumontite and wairakite we calculated the enthalpy of the laumontite-wairakite equilibrium. We used the value for entropy of laumontite of 485.76 J/(mol·K) from Helgeson et al. (1978). Using the molar volumes of laumontite and wairakite from Helgeson et al. (1978) we calculated  $\Delta V_s^0 = -2.07$  J/bar. The resulting standard entropy of wairakite is 400.7 J/(mol·K). This is in the midrange of estimates from different datasets (Table 5).

The entropy of yugawaralite at 298 K can be derived analogously. At present there is only one experimental study (Zeng and Liou 1982) of the univariant equilibrium curve for the dehydration of yugawaralite:




Even though this reaction was studied in a metastable field (see Fig. 2), the steep slope of the  $P$ - $T$  curve was established reliably. We obtained a value for the standard entropy of yugawaralite of 609.8 J/(mol·K) (Table 5). Our value for  $S^0$  of yugawaralite is in excellent agreement with the value calculated by Zeng and Liou (1982).

The thermochemical calculations give stability fields of laumontite, wairakite, and yugawaralite that agree with reversed phase equilibria (Liou 1971; Zeng and Liou 1982) (see Fig. 2). Because the entropies used were based in part on these reversals, such agreement is expected. Nevertheless, the simultaneous good fit to several equilibria using the calorimetric results strongly suggests internal consistency in both the calorimetric and phase-equilibrium studies.

Recommended values for heats and entropies of formation, and for standard entropies based on the above considerations, are listed in Table 6 for all the zeolites studied.

### GEOLOGIC APPLICATIONS

Several reactions constrain minerals in the zeolite facies; one such reaction that determines the upper limits of wairakite stability is



Coombs et al. (1959) reported that the equilibrium temperature for this reaction at 2 kbar is near 623 K. Koizumi and Roy (1960) suggested that the maximum temperature for this reaction is 733 K at 1 kbar and 743 K at 2 kbar. These studies were synthesis experiments rather than reversals. Seki (1968) noted that the synthetic wairakite used in these studies was a tetragonal modification, disordered wairakite. The equilibrium dehydration of wairakite to anorthite, quartz, and water (Liou 1970) (see Fig. 2) yielded temperatures of the breakdown curve of wairakite much lower than the earlier data.

Using our new enthalpy and entropy values, we calculated the univariant curve of the equilibrium of wairakite with anorthite, quartz and fluid. The Gibbs free energy of reaction at  $P$  and  $T$  was calculated as above, and we assumed constant  $\Delta C_p$  and  $\Delta V_s$ . We used values for the molar volume, standard entropy, and enthalpy of quartz and  $\text{H}_2\text{O}$  from Robie et al. (1978), volume and entropy of anorthite from Robie et al. (1978), the enthalpy of synthetic anorthite from a recent publication of Zhu et al. (1994) and the volume of wairakite from Helgeson et al. (1978). Thermodynamic properties of  $\text{H}_2\text{O}$  at  $P$  and  $T$  were taken from Burnham et al. (1969).

The calculated breakdown curve of wairakite (Fig. 2) agrees reasonably with Liou's (1970) phase equilibria. The temperature of the calculated univariant curve at 500 bars is near the experimental value, but at 1000–2000 bars calculated temperatures are slightly lower than experimental temperatures. At a given pressure, the calculated temperature can change by as much as 50 K depending on the values of  $\Delta H$ ,  $\Delta S$ , and  $\Delta V$  chosen within their allowable ranges. The assumptions of negligible  $\Delta C_p$  and constant  $\Delta V_s$  add other uncertainties. However, these factors probably do not vary independently, and a rigorous assessment of uncertainties is impossible. We consider the agreement with experimental values encouraging, especially because the wairakite breakdown reaction was not used to constrain the thermochemical data. Thus, this thermochemical calculation of the upper thermal stability limit of wairakite supports the consistency and reliability of our new calorimetric data. The calculated  $P$ - $T$  field of wairakite agrees with field observations in low-grade metamorphic and hydrothermally altered rocks (Coombs et al. 1959). The thermochemical data confirm that wairakite requires higher temperatures for its formation than laumontite. They also support the topology of the lau-

TABLE 6. Recommended values of thermodynamic properties of zeolites obtained in this study

Mineral and formula	$\Delta H_f^{\circ}_{298.15}$ (kJ/mol)	$S^{\circ}_{298.15}$ [J/(mol·K)]	$\Delta G_f^{\circ}_{298.15}$ (kJ/mol)
Laumontite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ )	$-7251.0 \pm 8.5$	485.5*	-6698.9
Leonhardite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ )	$-7107.3 \pm 5.6$	461.1**	-6582.7
Metaleonhardite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ )	$-5964.3 \pm 5.1$	255.0	-5621.6
Wairakite ( $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ )	$-6646.7 \pm 6.3$	400.7	-6208.3
Yugawaralite ( $\text{CaAl}_2\text{Si}_6\text{O}_{16} \cdot 4\text{H}_2\text{O}$ )	$-9051.3 \pm 10.4$	609.8	-8402.6

\* Helgeson et al. (1978).  
\*\* King and Weller (1961).

montite-wairakite-yugawaralite phase diagram, with an invariant point (in the presence of excess quartz and water) near 500 bars and 500 K.

These thermochemical data can be used to calculate other equilibria involving these minerals.  $P_{\text{CO}_2}$  and  $P_{\text{H}_2\text{O}}$  may significantly affect the stability of calcium zeolites. These minerals are unstable in the presence of even a moderate amount of  $\text{CO}_2$  because the partial pressure of  $\text{CO}_2$  lowers the fugacity of water and because Ca is consumed in carbonates. Such conditions are rather common for the calcium zeolite assemblages in uranium deposits (Mironenko and Naumov 1982). Thus,  $P$ - $T$  conditions of calcium zeolite formation strongly depend on the geothermal gradient,  $P_{\text{H}_2\text{O}}/P_{\text{tot}}$ ,  $P_{\text{CO}_2}$ , and other factors, including solution composition (Na,K). With appropriate care, the thermochemical data produced by this study are useful in understanding zeolite stability in complex natural systems. Equally important, a knowledge of thermodynamic stability helps distinguish equilibrium from nonequilibrium conditions, a constant problem in low-temperature environments.

### ACKNOWLEDGMENTS

This work was supported by the U.S. Department of Energy (grant DE-FG02-85ER 13437) and the RFFI of the Academy of Sciences, Russia. We thank A. Godovikov for providing samples. We thank L. Topor, L. Chai, E. Smelik, and K. Bose for advice and help with experimental procedures, and S. Wasylenko and K. Rozenberg for help with manuscript preparation.

### REFERENCES CITED

- Aoki, M., and Minato, H. (1980) Lattice constants of wairakite as a function of chemical composition. *American Mineralogist*, 65, 1212–1216.
- Barany, R. (1961) Heats and free energies of formation of some hydrated and anhydrous sodium- and calcium-aluminum silicates. U.S. Bureau of Mines, 5900, 1–18.
- Barany, R., and Kelley, K.K. (1961) Heats of formation of gibbsite, kaolinite, halloysite, and dickite. U.S. Bureau of Mines Report Inv. 5825, 1–13.
- Belitsky, I.A., Fursenko, B.A., Gabuda, S.P., Kholdeev, O.V., and Seryotkin, Y.V. (1992) Structural transformations in natrolite and edingtonite. *Physics and Chemistry of Minerals*, 18, 497–505.

- Belitsky, I.A., Fursenko, B.A., Seryotkin Yu.A., Goryainov, S.V., and Drebuschak, V.A. (1993) High-pressure, high-temperature behavior of wairakite. In Zeolite 1993 4th International Conference on the Occurrence, Properties, and Utilization of Natural Zeolites, Boise, Idaho, June 20–28, 1993, Program and Abstracts, 45–48.
- Berman, R.G. (1988) Internally-consistent thermodynamic data for minerals in  $\text{Na}_2\text{O}-\text{K}_2\text{O}-\text{CaO}-\text{MgO}-\text{FeO}-\text{Fe}_2\text{O}_3-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{TiO}_2-\text{H}_2\text{O}-\text{CO}_2$ . *Journal of Petrology*, 29, 445–522.
- Burnham, C.W., Holloway, J.R., and Davis, N.F. (1969) Thermodynamic properties of water to 1000 °C and 10,000 bars. *Geological Society America*, 132, 1–96.
- Chai, L., and Navrotsky, A. (1993) Thermochemistry of carbonate-pyroxene equilibria. *Contributions to Mineralogy and Petrology*, 114, 139–147.
- Chermak, J.A., and Rimstidt, J.D. (1989) Estimating the thermodynamic properties ( $\Delta G_f^\circ$  and  $\Delta H_f^\circ$ ) of silicate minerals at 298 K from the sum of polyhedral contributions. *American Mineralogist*, 74, 1023–1031.
- Circone, S., and Navrotsky, A. (1992) Substitution of  $^{6,9}\text{Al}$  in phlogopite: High-temperature solution calorimetry, heat capacities, and thermodynamic properties of the phlogopite-eastonite join. *American Mineralogist*, 77, 1191–1205.
- Coombs, D.S., Ellis, A.J., Fyfe, W.S., and Taylor, A.M. (1959) The zeolite facies with comments on the interpretation of hydrothermal syntheses. *Geochimica et Cosmochimica Acta*, 17, 53–107.
- Crawford, W.A., and Fyfe, W.S. (1965) Lawsonite equilibria. *American Journal of Science*, 263, 262–270.
- Eberlein, G.D., Erd, R.C., Weber, F., and Beatty, L.B. (1971) New occurrence of yugawaralite from the Chena Hot Springs area, Alaska. *American Mineralogist*, 56, 1699–1717.
- Ellison, A.J.G., and Navrotsky, A. (1992) Enthalpy of formation of zircon. *Journal of American Ceramic Society*, 75(6), 1430–1433.
- Gerardin, C., Sundaresan, S., Benziger, J., and Navrotsky, A. (1994) Structural investigation and energetics of mullite formation from sol-gel precursors. *Chemistry of Materials*, 6, 160–170.
- Glushko, V.P. (1981) Thermodynamic properties of individual substances, vol. 3, 251 p. Nauka, Moscow.
- Gottardi, G., and Galli, E. (1985) Natural zeolites, 409 p. Springer, Berlin.
- Helgeson, H.C., Delany, J.M., Nesbitt, H.W., and Bird, D.K. (1978) Summary and critique of the thermodynamic properties of rock-forming minerals. *American Journal of Science*, 278A, 1–229.
- Hemingway, B.S. (1987) Quartz: Heat capacities from 340 to 1000 K and revised values for the thermodynamic properties. *American Mineralogist*, 72, 273–279.
- Hemingway, B.S., and Robie, R.A. (1977) Enthalpies of formation of low albite  $\text{NaAlSi}_3\text{O}_8$ , gibbsite  $\text{Al}(\text{OH})_3$  and  $\text{NaAlO}_2$ ; revised values for  $\Delta H_f^\circ$  and  $\Delta G_f^\circ$  of some aluminosilicate minerals. *Journal of Research U.S. Geological Survey*, 5, 413–429.
- Holland, T.J.B., and Powell, R. (1990) An enlarged and updated internally consistent thermodynamic dataset with uncertainties and correlations: The system  $\text{K}_2\text{O}-\text{Na}_2\text{O}-\text{CaO}-\text{MgO}-\text{MnO}-\text{FeO}-\text{Fe}_2\text{O}_3-\text{Al}_2\text{O}_3-\text{TiO}_2-\text{SiO}_2-\text{C}-\text{H}_2-\text{O}_2$ . *Journal of Metamorphic Geology*, 8, 89–124.
- Hu, Y., Navrotsky, A., Chen, C.-Y., and Davis, M.E. (1995) Thermochemical study of the relative stability of dense and microporous aluminophosphate frameworks. *Chemistry of Materials*, 7, 1816–1823.
- Ivanov, P.P., and Gurevich, L.P. (1975) Experimental study of  $T-X(\text{CO}_2)$  boundaries of metamorphic zeolite facies. *Contributions to Mineralogy and Petrology*, 53, 55–60.
- Kerr, I.S., and Williams, D.J. (1969) The crystal structure of yugawaralite. *Acta Crystallography*, B25, 1183–1190.
- King, E.G., and Weller, W.W. (1961) Low-temperature heat capacities and entropies at 298.15 K of some sodium- and calcium-aluminum silicates. U.S. Bureau of Mines Report Inv. 5855, 1–8.
- Kiseleva, I.A., and Ogorodova, L.P. (1983) On using of the high-temperature calorimetry of dissolution for determination of enthalpies of formation of hydroxyl bearing minerals (using the examples of talc and tremolite). *Geokhimiya*, 12, 1745–1756 (in Russian).
- Koizumi, M., and Roy, R. (1960) Zeolite studies: 1. Synthesis and stability of the calcium zeolites. *Journal of Geology*, 68, 41–53.
- Liou, J.G. (1970) Synthesis and stability relations of wairakite  $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ . *Contributions to Mineralogy and Petrology*, 27, 259–282.
- (1971)  $P$ - $T$  stabilities of laumontite, wairakite, lawsonite, and related minerals in the system  $\text{CaAl}_2\text{Si}_2\text{O}_8-\text{SiO}_2-\text{H}_2\text{O}$ . *Journal of Petrology*, 12, 379–411.
- Mironenko, M.V., and Naumov, G.B. (1982) Physical and chemical conditions of laumontite formation in  $\text{CO}_2$ -hydrothermal systems. *Geokhimiya*, 4, 18–22 (in Russian).
- Navrotsky, A. (1977) Progress and new directions in high temperature calorimetry. *Physics and Chemistry of Minerals*, 2, 89–104.
- Navrotsky, A., Rapp, R.P., Smelik, E., Burnley, P., Circone, S., Chai, L., Bose, K., and Westrich, H.R. (1994) The behavior of  $\text{H}_2\text{O}$  and  $\text{CO}_2$  in high-temperature lead borate solution calorimetry of volatile-bearing phases. *American Mineralogist*, 79, 1099–1109.
- Petrovic, I., Navrotsky, A., Davis, M.E., and Zones, S.I. (1993) Thermochemical study of the stability of frameworks in high silica zeolites. *Chemistry of Materials*, 5, 1805–1813.
- Pipping, F. (1966) The dehydration and chemical composition of laumontite. *Mineral Society, India IMA*, 1, 159–166.
- Richet, P., Bottinga, Y., Denielou, L., Pettet, J.P., and Tequi, C. (1982) Thermodynamic properties of quartz, cristobalite and amorphous  $\text{SiO}_2$ : Drop calorimetry measurements between 1000 and 1800 K and a review from 0 to 2000 K. *Geochimica et Cosmochimica Acta*, 46, 2639–2658.
- Robie, R.A., Hemingway, B.S., and Fisher, J.R. (1978) Thermodynamic properties of minerals and related substances at 298.15 K and 1 bar (105 Pascals) pressure and at higher temperatures. U.S. Geological Survey Bulletin 1452, 456 p.
- Seki, Y. (1968) Synthesized wairakites their difference from natural wairakites. *Journal of Geological Society, Japan*, 74, 457–458.
- Senderov, E.E. (1980) Estimation of gibbs energy for laumontite and wairakite from conditions of their formation in geothermal areas. *Proceedings of The Fifth International Conference on Zeolites, Naples, Italy, June 2–6*.
- (1988) Physical-chemical aspects of zeolite formation in nature. In *Occurrence, properties and utilization of natural zeolites*, 111, 147 p.
- Smelik, E.A., Jenkins, D.M., and Navrotsky, A. (1994) A calorimetric study of synthetic amphiboles along the tremolite-tschermakite join and the heats of formation of magnesiohornblende and tschermakite. *American Mineralogist*, 79, 1110–1122.
- Takéuchi, Y., Mazzi, F., Haga, N., and Galli, E. (1979) The crystal structure of wairakite. *American Mineralogist*, 64, 993–1001.
- Thompson, A.B. (1970) Laumontite equilibria and the zeolite facies. *American Journal of Science*, 269, 267–275.
- Yakovovich, O.V., and Simonov, M.A. (1985) Refined crystal structure of the zeolite laumontite  $\text{Ca}(\text{H}_2\text{O})_{2.6}[\text{Al}_2\text{Si}_4\text{O}_{12}] \cdot 0.5\text{H}_2\text{O}$ . *Society of Physics and Crystals*, 30, 624–626.
- Yamazaki, A., Shiraki, T., Mishido, H., and Otsuka, R. (1991) Phase change of laumontite under relatively humidity-controlled conditions. *Clay Science*, 8, 79–86.
- Zen, E-an (1972) Gibbs free energy, enthalpy, and entropy of ten rock-forming minerals: Calculations, discrepancies, implications. *American Mineralogist*, 57, 524–553.
- Zeng, Y., and Liou, J.G. (1982) Experimental investigation of yugawaralite-wairakite equilibrium. *American Mineralogist*, 67, 937–943.
- Zhu, H., Newton, R.C., and Kleppa, O.J. (1994) Enthalpy of formation of wollastonite ( $\text{CaSiO}_3$ ) and anorthite ( $\text{CaAl}_2\text{Si}_2\text{O}_8$ ) by experimental phase equilibrium measurements and high-temperature solution calorimetry. *American Mineralogist*, 79, 134–144.

MANUSCRIPT RECEIVED MARCH 9, 1995

MANUSCRIPT ACCEPTED JANUARY 20, 1996

APPENDIX TABLE. Thermodynamic cycles

<b>A. Leonhardite dehydration</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) + $3.5\text{H}_2\text{O}$ (g, 975 K)	(1) <sup>a</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K)	(2) <sup>a</sup>
$3.5\text{H}_2\text{O}$ (g, 975 K) = $3.5\text{H}_2\text{O}$ (l, 298 K)	(3) <sup>b</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) + $3.5\text{H}_2\text{O}$ (l, 298 K)	(4)
$\Delta H_4 = \Delta H_1 + \Delta H_2 + \Delta H_3$	
<b>B. Laumontite dehydration</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) + $4\text{H}_2\text{O}$ (g, 298 K)	(5) <sup>a,c</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K)	(2) <sup>a,b</sup>
$4\text{H}_2\text{O}$ (g, 975 K) = $4\text{H}_2\text{O}$ (l, 298 K)	(6)
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) + $4\text{H}_2\text{O}$ (l, 298 K)	(7)
$\Delta H_7 = \Delta H_5 + \Delta H_2 + \Delta H_6$	
<b>C. Laumontite-leonhardite reaction</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) + $4\text{H}_2\text{O}$ (g, 298 K)	(5) <sup>a</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) + $3.5\text{H}_2\text{O}$ (g, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K)	-(1) <sup>a</sup>
$0.5\text{H}_2\text{O}$ (g, 975 K) = $0.5\text{H}_2\text{O}$ (l, 298 K)	(8) <sup>b</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K) + $0.5\text{H}_2\text{O}$ (l, 298 K)	(9)
$\Delta H_9 = \Delta H_5 - \Delta H_1 + \Delta H_8$	
<b>D. Formation of leonhardite (using drop-solution calorimetry of leonhardite)</b>	
$\text{CaCO}_3$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 975 K)	(10) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{CO}_2$ (g, 298 K) = $\text{CaCO}_3$ (xl, 298 K)	(11) <sup>b</sup>
$\text{CO}_2$ (g, 975 K) = $\text{CO}_2$ (g, 298 K)	(12) <sup>b</sup>
$\text{Al}_2\text{O}_3$ (xl, 298 K) = $\text{Al}_2\text{O}_3$ (sol, 975 K)	(13) <sup>a</sup>
$4\text{SiO}_2$ (xl, 298 K) = $4\text{SiO}_2$ (sol, 975 K)	(14) <sup>a</sup>
$3.5\text{H}_2\text{O}$ (l, 298 K) = $3.5\text{H}_2\text{O}$ (g, 975 K)	(3) <sup>b</sup>
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $4\text{SiO}_2$ (sol, 975 K) + $3.5\text{H}_2\text{O}$ (g, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$	(15) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $3.5\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K)	(16)
$\Delta H_{16} = \Delta H_{10} + \Delta H_{11} + \Delta H_{12} + \Delta H_{13} + \Delta H_{14} + \Delta H_3 + \Delta H_{15}$	
<b>E. Formation of leonhardite (using dehydration and drop-solution calorimetry of metaleonhardite)</b>	
$\text{CaCO}_3$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 975 K)	(10) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{CO}_2$ (g, 298 K) = $\text{CaCO}_3$ (xl, 298 K)	(11) <sup>b</sup>
$\text{CO}_2$ (g, 975 K) = $\text{CO}_2$ (g, 298 K)	(12) <sup>b</sup>
$\text{Al}_2\text{O}_3$ (xl, 298 K) = $\text{Al}_2\text{O}_3$ (sol, 975 K)	(13) <sup>a</sup>
$4\text{SiO}_2$ (xl, 298 K) = $4\text{SiO}_2$ (sol, 975 K)	(14) <sup>a</sup>
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $4\text{SiO}_2$ (sol, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K)	(17) <sup>a</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K)	-(2) <sup>a</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 975 K) + $3.5\text{H}_2\text{O}$ (g, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K)	(1) <sup>a</sup>
$3.5\text{H}_2\text{O}$ (l, 298 K) = $3.5\text{H}_2\text{O}$ (g, 975 K)	-(1) <sup>a,c</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $3.5\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K)	(16)
$\Delta H_{16} = \Delta H_{10} + \Delta H_{11} + \Delta H_{12} + \Delta H_{13} + \Delta H_{14} + \Delta H_{17} - \Delta H_2 - \Delta H_1$	
<b>F. Formation of metaleonhardite (fully dehydrated leonhardite)</b>	
$\text{CaCO}_3$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 975 K)	(10) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 298 K) = $\text{CaCO}_3$ (xl, 298 K)	(11) <sup>b</sup>
$\text{CO}_2$ (g, 975 K) = $\text{CO}_2$ (g, 298 K)	(12) <sup>b</sup>
$\text{Al}_2\text{O}_3$ (xl, 298 K) = $\text{Al}_2\text{O}_3$ (sol, 975 K)	(13) <sup>a</sup>
$4\text{SiO}_2$ (xl, 298 K) = $4\text{SiO}_2$ (sol, 975 K)	(14) <sup>a</sup>
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $4\text{SiO}_2$ (sol, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K)	(17) <sup>a</sup>
$\text{CaO}$ + $\text{Al}_2\text{O}_3$ + $4\text{SiO}_2$ = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$	(18)
$\Delta H_{18} = \Delta H_{10} + \Delta H_{11} + \Delta H_{12} + \Delta H_{13} + \Delta H_{14} + \Delta H_{17}$	
<b>G. Metaleonhardite = anorthite + quartz</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $4\text{SiO}_2$ (sol, 975 K)	-(17) <sup>a</sup>
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $2\text{SiO}_2$ (sol, 975 K) = $\text{CaAl}_2\text{Si}_2\text{O}_8$ (xl, 975 K)	(19) <sup>d</sup>
$\text{CaAl}_2\text{Si}_2\text{O}_8$ (xl, 975 K) = $\text{CaAl}_2\text{Si}_2\text{O}_8$ (xl, 298 K)	(20) <sup>d</sup>

APPENDIX TABLE.—Continued

$2\text{SiO}_2$ (sol, 975 K) = $2\text{SiO}_2$ (xl, 298 K)	(21) <sup>a</sup>
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_2\text{O}_8$ (xl, 298 K) + $2\text{SiO}_2$ (xl, 298 K)	(22)
$\Delta H_{22} = -\Delta H_{17} + \Delta H_{19} + \Delta H_{20} + \Delta H_{21}$	
<b>H. Laumontite formation (first cycle)</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K) + $0.5\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	-(9) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $3.5\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 3.5\text{H}_2\text{O}$ (xl, 298 K)	(16) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $4\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	(23)
$\Delta H_{23} = -\Delta H_5 + \Delta H_{16}$	
<b>I. Laumontite formation (second cycle)</b>	
$\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K) + $4\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	(24) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12}$ (xl, 298 K)	(18) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $4\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	(25)
$\Delta H_{25} = \Delta H_{24} + \Delta H_{18}$	
<b>J. Wairakite formation</b>	
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $4\text{SiO}_2$ (sol, 975 K) + $2\text{H}_2\text{O}$ (g, 975 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ (xl, 298 K)	(26) <sup>a</sup>
$\text{CaCO}_3$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 975 K)	(10) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{CO}_2$ (g, 298 K) = $\text{CaCO}_3$ (xl, 298 K)	(11) <sup>b</sup>
$\text{CO}_2$ (g, 975 K) = $\text{CO}_2$ (g, 298 K)	(12) <sup>b</sup>
$\text{Al}_2\text{O}_3$ (xl, 298 K) = $\text{Al}_2\text{O}_3$ (sol, 975 K)	(13) <sup>a</sup>
$4\text{SiO}_2$ (xl, 298 K) = $4\text{SiO}_2$ (sol, 975 K)	(14) <sup>a</sup>
$2\text{H}_2\text{O}$ (l, 298 K) = $2\text{H}_2\text{O}$ (g, 975 K)	(27) <sup>b</sup>
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $4\text{SiO}_2$ (xl, 298 K) + $2\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_4\text{O}_{12} \cdot 2\text{H}_2\text{O}$ (xl, 298 K)	(28)
$\Delta H_{28} = \Delta H_{26} + \Delta H_{10} + \Delta H_{11} + \Delta H_{12} + \Delta H_{13} + \Delta H_{14} + \Delta H_{27}$	
<b>K. Yugawaralite formation</b>	
$\text{CaO}$ (sol, 975 K) + $\text{Al}_2\text{O}_3$ (sol, 975 K) + $6\text{SiO}_2$ (sol, 975 K) + $4\text{H}_2\text{O}$ (g, 975 K) = $\text{CaAl}_2\text{Si}_6\text{O}_{16} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	(29) <sup>a</sup>
$\text{CaCO}_3$ (xl, 298 K) = $\text{CaO}$ (sol, 975 K) + $\text{CO}_2$ (g, 975 K)	(10) <sup>a</sup>
$\text{CaO}$ (xl, 298 K) + $\text{CO}_2$ (g, 298 K) = $\text{CaCO}_3$ (xl, 298 K)	(11) <sup>b</sup>
$\text{CO}_2$ (g, 975 K) = $\text{CO}_2$ (g, 298 K)	(12) <sup>b</sup>
$\text{Al}_2\text{O}_3$ (xl, 298 K) = $\text{Al}_2\text{O}_3$ (sol, 975 K)	(13) <sup>a</sup>
$6\text{SiO}_2$ (xl, 298 K) = $6\text{SiO}_2$ (sol, 975 K)	(30) <sup>a</sup>
$4\text{H}_2\text{O}$ (l, 298 K) = $4\text{H}_2\text{O}$ (g, 975 K)	-(6)
$\text{CaO}$ (xl, 298 K) + $\text{Al}_2\text{O}_3$ (xl, 298 K) + $6\text{SiO}_2$ (xl, 298 K) + $4\text{H}_2\text{O}$ (l, 298 K) = $\text{CaAl}_2\text{Si}_6\text{O}_{16} \cdot 4\text{H}_2\text{O}$ (xl, 298 K)	(31)
$\Delta H_{31} = \Delta H_{29} + \Delta H_{10} + \Delta H_{11} + \Delta H_{12} + \Delta H_{13} + \Delta H_{30} - \Delta H_6$	

Note: a = this work; b = Robie et al. (1978); c = the materials produced by dehydrating laumontite and leonhardite are essentially the same and have the same heat contents, called  $\Delta H_2$  in the cycles above; and d = Zhu et al. (1994).