# Crystal chemistry of phlogopite from Vulture-S. Michele Subsynthem volcanic rocks (Mt. Vulture, Italy) and volcanological implications

# SILVIA MATARRESE,<sup>1</sup> EMANUELA SCHINGARO,<sup>1</sup> FERNANDO SCORDARI,<sup>1,\*</sup> FRANCESCO STOPPA,<sup>2</sup> GIANLUIGI ROSATELLI,<sup>2</sup> GIUSEPPE PEDRAZZI,<sup>3</sup> AND LUISA OTTOLINI<sup>4</sup>

<sup>1</sup>Dipartimento Geomineralogico, Università degli Studi di Bari, via E. Orabona 4, I-70125 Bari, Italy
 <sup>2</sup>Dipartimento di Scienze della Terra, Università G.d'Annunzio, via dei Vestini, 3, I-66100 Chieti, Italy
 <sup>3</sup>Dipartimento di Sanita' Pubblica, Sezione di Fisica, Plesso Biotecnologico Integrato, via Volturno 39, I-43100, Parma, Italy
 <sup>4</sup>CNR-Istituto di Geoscienze e Georisorse (IGG), Sezione di Pavia, via Ferrata, 1, I-27100 Pavia, Italy

#### ABSTRACT

Volcanic activity at Mt. Vulture lasted about 750 ka and produced SiO<sub>2</sub>-undersaturated volcanic rocks that can be classified as old (~700 ka), intermediate (~600–550 ka), and young (~130 ka). The intermediate deposits consist of pyroclastic falls and flows and lavas with compositions ranging from phonolite to foidite. A recent revision of the stratigraphic setting allowed these deposits to be classified into one synthem (the Barile Synthem) and further subdivided into four subsynthems (Toppo S. Paolo, Rionero, Vulture-S. Michele, and Ventaruolo). In the present investigation, trioctahedral micas from sample VUT191 in the Vulture-S. Michele Subsynthem are considered. The host rock has modal diopside (20.2%), analcime (22.8%), plagioclase (27.8%), haüyne (5%), phlogopite (8.9%), and magnetite (6.3%). The micas were studied using chemical (EPMA, C-H-N, SIMS), structural (SCXRD), and spectroscopic (Mössbauer) methods.

EPMA of 36 crystals from thin sections and 6 discrete crystals selected for the structural analysis showed remarkable compositional variability, as follows (in wt%):  $SiO_2 = 33.14-38.01$ ,  $Al_2O_3 =$ 15.56-20.45, MgO = 13.02-20.81, FeO<sub>tot</sub> = 6.34-14.08, TiO<sub>2</sub> = 2.34-6.02, K<sub>2</sub>O = 6.03-9.48, Na<sub>2</sub>O = 0.50-0.78, and BaO = 0.89-4.06; all crystals proved to be phlogopite. Elemental C-H-N analyses yielded  $H_2O = 2.86 \pm 0.36$  wt%. The water content was also determined by SIMS on two single crystals, labeled VUT191 2 and VUT191 19, which yielded values of  $3.81 \pm 0.12$  and  $1.72 \pm 0.08$  wt% H<sub>2</sub>O, respectively. Mössbauer investigation showed that all the iron in VUT191 mica is octahedral with  $Fe^{2+} = 25.5\%$  and  $Fe^{3+} = 74.5\%$ , confirming that Vulture micas are particularly  $Fe^{3+}$ -rich, as also found from previous investigations. Structure refinements using anisotropic displacement parameters were performed in space group C2/m and converged at  $1.89 \le R \le 3.17$ ,  $2.09 \le R_w \le 3.43\%$ . All of the analyzed micas belong to the 1M polytype but exhibit remarkable variations in the c parameter from 10.1569(4) to 10.2458(4) Å. The chemical and structural parameters indicate that the studied micas can be divided into two groups: the first encompassing strongly dehydrogenated micas affected mainly by Ti-oxy  $[^{VI}M^{2+} + 2(OH)^- \leftrightarrow ^{VI}Ti^{4+} + 2O^{2-} + H_2]$  and  $M^{3+}$ -oxy  $[^{VI}M^{2+} + (OH)^- \leftrightarrow ^{VI}M^{3+} + O^{2-} + \frac{1}{2}H_2]$ , with  $M^{3+} = Fe^{3+}$ ,  $A^{3+}$ ] substitutions. The second group consist of samples in which vacancy-bearing mechanisms,  $2^{VI}M^{2+} \leftrightarrow {}^{VI}Ti^{4+} + {}^{VI}\Box$  and  $3^{VI}M^{2+} \leftrightarrow 2^{VI}M^{3+} + {}^{VI}\Box$  occur.

**Keywords:** Volcanic phlogopite, Mössbauer spectroscopy, structure refinement, CHN, SIMS, crystal chemistry, substitution mechanisms

## INTRODUCTION

Phlogopite and biotite are common phases in many volcanic rocks, including andesites, dacites, and rhyolites, as well as potassic and ultrapotassic varieties such as phonolite, foidites, lamprophyres, and lamproites. In principle, these minerals can be used to infer pre-eruptive magmatic conditions, provided that complete chemical analysis, including Fe<sup>3+</sup>, Fe<sup>2+</sup>, and H<sub>2</sub>O contents, are available. In addition, mica is an important scavenger of some lithophile elements (LILE) (such as Ba) and high-field strength elements (HFSE) (Ti) that are particular soluble in high K, OH<sup>-</sup>, and CO<sub>2</sub> melts (Chakmouradian 2006). LILE/HFSE and HFSE concentration/dilution relationships are considered crucial in petrological investigations and magma tectonic assignments. In practice, volcanic micas may undergo complex geologic histories and non-equilibrium conditions. This makes it more difficult to develop activity-composition models for solid solutions in volcanic micas as are needed to determine magmatic intensive variables (Feldstein et al. 1996). In addition, the possibility that the structures and crystal chemistry of micas may preserve a record of the geologic processes they underwent has often been neglected in the past. Only in recent works has the

<sup>\*</sup> E-mail: f.scordari@geomin.uniba.it

crystallographic approach been systematically adopted in the study of geologically well-characterized volcanic micas (Brigatti et al. 2005; Scordari et al. 2006; Matarrese et al. 2006a, 2006b; Matarrese 2007). In contrast, the relationship between crystal chemistry and petrogenesis for metamorphic biotites has been well assessed (Henry and Guidotti et al. 2002; Henry et al. 2005; Cesare et al. 2003).

It is known that understanding the crystal chemistry and solid solution of natural micas is quite a challenge because these minerals are affected by multiple cationic and anionic substitutions. Not only does the chemical disorder involve the tetrahedral (Si, Al, Fe, Ti), octahedral (Mg, Mn, Fe<sup>2+</sup>, Fe<sup>3+</sup>, Ti, Al, Cr,  $\Box$ ), interlayer (K, Ba, Ca, Na, NH<sub>4</sub>,  $\Box$ ), and anion (OH<sup>-</sup>, O<sup>2-</sup>, Cl<sup>-</sup>, F<sup>-</sup>) sites, but the same chemical constituents may enter the mica structure according to different heterovalent substitutions. For example, the most likely mechanisms of Ti substitution are (1) 2 <sup>VI</sup>M<sup>2+</sup>  $\leftrightarrow$  <sup>VI</sup>Ti<sup>4+</sup> + <sup>VI</sup> $\Box$ , known as Ti-vacancy (Forbes and Flower 1974); (2) <sup>VI</sup>M<sup>2+</sup> + 2 <sup>IV</sup>Si<sup>4+</sup>  $\leftrightarrow$  <sup>VI</sup>Ti<sup>4+</sup> + 2(OH)<sup>-</sup>  $\leftrightarrow$  <sup>VI</sup>Ti<sup>4+</sup> + 2O<sup>2-</sup> + H<sub>2</sub>, known as Ti-oxy (Bohlen et al. 1980).

Different authors have ascertained that  $Fe^{3+}$ -oxy [<sup>VI</sup>M<sup>2+</sup> + (OH)<sup>-</sup>  $\leftrightarrow$  <sup>VI</sup>Fe<sup>3+</sup> + O<sup>2-</sup> + <sup>1</sup>/<sub>2</sub>H<sub>2</sub>] and Ti-oxy substitutions (mechanism 3 above) play a major role both in metamorphic and igneous phlogopites (Virgo and Popp 2000; Righter et al. 2002; Cesare et al. 2003).

The synergy of several independent analytical techniques is required to correctly assess the substitution mechanisms and correct site populations that form the basis from which reliable activity-composition models may be developed. In the present study, the combination of electron probe micro analysis (EPMA), secondary ion mass spectrometry (SIMS), C-H-N analysis, structural (single-crystal X-ray diffraction, SCXRD), and spectroscopic (Mössbauer) analyses has been used to understand the crystal chemistry of trioctahedral micas from foiditic-phonolitic pyroclastics of Monte Vulture (Potenza, Italy).

The explosive-effusive activity of Mt. Vulture lasted about 750 ka and produced a large variety of alkaline, SiO<sub>2</sub>-undersaturated volcanic rocks ranging from carbonatites and melilitites, to foidite, phonolitic-foidite, tephritic-phonolite, and phonolites (Stoppa et al. 2006). Based on their ages (Brocchini et al. 1994), the Vulture products may be classified as old (~700 ka), intermediate (~600–550 ka), and young (~130 ka). Micas are ubiquitous in Mt. Vulture volcanic rocks. According to a recent revision of the stratigraphic setting (Giannandrea et al. 2006) based on Unconformity Bounded Stratigraphic Units (UBSU, Salvador 1987), the intermediate deposits in the central edifice of Monte Vulture consist of four subsynthems (Toppo S. Paolo, Rionero, Vulture-S. Michele, and Ventaruolo) formed by pyroclastic fall and flows and lava products. The four subsynthems are clustered together into the Barile Synthem. The present investigation is focused on the crystal chemistry of trioctahedral micas belonging to this synthem and in particular, originating in the pyroclastic flow deposits, dated  $601 \pm 7$  to  $629.6 \pm 4.7$  ka (Brocchini et al. 1994), belonging to Vulture-S. Michele Subsynthem. The ultimate aim of the investigations on the Mt. Vulture micas is to gain insights into the chemical and structural responses of these minerals to the geologic history of the host rocks.

#### MATERIALS AND METHODS

The micas of the present work were separated from sample VUT191, which represents the juvenile fraction of the Masseria Saraceno pyroclastic flow, Vulture-San Michele Subsynthem. Micas were given the same label as their host rock. The mode of the phenocrysts in the host rock is 20.2% clinopyroxene, 22.8% analcime, 5.0% haüyne, 8.9% mica, 27.8% plagioclase, 6.3% magnetite, 2.5% apatite, and 5.0% others. The rock name should therefore be pheno-tephrite. However, the presence of a very fine-grained matrix and glass suggests discontinuing pheno-modal classification. The fresh, porphyritic-microcrystalline lava, VUT193, associated with the Masseria Saraceno pyroclastic flow, has a mode of 18.8% diopside, 5.8% leucite, 2.4% K-feldspar, 34.4% anorthoclase, 10.1% haüyne, 8% phlogopite, 6.5% nepheline, and 12.7% plagioclase, along with accessory minerals such as apatite and spinel (<1% each). It is a tephri-phonolite (see Stoppa et al. 2006 for more details). The subhedral mica in VUT191 is not fractured. It contains apatite inclusions and does not show notable deformation features.

Mica separates were obtained by electrostatic separation followed by hand picking. The purity of the concentrates was estimated optically by a reflection microscope to be 99%.

# EPMA

Major-elements compositions were determined on 36 crystals from polished thin sections and on six discrete crystals selected from the concentrates and embedded in epoxy resin. The latter crystals were also used for structural analysis.

Chemical compositions of samples in thin sections were measured with a Cameca SX50 in full WDS mode. Analyses were performed at the Natural History Museum (EMMA facilities) of London. Operating conditions were 15 kV, ~20 nA, and 5  $\mu$ m beam diameter. The standards used were jadeite (Na, Si), periclase (Mg), wollastonite (Ca), rutile (Ti), aluminum oxide (Al), iron oxide (Fe), potassium bromide (K), barium fluoride (Ba), rubidium manganese fluoride (F), halite (Cl), nickel metal (Ni), cromite (Cr), manganese metal (Mn), apatite (P), and celestine (Sr).

The compositions of the six single crystals that were used for structure refinement were measured by wavelength-dispersive spectrometry using a Cameca SX-50 electron microprobe at the Istituto di Geologia Ambientale e Geoingegneria, CNR, Rome, with the following operating conditions: 15 kV accelerating voltage, 15 nA specimen beam current, and 10  $\mu$ m beam diameter. The following standards were employed: jadeite (Na), periclase (Mg), wollastonite (Si and Ca), rutile (Ti), corundum (Al), magnetite (Fe), orthoclase (K), barite (Ba), fluor-phlogopite (F), sylvite (Cl). A conversion from X-ray counts to oxide weight percentages (wt%) was made with the PAP procedure (Pouchou and Pichoir 1985).

The chemical composition of sample VUT191 bulk rock was determined by multi-method analysis (combination of X-ray Fluorescence, Titration, Atomic Absorption Spectrometry) at the SGS Canada INC Mineral Services, Toronto.

#### C-H-N

Bulk C, H, and N analysis of mica concentrates was performed with the EA 1108 elemental analyser of CE Instruments at CNR-IGG, Padova, calibrated with standard Acetanilide ( $C_8H_9NO$ ). The analysis was carried out on ~30 mg of powdered sample.

# SIMS

SIMS measurements were performed with a CAMECA IMS 4f ion microprobe installed at the CNR-IGG at Pavia, Italy. A -12.5 kV accelerated 16O- primary ion beam was used having a current intensity of 3-7 nA and a diameter of <5-10 µm beam, following procedures similar to those adopted by L.O. in the paper by Mesto et al. (2006). Sample mounts and standards were left to de-gas overnight in the ion-microprobe sample chamber. Secondary ion signals of the following isotopes were monitored at the electron multiplier: 1H+, 7Li+, 19F+, and 30Si+ (the latter was used as an inner reference for the matrix). Acquisition times were 20 s (H), 10 s (Li), 50 s (F), and 15 s (Si) over 5 analytical cycles. Detection of positive secondary ions having kinetic energies in the range of 75-125 eV was obtained under steady-state sputtering conditions after 15 min of pre-sputtering. According to previous SIMS work on light elements in silicates, the analysis of "filtered" secondary ions is useful in reducing most chemical matrix effects and in improving the overall measurement reproducibility (Ottolini et al. 1993, 1995 and references therein). Several medium-silica silicate standards were employed for the quantification of the ion signals for H, Li, and F: Finero phlogopite (SiO2 = 40.04 wt%, 4.20 wt% H<sub>2</sub>O, 712 ppm F), kornerupine no.6, schorl no.16, dravite

no.18, and elbaite no.19 (from Ottolini et al. 2002). In particular, we adopted empirical corrections to the ion yield (IY) (H/Si) to consider the variation of the ion signals with increasing (Fe + Mn) content in the sample, as fully described in Ottolini and Hawthorne (2001) and Ottolini et al. (2002). The sample mounts were then re-polished smoothly, carbon coated, and investigated with the electron microprobe proximal to the SIMS craters. These EMPA data were then used in the final SIMS quantification procedures.

#### Mössbauer spectroscopy

The sample for Mössbauer investigation was prepared by crushing the mica under acetone to minimize atmospheric oxidation and the powdered material was dispersed in a 0.5 inch diameter holder. Transmission <sup>57</sup>Fe Mössbauer spectra were obtained on 20 mg of powdered sample at room temperature (RT) using a <sup>57</sup>Co thin source in a Rh matrix. The transducer was employed in constant acceleration mode over a Doppler velocity range of ±4 mm/s. Data were acquired on 512 channels and folded to give a flat background and a zero velocity position corresponding to the center shift (CS, also called isomer shift) of metallic  $\alpha$ -Fe at RT. The spectrum was fitted using the Voigt-based Quadrupole Splitting Distribution (QSD) method developed by Rancourt and Ping (1991) and implemented in the software RECOIL (Lagarec and Rancourt 1997, 1998). This method assumes that the QSD for each generalized site is composed of a number (*N*) of Gaussian components. The number of Gaussian components, but has to ensure that the true QSD of the spectral contribution is well-represented by the sum of Gaussians (Rancourt et al. 1994a, 1994b).

#### SCXRD

X-ray diffraction (XRD) data were collected from single crystals selected from the mica concentrates, using a Bruker AXS X8 APEX II CCD automated diffractometer equipped with a four-circle Kappa goniometer and graphitemonochromatized MoKa radiation (50 kV and 30 mA operating conditions). Several sets of frames were recorded with a crystal-to-detector distance of 40 mm and a strategy optimized by the APEX suite program (Bruker 2003b). A set of 12 frames was used for initial cell determination, whereas complete data collection was accomplished by several  $\phi$  and  $\omega$  scans with 0.5° rotation and 10 s exposure time per frame. The whole Ewald sphere  $(\pm h, \pm k, \pm l)$  was recorded in  $\theta$  ranges up to ~40°. Reflection intensities were extracted and corrected for Lorentz-polarization using the SAINT program (Bruker 2003a). Absorption correction was applied using SADABS (Sheldrick 2003). Least-squares refinements were performed using the program CRYSTALS (Betteridge et al. 2003) in the space group C2/m. Starting atomic coordinates were taken from Mesto et al. (2006). The refined parameters were scale factors, atomic positions, cation occupancy factors, and anisotropic displacement parameters. Ionized X-ray scattering curves were employed for non-tetrahedral cations, whereas ionized vs. neutral species were used for Si and O (Hawthorne et al. 1995). At octahedral sites, the electron density was fitted by varying Fe2+ vs. Mg2+ occupancies with full occupancy constraints, whereas for interlayer and tetrahedral sites, the use of restraints (Watkin 1994) allowed the occupancy to assume values greater than or less than 1. This procedure accounts for interlayer and tetrahedral substitutions and provides a better fit to the electron density, particularly at tetrahedral sites.

### **RESULTS AND DISCUSSION**

### **Chemical analyses**

**Bulk-rock analysis.** The chemical composition of sample VUT191 bulk rock is (in oxide wt%): 48.07% SiO<sub>2</sub>, 1.04% TiO<sub>2</sub>, 18.15% Al<sub>2</sub>O<sub>3</sub>, 4.24% Fe<sub>2</sub>O<sub>3</sub>, 3.09% FeO, 0.21% MnO, 4.54% MgO, 9.51% CaO, 4.38% Na<sub>2</sub>O, 1.07% K<sub>2</sub>O, 0.94% P<sub>2</sub>O<sub>5</sub>, and 3.91% loss on ignition (L.O.I), with a sum of 99.15%. Sample VUT191 is thus an SiO<sub>2</sub>-undersaturated, CaO-rich, alkaline rock with a sodic character having Na<sub>2</sub>O > K<sub>2</sub>O (Irvine and Bargar 1971). Its high  $X_{Mg}$  [ $X_{Mg}$  = Mg/(Mg + Fe) ~72] indicates that it is still a primitive rock. High L.O.I. prevents the use of the TAS diagram, but according to the De La Roche diagram (De La Roche et al. 1980), the rock is a "trachy-basalt" that clearly contrasts with its pheno-crystals modal content and associated rocks. Inconsistent chemical classification is explained by the

remarkable content of analcime that may replace foids. The associated lava VUT193 is chemically a phonolitic-tephrite (according de La Roche's R1-R2 diagram). Its modal composition is tephritic-phonolite, and may represent an evolved rock derived from VUT191. Thus, the VUT 191 rock sample may be related to a relatively primitive liquid (tephri-foidite) that was able to evolve in composition toward tephritic-phonolitc, possibly through clinopyroxene crystal settling in a magma chamber and an increase in feldspar content.

**Micas analyses.** The results of electron microprobe analysis are reported on Table 1 for micas from thin section and in Tables 2 and 3, for mica single crystals that were used structure refinement. The latter analyses represent averages of 5–10 spots per crystal. Considering the whole data set, VUT191 phlogopite is characterized by remarkable chemical inhomogeneity, as revealed by the generally large ranges in oxide wt%: 33.14-38.01 for SiO<sub>2</sub>, 15.56-20.45 for Al<sub>2</sub>O<sub>3</sub>, 13.02-20.81 for MgO, 6.34-4.08 for FeO<sub>tot</sub>, 2.34-6.02 for TiO<sub>2</sub>, 6.03-9.48 for K<sub>2</sub>O, 0.50-0.78 for Na<sub>2</sub>O, and 0.89-4.06 for BaO.

The H<sub>2</sub>O content determined from C-H-N analysis is  $2.86 \pm 0.36$  wt%. This value is the average of two different measurements (that yielded 3.11 and 2.60 wt%) performed on different aliquots of the same mica sample. This result suggests a degree of inhomogeneity of OH concentration, which also has been documented for other Mt. Vulture phlogopites (Mesto et al. 2006). No nitrogen was detected in the analyzed samples, whereas the carbon content was  $0.18 \pm 0.02$  wt% C.

SIMS measurements were performed on two out of the six single crystals used for structure refinement (VUT191\_2 and VUT191\_19), selected on the basis of their different chemistries and *c*-parameters (see below). The grain to grain chemical variability noted above extends to light elements (H, Li, F, see Tables 2 and 3).

All considered, the analyzed samples are solid solutions between phlogopite  $[KMg_3AlSi_3O_{10}(OH)_2]$  and annite  $[KFe_3^2+AlSi_3O_{10}(OH)_2]$  with Ti 0.10–0.38 apfu, Ba 0.03–0.12 apfu, and a variable degree of dehydrogenation. The atomic ratios for data from single crystals, as listed in Table 3, were calculated on the basis on 12(O, OH, Cl, F), assuming all Ti as Ti<sup>4+</sup> and using the Fe<sup>2+</sup>/Fe<sup>3+</sup> ratios provided by the Mössbauer investigation (see below). Final crystal-chemical formulae and a critical discussion of data in Table 3 are postponed to the section on crystal chemistry below.

Figure 1 is a plot of the annite component  $[Fe_{tot} + Mg)]$ , vs. the Ti content (apfu) for the samples of the present study. For comparison, data from other well-characterized micas from Mt. Vulture, all from the Barile Synthem but from different subsynthems, are also shown. Unlike micas belonging to other subsynthems (such as SA sample, from Rionero Subsynthem as discussed in Scordari et al. 2006) and the VUT187 sample from the Ventaruolo Subsynthem, the VUT191 phlogopite has an annite component that spans almost the whole interval of variation (0.15–0.40) found so far for Monte Vulture micas (Matarrese et al. 2006, 2007).

# Mössbauer spectroscopy

The results of the Mössbauer investigation are reported in Table 4 and typical Mössbauer spectra of the micas are displayed in Figure 2. Three absorption bands occur at approximately -0.2,

TABLE 1. Electron microprobe data (wt%) for VUT191 phlogopites in thin sections

				(										
Sample	191_1	191_3	191_4	191_5	191_6	191_7	191_8	191_9	191_10	191_12	191_14	191_15	191_16	191_17
SiO <sub>2</sub>	37.416	36.616	35.596	33.848	33.574	34.105	33.805	36.706	36.868	37.459	34.291	33.257	33.138	36.293
Al <sub>2</sub> O <sub>3</sub>	17.207	17.430	16.808	15.767	15.909	15.883	15.898	16.606	17.713	16.801	16.308	15.690	15.563	17.534
MgO	19.623	18.148	15.619	14.556	14.550	14.818	14.445	20.219	18.889	19.872	14.369	14.065	13.434	18.149
FeO <sub>tot</sub>	6.337	8.268	12.133	12.835	12.525	12.864	12.960	7.649	7.432	6.917	12.954	12.997	13.445	8.392
TiO <sub>2</sub>	2.344	3.022	3.876	5.721	5.539	5.464	5.816	2.891	2.585	2.539	5.503	6.022	5.706	2.951
$Cr_2O_3$	0.140	0.000	0.000	0.000	0.000	0.000	0.000	0.092	0.023	0.067	0.026	0.007	0.018	0.094
MnO	0.017	0.035	0.179	0.231	0.245	0.192	0.152	0.037	0.022	0.045	0.205	0.178	0.209	0.070
K₂O	9.009	8.760	9.154	8.298	8.201	8.357	8.155	9.404	8.916	9.140	8.772	8.014	7.417	8.819
Na₂O	0.538	0.656	0.594	0.569	0.597	0.567	0.541	0.604	0.613	0.603	0.569	0.527	0.677	0.616
BaO	0.909	1.207	1.278	3.680	3.520	3.306	3.564	1.106	0.933	0.972	1.459	4.058	3.472	1.282
CaO	0.194	0.115	0.050	0.038	0.057	0.069	0.105	0.127	0.206	0.173	0.137	0.153	0.554	0.081
F	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Cl	0.039	0.016	0.072	0.136	0.112	0.076	0.098	0.017	0.023	0.027	0.095	0.103	0.094	0.040
Sum	94.30	94.96	96.367	96.745	95.86	96.77	96.616	96.106	94.893	95.19	95.764	96.151	94.844	95.018
FeO*	1.606	2.096	3.076	3.254	3.175	3.261	3.285	1.939	1.884	1.753	3.284	3.295	3.408	2.127
Fe <sub>2</sub> O <sub>3</sub> *	5.257	6.859	10.065	10.648	10.391	10.672	10.751	6.345	6.165	5.738	10.746	10.782	11.154	6.962
	101 10	101 20	101 21	101 22	101 23	101 24	101 25	101 26	101 27	101 28	101 20	101 30	101 33	101 3/
SIO	20 011	26 / 20	26.062	24.540	24.400	22.050	24.607	22 655	22 226	27.061	26 100	22 004	25 021	22 /65
	20 447	16 082	18 052	16 1/17	16 100	15 820	16 100	15 702	15 7/1	17 031	17 670	16 /08	17 230	15 888
MaO	20.447	10.902	10.032	14 014	1/ 205	14.240	15.025	14 775	14666	10 101	1/.0/9	12 021	16 251	11,000
FoO	7 8 2 7	7 701	8 0/18	12 018	14.303	12 606	12 212	14.773	12 027	8 268	14.313	1/ 070	11 05 8	12/125
	2 560	2 1/12	2 060	5 449	5 671	5 650	5 1 8 6	6.003	5 5/18	2 081	3 /13	5 308	3 110	5 608
	0.012	0.051	0.000	0.000	0.042	0.032	0.000	0.000	0.000	0.060	0.070	0.063	0.000	0.037
MnO	0.012	0.051	0.000	0.000	0.042	0.052	0.000	0.000	0.000	0.000	0.070	0.005	0.000	0.037
K.O	7 477	8 999	8 642	8 862	8 1 2 1	8 3 3 6	8 74 2	8 112	8 313	8 542	6.033	8 245	8 944	8 1 5 6
Na-O	0.569	0.656	0.660	0.549	0.121	0.550	0.742	0.112	0.543	0.542	0.033	0.245	0.534	0.150
BaO	1 073	1 242	1 284	2 039	3 2 2 8	3 343	1 901	3 873	3 176	1 310	0.705	2 981	1 1 2 4	3 309
CaO	0 189	0.071	0 118	0.077	0 108	0 1 1 3	0.095	0 143	0 162	0 164	2 332	0.090	0 144	0.078
F	n d	n d	n d	n d	nd	nd	n d	n d	n d	n d	n d	n d	nd	n d
Cl	0.027	0.060	0.028	0.097	0.107	0.100	0.104	0.090	0.091	0.037	0.086	0.079	0.087	0.097
Sum	94 511	95 39	95 533	96.887	96.654	96.048	95.858	97.056	95.673	95,801	96.301	96.114	96,386	95 236
FeO*	1.984	1.975	2 040	3,275	3,222	3,218	3,121	3,225	3 277	2 096	3,297	3,569	3.031	3,152
Fe <sub>2</sub> O <sub>3</sub> *	6.493	6.463	6.677	10.717	10.543	10.532	10.215	10.555	10.724	6.859	10.789	11.680	9.920	10.316
	191_35	191_36	191_37	191_38	191_39	191_40	191_204	191_214						
SIO <sub>2</sub>	35.630	33.568	33.991	36.511	36.607	36.552	34.845	37.331						
AI <sub>2</sub> O <sub>3</sub>	16.404	15.858	16.088	16.682	16.612	16.910	15.801	17.007						
MgO	17.834	14./65	14.243	20.663	20.810	18./54	15.526	19.9/6						
FeO <sub>tot</sub>	11.098	13.379	13.210	6.632	6./60	7.725	12.542	8.367						
	3.099	5.226	5.406	2.804	2.809	3.049	4.435	3.238						
$Cr_2O_3$	0.004	0.000	0.000	0.056	0.000	0.009	0.000	0.091						
MINU	0.129	0.213	0.201	0.090	0.084	0.008	0.183	0.059						
K <sub>2</sub> O	9.298	8.634	8.318	9.217	9.411	9.028	9.480	9.119						
Na <sub>2</sub> O	0.543	0.566	0.554	0.567	0.578	0.600	0.576	0.581						
BaO C=O	1.103	3.207	3.281	1.154	1.185	1.351	0.012	0.067						
	0.007	0.050	0.067	0.064	0.087	0.119	0.015	0.007						
	n.a.	n.a.	n.a.	n.a.	n.a.	n.a	n.a	n.a						
Sum	0.049	06 600	0.007	0.004	0.02/	0.05/	0.050	0.010						
	20.24	2 202	2 2 4 4	1 6 0 1	55.55∠ 1 714	74./04 1.050	2 170	20.341						
Fe <sub>0</sub> 0.*	2.013	5.592 11.090	5.549 10.959	5 502	5.608	6 4 0 9	5.179 10.405	2.121 6.941						
Noto: n d	- not dot	arminod	10.739	5.502	5.000	0.709	10.405	0.71						
* From N	. – not det lössbauer	spectrosco	ру.											
	-													

1.0, and 2.3 mm/s that correspond respectively to (1) the sum of low-energy lines of <sup>VI</sup>Fe<sup>2+</sup> and <sup>VI</sup>Fe<sup>3+</sup> doublets; (2) the high-energy line of the <sup>VI</sup>Fe<sup>3+</sup> doublet; and (3) the high-energy component of the <sup>VI</sup>Fe<sup>2+</sup> doublet. No shoulder at ~0.5 mm/s (which would be evidence for the occurrence of tetrahedral Fe<sup>3+</sup>; see Redhammer et al. 2005) was observed. In Table 4, the results of  $n_1$ - $n_2$  fittings are compiled, where  $n_1$  and  $n_2$  are the numbers of assumed QSD Gaussian components for <sup>VI</sup>Fe<sup>2+</sup> VIFe<sup>3+</sup> sites, respectively. For the sample analyzed here,  $n_1 = n_2 = 1$ . A<sup>-/</sup>A<sup>+</sup> ratios were allowed to vary during fitting (see Table 4) to account for the possibility that the orientation of crystallites in the analyzed powder is not completely random (Rancourt et al. 1994b).

The fitting parameters are consistent with those reported in the literature for synthetic and natural trioctahedral micas (Redhammer 1998; Rancourt et al. 1993, 1994a; Shabani 1999; Dyar 2002, Redhammer et al. 2005). The sample is Fe<sup>3+</sup>-rich, having Fe<sup>2+</sup>  $\approx$  25% and Fe<sup>3+</sup>  $\approx$  75%.

Previous investigations have shown that Mt. Vulture micas are characterized by a variable but nevertheless high ratio of Fe<sup>3+</sup>/ Fe<sub>tot</sub>, ranging from 42 to 88% (Matarrese et al. 2005, 2006, 2007; Scordari et al. 2006; Schingaro et al. 2006; Matarrese 2007). Focusing only on micas from intermediate Vulture volcanics, Scordari et al. (2006) have found 44% Fe<sup>2+</sup> and 56% Fe<sup>3+</sup> in their SA sample (Rionero Subsynthem), whereas Schingaro et al. (in review) studied a phlogopite from the Ventaruolo Subsynthem that has 13% Fe<sup>2+</sup> and 87% Fe<sup>3+</sup>. However, the typical Fe<sup>3+</sup>/Fe<sub>tot</sub> for Mt. Vulture micas is ~50% (Matarrese 2007), and in all cases where an Fe<sup>3+</sup>/Fe<sub>tot</sub> ratio >> 50% has been found, late- and/or post-magmatic events can be invoked, which affected the original Fe<sup>3+</sup> contents.

			ci accare i			
Sample	VUT191	VUT191	VUT191	VUT191	VUT 191	VUT191
name	_1	_2	_10	_11	_13	_19
SiO <sub>2</sub>	37.036	37.198	33.600	33.980	35.078	34.360
$AI_2O_3$	17.098	16.354	15.879	15.730	16.343	15.969
MgO	20.238	19.404	15.586	14.716	14.677	14.699
FeO <sub>tot</sub>	6.620	6.800	11.599	11.682	11.696	12.214
TiO <sub>2</sub>	2.405	2.640	4.623	4.654	4.579	5.063
$Cr_2O_3$	0.147	0.053	0.007	0.003	0.011	0.007
MnO	0.046	0.042	0.124	0.178	0.210	0.175
NiO	0.022	0.041	0.015	0.028	0.025	0.020
Li₂O	-	0.009	-	-	-	0.027
K₂O	8.488	8.749	8.088	7.749	7.851	7.877
Na <sub>2</sub> O	0.608	0.600	0.592	0.599	0.631	0.565
BaO	1.035	0.971	2.398	2.976	2.474	2.836
CaO	0.011	0.012	0.026	0.075	0.033	0.021
F	0.415	0.640‡	0.621	0.945	0.884	1.280‡
CI	0.024	0.037	0.084	0.088	0.111	0.09
H₂O	2.860†	3.810‡	2.86†	2.86†	2.86†	1.720‡
Total	97.683	96.461	95.947	95.635	96.865	95.829
FeO*	1.678	1.718	2.940	2.979	2.982	3.115
$Fe_2O_3^*$	5.492	5.622	9.623	8.703	9.683	10.112

 TABLE 2.
 Average electron microprobe data (wt%) for VUT191 single

 crystals used for structure refinement

\* Mössbauer results used to recast FeO<sub>tot</sub> into FeO and Fe<sub>2</sub>O<sub>3</sub>.

† Average value from C-H-N analysis (see text).

‡ From SIMS.



**FIGURE 1.** A plot of Fe/(Fe + Mg), thought to represent the amount of the annite component vs. Ti content for micas from Monte Vulture intermediate pyroclastics (Barile Synthem). Diamonds = Barred symbols are VUT191 phlogopites from thin section, this work; full symbols are single crystals of VUT191 phlogopite, this work; open triangles = VUT187 phlogopite, Schingaro et al. (in review); open circles = SA1 phlogopite single crystals data from Scordari et al. (2006).

### Structural features

Details on data collection, lattice parameters, results of structure refinements, fractional atomic coordinates, partial occupancies, anisotropic, and equivalent isotropic refinements are given in Tables 5 and 6. Tables 7 and 8 contain selected bond lengths and distortion parameters describing the polyhedral and layer geometry for the studied samples. Atomic labeling is the same as in Hazen and Burnham (1973). The refinements performed in space group C2/m converged at  $1.89 \le R \le 3.17$ ,  $2.09 \le R_w \le$ 3.43%. The difference Fourier maps revealed hydrogen in the expected position for all samples, but extra peaks were observed for samples VUT191\_10 and VUT191\_11. In particular for sample VUT191\_10, the initial structure refinement converged

Sample	VUT 191	VUT 191	VUT191	VUT191	VUT191	VUT191
name	_1	_2	_10	_11	_13	_19
Si	2.735	2.715	2.584	2.632	2.652	2.658
™AI	1.265	1.285	1.368	1.294	1.348	1.342
ΣΤ	4.000	4.000	4.000	4.000	4.000	4.000
<sup>VI</sup> AI	0.223	0.122	0.068	0.182	0.108	0.114
Mg	2.228	2.112	1.699	1.747	1.654	1.695
Fe <sup>+2</sup>	0.104	0.105	0.193	0.197	0.189	0.202
Fe <sup>+3</sup>	0.305	0.310	0.507	0.581	0.551	0.589
Ti	0.134	0.145	0.271	0.279	0.260	0.295
Cr	0.019	0.007	0.000	0.000	0.001	0.001
Mn	0.003	0.003	0.012	0.012	0.001	0.011
Ni	0.001	0.002	0.001	0.002	0.002	0.001
Li	-	0.003	-	-	-	0.008
$\Sigma$ Oct	3.015	2.809	2.832	2.751	2.764	2.916
K	0.800	0.815	0.766	0.787	0.757	0.777
Na	0.087	0.085	0.090	0.093	0.092	0.085
Ba	0.030	0.028	0.090	0.093	0.073	0.086
Ca	0.001	0.001	0.000	0.006	0.003	0.002
ΣΙ	0.917	0.927	0.946	0.979	0.923	0.948
OH⁻	1.410	1.857	1.479	0.670	1.443	0.888
F	0.097	0.148	0.232	0.238	0.211	0.313
Cl	0.003	0.005	0.012	0.012	0.014	0.012
0	0.490	0.000	0.278	1.080	0.331	0.787

TABLE 3. Atomic proportions (apfu) as determined from data in Table 2



FIGURE 2. Room-temperature Mössbauer spectrum of VUT191 phlogopite.

at R = 7.49,  $R_w = 7.88\%$  and the residual electron density gave  $\rho_{min} = -0.75$ , and  $\rho_{max} = 7.24 \text{ e}^{-/}\text{Å}^3$ . The same parameters for sample VUT191\_11 were R = 4.30,  $R_w = 4.75\%$ ,  $\rho_{min} = -0.45$ , and  $\rho_{max} = 2.68 \text{ e}^{-/}\text{Å}^3$ . Extra peaks in difference Fourier syntheses apparently pointed to  $\pm b/3$  shifts of the T and K atomic positions (Schingaro et al. 2001). It turned out that these samples were affected by the Ďurovič effect (Nespolo and Ferraris 2001), so that the large residues in electron density maps could be ascribed to systematic wrong measurements of non-family reflections. In subsequent refinements, different scales were allowed to vary for reflections *hkl* with k = 3n and *hkl* with  $k \neq 3n$ , which refined to 0.563(1), 0.437(1) and 0.533(1), 0.467(1) for VUT191\_10, and VUT191\_11, respectively.

	iossouuci p	arameters obtai		ang method					
	χ²	Species	Γ/2	$\delta_0^*$	A-/A+	$\Delta E_Q$	σ	P (%)	A (%)
			(mm/s)	(mm/s)		(mm/s)	(mm/s)		
VUT191	1.75	Fe <sup>2+</sup>	0.097*	1.083(10)	1†	2.333(21)	0.338(16)	100*	25.5(6)
		Fe <sup>3+</sup>		0.3987(57)	0.9147(89)	1.151(10)	0.5197(82)	100*	74.5(6)
$* \delta_1 = 0.$									
+ Fixed valu	ie.								

TABLE 4. Mössbauer	parameters obtained b	v C	)SD	fitting	method

#### TABLE 5. Crystal, experimental, and refinement data for VUT191 phlogopite

	VUT191_1	VUT191_2	VUT191_10	VUT191_11	VUT191_13	VUT191_19
Space group	C2/m	C2/m	C2/m	C2/m	C2/m	C2/m
a (Å)	5.3252(2)	5.3254(4)	5.3380(2)	5.3367(3)	5.3377(2)	5.3341(2)
b (Å)	9.2205(3)	9.2216(7)	9.2430(3)	9.2431(5)	9.2424(4)	9.2387(3)
<i>c</i> (Å)	10.2458(4)	10.2411(9)	10.1923(3)	10.1806(5)	10.1654(4)	10.1569(4)
β (°)	99.9938(26)	100.0223(64)	100.0814(21)	100.1033(42)	100.0990(32)	100.0597(27)
Cell volume (Å <sup>3</sup> )	495.442	495.255	495.116	494.404	493.721	492.836
Ζ	2	2	2	2	2	2
Crystal size (mm)	$0.17 \times 0.31 \times 0.03$	$0.27 \times 0.25 \times 0.01$	$0.37 \times 0.33 \times 0.02$	$0.22\times0.29\times0.02$	$0.28 \times 0.14 \times 0.03$	0.27 × 0.24 × 0.01
$\theta$ range for data collection	2-31°	2-39°	2-36°	2-36°	2-44°	4-36°
Reflections measured /unique	11606/836	23889/1635	1257/1257	1258/1258	8468/1944	6972/1256
	$[R_{int} = 0.039]$	$[R_{int} = 0.054]$	$[R_{int} = 0.053]$	$[R_{int} = 0.047]$	$[R_{int} = 0.035]$	$[R_{int} = 0.031]$
Reflections used $[l > 3\sigma(l)]$	438	446	729	517	711	612
No. of refined parameters	71	71	73	73	71	71
Goof*	0.83	0.83	0.43	0.62	0.69	0.75
<i>R</i> <sub>1</sub> †	3.17	3.14	1.89	2.33	2.67	2.77
Rw†	3.43	3.29	2.09	2.46	2.94	3.14
$\Delta \rho_{min}, \Delta \rho_{max}$ (e/Å <sup>3</sup> )	-0.21, 0.44	-0.20, 0.41	-0.17, 0.24	-0.21, 0.20	-0.22, 0.29	-0.16, 0.38
K C L C C C C C C C C C C C C C C C C C		1.0	<i>c a i i</i>			

\* Goodness-of-fit =  $\left[\Sigma[w(F_o^2 - F_c^2)^2]/(N - P)\right]^{1/2}$ . where N and P are the number of reflections and parameters, respectively. +  $R_1 = \Sigma[[F_o] - [F_c]]/\Sigma[F_o]; R_w = [\Sigma[w([F_o] - [F_c])^2]/\Sigma w[F_o]^2]^{1/2}$ .

 TABLE 6.
 Results of structure refinement in space group C2/m: crystallographic coordinates, equivalent isotropic displacement parameters

 (Å<sup>2</sup>), partial occupancies and anisotropic displacement parameters

Site	Atom	x/a	v/h	7/0	11	Occupancy	11.	11	11.	11	11.	11.
Jite	Atom	x/u	y/0	2/1	U <sub>iso/equiv</sub>	Occupancy	011	022	033	023	013	012
VUT1	91_1											
K	K+	0.0000	0.0000	0.0000	0.0342	1.0049(9)	0.0328(12)	0.0334(13)	0.0363(14)	0.0000	0.0060(10)	0.0000
M1	Mg <sup>2+</sup>	0.0000	0.5000	0.5000	0.0123	0.8468(8)	0.0098(11)	0.0077(10)	0.0203(13)	0.0000	0.0052(10)	0.0000
	Fe <sup>2+</sup>	0.0000	0.5000	0.5000	0.0123	0.1532(7)	0.0098(11)	0.0077(10)	0.0203(13)	0.0000	0.0052(10)	0.0000
M2	Mg <sup>2+</sup>	0.0000	0.83464(19)	0.5000	0.0138	0.8319(8)	0.0091(7)	0.0127(7)	0.0196(8)	0.0000	0.0029(6)	0.0000
	Fe <sup>2+</sup>	0.0000	0.83464(19)	0.5000	0.0138	0.1681(8)	0.0091(7)	0.0127(7)	0.0196(8)	0.0000	0.0029(6)	0.0000
Т	Si <sup>4+</sup>	0.57542(19)	0.16675(12)	0.22673(12)	0.0136	0.4137(10)	0.0114(4)	0.0103(4)	0.0193(5)	0.0003(6)	0.0032(4)	0.0000(5)
	Si	0.07542(19)	0.16675(12)	0.22673(12)	0.0136	0.5765(10)	0.0114(4)	0.0103(4)	0.0193(5)	0.0003(6)	0.0032(4)	0.0000(5)
01	0,02-	0.8310(6)	0.2255(4)	0.1694(3)	0.0219	1	0.0189(14)	0.0228(16)	0.0245(16)	-0.0032(14)	0.0051(12)	-0.0039(13)
O2	0,0 <sup>2-</sup>	0.5067(8)	0.0000	0.1702(5)	0.0216	1	0.023(2)	0.016(2)	0.025(2)	0.0000	0.0011(18)	0.0000
O3	0,0 <sup>2-</sup>	0.6306(5)	0.1673(3)	0.3908(3)	0.0146	1	0.0112(12)	0.0131(12)	0.0195(14)	-0.0004(14)	0.0027(10)	-0.0003(12)
04	0,02-	0.1321(8)	0.0000	0.3992(5)	0.0148	1	0.015(2)	0.014(2)	0.016(2)	0.0000	0.0033(18)	0.0000
	Н	0.106(18)	0.0000	0.321(10)	0.0148	0.7607(10)	0.015(2)	0.014(2)	0.016(2)	0.0000	0.0033(18)	0.0000
VI IT 1	01 2											
VUII	91_Z					4 04 00 (0)	0.0075(4.0)	0.0050(40)	0.0454(4.6)		0.0007(4.4)	
K	K <sup>*</sup>	0.0000	0.0000	0.0000	0.0394	1.0188(9)	0.0375(13)	0.0359(13)	0.0454(16)	0.0000	0.0087(11)	0.0000
MI	Mg <sup>2+</sup>	0.0000	0.5000	0.5000	0.01/3	0.8478(8)	0.0136(12)	0.0111(12)	0.0286(16)	0.0000	0.0074(11)	0.0000
	Fe2+	0.0000	0.5000	0.5000	0.0173	0.1521(7)	0.0136(12)	0.0111(12)	0.0286(16)	0.0000	0.0074(11)	0.0000
M2	Mg <sup>2+</sup>	0.0000	0.8360(2)	0.5000	0.0194	0.8109(8)	0.0135(7)	0.0170(8)	0.0282(10)	0.0000	0.0050(7)	0.0000
	Fe <sup>2+</sup>	0.0000	0.8360(2)	0.5000	0.0194	0.1891(8)	0.0135(7)	0.0170(8)	0.0282(10)	0.0000	0.0050(7)	0.0000
Т	Si <sup>4+</sup>	0.57527(19)	0.16678(14)	0.22626(12)	0.0180	0.4483(10)	0.0142(4)	0.0127(5)	0.0279(6)	-0.0005(8)	0.0057(4)	-0.0002(6)
	Si	0.57527(19)	0.16678(14)	0.22626(12)	0.0180	0.5479(10)	0.0142(4)	0.0127(5)	0.0279(6)	-0.0005(8)	0.0057(4)	-0.0002(6)
01	0,02-	0.8315(6)	0.2245(4)	0.1692(3)	0.0271	1	0.0226(14)	0.0287(18)	0.0315(17)	-0.0032(15)	0.0085(13)	-0.0060(13)
02	0,02-	0.5055(9)	0.0000	0.1696(5)	0.0269	1	0.031(3)	0.017(2)	0.031(3)	0.0000	0.002(2)	0.0000
O3	0,0 <sup>2-</sup>	0.6302(5)	0.1675(4)	0.3913(3)	0.0181	1	0.0135(12)	0.0136(13)	0.0280(17)	0.0004(16)	0.0055(11)	0.0002(14)
04	0,02-	0.1337(9)	0.0000	0.4006(5)	0.0181	1	0.016(2)	0.015(2)	0.024(3)	0.0000	0.005(2)	0.0000
	Н	0.103(15)	0.0000	0.327(6)	0.0181	0.9009(10)	0.016(2)	0.015(2)	0.024(3)	0.0000	0.005(2)	0.0000

In the final step of the anisotropic refinements, the hydrogen atom was added to the model. Its coordinates and occupancy were refined (see Table 6), whereas the displacement parameters were made to ride those of the O4 atom. The final O-H distances were in the range 0.74(6)-0.87(5) Å.

The variation of the *c*-parameter, which ranges from 10.1569(4) Å (sample VUT191\_19) to 10.2458(4) Å (sample VUT191\_1, see Table 5) is notable. This behavior is peculiar for micas of the Vulture-S. Michele Subsynthem, because in all

other subsynthems so far investigated, the relevant phlogopites have much smaller ranges of variation (Matarrese et al. 2006; Schingaro et al. 2006; Matarrese 2007). This feature seems to reflect the grain-to-grain variability of chemical composition already highlighted above. However, the *c* parameters observed for the micas of this study are small when compared with those of close-to-end-member annite [10.3235(4) Å, Redhammer and Roth (2002)] and close-to-end-member phlogopite [10.291(4) Å, Alietti et al. (1995); 10.3100(54) Å, synthetic sample Phl no. 2

TABLE 6. —CONTINUED

Site	Aton	n x/a	y/b	z/c	U <sub>iso/equiv</sub>	Occupancy	U <sub>11</sub>	U <sub>22</sub>	U <sub>33</sub>	U <sub>23</sub>	U <sub>13</sub>	U <sub>12</sub>
VUT19	01 10					. ,						
ĸ	K+	0.0000	0.0000	0.0000	0.0292	1,1088(8)	0.0272(4)	0.0277(5)	0.0331(5)	0.0000	0.0061(4)	0.0000
M1	Ma <sup>2+</sup>	0.0000	0.5000	0.5000	0.0115	0.6593(8)	0.0116(4)	0.0072(4)	0.0163(5)	0.0000	0.0043(3)	0.0000
	Fe <sup>2+</sup>	0.0000	0.5000	0.5000	0.0115	0.3407(7)	0.0116(4)	0.0072(4)	0.0163(5)	0.0000	0.0043(3)	0.0000
M2	Mg <sup>2+</sup>	0.0000	0.83813(9)	0.5000	0.0124	0.6424(8)	0.0093(2)	0.0135(3)	0.0142(3)	0.0000	0.00149(19)	0.0000
	Fe <sup>2+</sup>	0.0000	0.83813(9)	0.5000	0.0124	0.3569(7)	0.0093(2)	0.0135(3)	0.0142(3)	0.0000	0.00149(19)	0.0000
Т	Si <sup>4+</sup>	0.57472(8)	0.16706(7)	0.22445(4)	0.0101	0.2912(9)	0.00869(16)	0.00913(17)	0.01244(18)	-0.0003(3)	0.00202(13)	0.0000(2)
	Si	0.57472(8)	0.16706(7)	0.22445(4)	0.0101	0.7013(9)	0.00869(16)	0.00913(17)	0.01244(18)	-0.0003(3)	0.00202(13)	0.0000(2)
01	0,02-	0.8283(3)	0.22651(18)	0.16700(14)	0.0187	1	0.0166(6)	0.0231(7)	0.0170(6)	-0.0028(6)	0.0044(5)	-0.0057(6)
02	0,02-	0.5089(4)	0.0000	0.1682(2)	0.0189	1	0.0239(10)	0.0145(9)	0.0169(9)	0.0000	-0.0008(7)	0.0000
O3	0,0 <sup>2-</sup>	0.6307(2)	0.16822(18)	0.39132(13)	0.0118	1	0.0114(5)	0.0109(5)	0.0134(5)	0.0000(6)	0.0023(4)	0.0003(6)
04	0,02-	0.1326(4)	0.0000	0.3996(2)	0.0123	1	0.0112(8)	0.0131(9)	0.0125(9)	0.0000	0.0018(7)	0.0000
	Н	0.108(9)	0.0000	0.313(5)	0.0123	0.7867(10)	0.0112(8)	0.0131(9)	0.0125(9)	0.0000	0.0018(7)	0.0000
VUITS	/1_11 K+	0.0000	0.0000	0.0000	0 0200	1 1210/0)	0.0257(7)	0.0269(7)	0.0276(0)	0.0000	0.0074(6)	0.0000
N 1	Ma <sup>2+</sup>	0.0000	0.0000	0.0000	0.0290	0.6907(9)	0.0237(7)	0.0208(7)	0.0370(9)	0.0000	0.0074(0)	0.0000
IVII	Eo2+	0.0000	0.5000	0.5000	0.0120	0.0097(0)	0.0119(7)	0.0080(7)	0.0195(9)	0.0000	0.0057(6)	0.0000
MO	Ma <sup>2+</sup>	0.0000	0.3000	0.5000	0.0120	0.5105(7)	0.0119(7)	0.0080(7) 0.0147(5)	0.0193(9)	0.0000	0.0037(0)	0.0000
1012	Fo <sup>2+</sup>	0.0000	0.83933(14)	0.5000	0.0141	0.3542(8)	0.0095(4)	0.0147(5)	0.0182(6)	0.0000	0.0027(4)	0.0000
т	Si <sup>4+</sup>	0.57436(12)	0.05755(14)	0.22395(8)	0.0123	0.53942(0)	0.0094(3)	0.0110(3)	0.0169(4)	0.0002(5)	0.0027(4)	-0.0004(4)
	Si	0.57436(12)	0.16733(11)	0.22395(8)	0.0123	0.4532(10)	0.0094(3)	0.0110(3)	0.0169(4)	0.0002(5)	0.0034(3)	-0.0004(4)
01	0.0 <sup>2-</sup>	0.8277(4)	0.2269(3)	0.1662(2)	0.0210	1	0.0170(9)	0.0241(12)	0.0226(11)	-0.0033(10)	0.0050(8)	-0.0058(9)
02	0,0 <sup>2-</sup>	0.5087(6)	0.0000	0.1673(3)	0.0202	1	0.0256(16)	0.0151(14)	0.0186(16)	0.0000	0.0006(12)	0.0000
O3	0,0 <sup>2-</sup>	0.6308(4)	0.1683(3)	0.3915(2)	0.0143	1	0.0129(9)	0.0130(9)	0.0174(10)	0.0005(12)	0.0034(7)	0.0006(10)
04	0,0 <sup>2-</sup>	0.1331(6)	0.0000	0.4001(4)	0.0137	1	0.0107(14)	0.0154(16)	0.0146(17)	0.0000	0.0014(13)	0.0000
	Н	0.092(15)	0.0000	0.321(6)	0.0137	0.6645(10)	0.0107(14)	0.0154(16)	0.0146(17)	0.0000	0.0014(13)	0.0000
VUT19	1_13	0.0000	0.0000	0.0000	0 0 0 0 0 0 0	1 1 2 2 4 (0)	0.0051(6)	0.0050(6)	0.02(7(0)	0.0000	0.0050(5)	0.0000
K M1	K <sup>+</sup>	0.0000	0.0000	0.0000	0.0292	1.1294(9)	0.0251(6)	0.0259(6)	0.0367(8)	0.0000	0.0058(5)	0.0000
IVII	To2+	0.0000	0.5000	0.5000	0.0107	0.7072(8)	0.0062(6)	0.0078(6)	0.0169(7)	0.0000	0.0045(5)	0.0000
MO	re Ma <sup>2+</sup>	0.0000	0.5000	0.5000	0.0107	0.2927(7)	0.0062(0)	0.0078(8)	0.0169(7)	0.0000	0.0045(5)	0.0000
IVIZ	Fo <sup>2+</sup>	0.0000	0.84067(11)	0.5000	0.0129	0.3598(8)	0.0008(3)	0.0159(4)	0.0160(4)	0.0000	0.0017(3)	0.0000
т	Si <sup>4+</sup>	0.57405(12)	0.04007(11)	0.22365(7)	0.0122	0.3577(10)	0.0000(3)	0.0102(3)	0.0141(3)	0.0001(4)	0.0017(3)	-0.0001(3)
	Si	0.57405(12)	0.16701(9)	0.22365(7)	0.0112	0.6352(10)	0.0092(3)	0.0102(3)	0.0141(3)	0.0001(4)	0.0020(2)	-0.0001(3)
01	0.0 <sup>2-</sup>	0.8267(4)	0.2278(3)	0.1657(2)	0.0202	1	0.0178(9)	0.0247(11)	0.0189(9)	-0.0039(9)	0.0049(7)	-0.0066(8)
02	0,0 <sup>2-</sup>	0.5103(6)	0.0000	0.1676(3)	0.0207	1	0.0248(15)	0.0136(12)	0.0215(15)	0.0000	-0.0021(12)	0.0000
03	0,0 <sup>2-</sup>	0.6310(3)	0.1689(2)	0.39121(19)	0.0119	1	0.0106(7)	0.0111(7)	0.0140(8)	-0.0006(8)	0.0018(6)	0.0008(7)
04	0,0 <sup>2-</sup>	0.1326(6)	0.0000	0.4004(3)	0.0135	1	0.0122(12)	0.0144(13)	0.0143(14)	0.0000	0.0032(11)	0.0000
	Н	0.105(14)	0.0000	0.322(8)	0.0135	0.6826(10)	0.0122(12)	0.0144(13)	0.0143(14)	0.0000	0.0032(11)	0.0000
	יז_19 ⊮+	0.0000	0.0000	0.0000	0 0 2 9 7	1 1206/01	0 0252/7)	0 0245(7)	0.0265(10)	0.0000	0.0061(6)	0.0000
м1	Γ. Μα <sup>2+</sup>	0.0000	0.0000	0.0000	0.028/	1.1200(9)	0.0233(7)	0.0245(7)	0.0305(10)	0.0000	0.0001(0)	0.0000
1011	Eo2+	0.0000	0.5000	0.5000	0.0112	0.7130(8)	0.0080(7)	0.0005(0)	0.0192(9)	0.0000	0.0048(0)	0.0000
MO	Ma <sup>2+</sup>	0.0000	0.3000	0.5000	0.0112	0.2040(7)	0.0060(7)	0.0003(0)	0.0192(9)	0.0000	0.0048(0)	0.0000
IVIZ	Eo <sup>2+</sup>	0.0000	0.84109(13)	0.5000	0.0120	0.0439(8)	0.0009(4)	0.0140(5)	0.0169(5)	0.0000	0.0020(4)	0.0000
т	Si <sup>4+</sup>	0.57393(14)	0.16714(10)	0.22369(9)	0.0110	0.1900(10)	0.0086(3)	0.0089(3)	0.0154(4)	-0.0002(4)	0.0021(3)	-0.0003(3)
	Si	0.57393(14)	0.16714(10)	0.22369(9)	0.0110	0.8001(10)	0.0086(3)	0.0089(3)	0.0154(4)	-0.0002(4)	0.0021(3)	-0.0003(3)
01	0.0 <sup>2-</sup>	0.8267(4)	0.2274(3)	0.1661(3)	0.0204	1	0.0170(10)	0.0248(13)	0.0203(12)	-0.0026(11)	0.0052(9)	-0.0067(10)
02	0,02-	0.5093(7)	0.0000	0.1675(4)	0.0204	1	0.0283(19)	0.0107(14)	0.0206(18)	0.0000	-0.0001(15)	0.0000
O3	0,0 <sup>2-</sup>	0.6305(4)	0.1687(3)	0.3914(2)	0.0120	1	0.0107(8)	0.0089(8)	0.0165(10)	-0.0002(10)	0.0028(7)	0.0001(9)
04	0,0 <sup>2-</sup>	0.1334(6)	0.0000	0.4006(4)	0.0132	1	0.0111(14)	0.0128(14)	0.0160(17)	0.0000	0.0032(13)	0.0000
	H	0.098(15)	0.0000	0.316(9)	0.0132	0.7209(10)	0.0111(14)	0.0128(14)	0.0160(17)	0.0000	0.0032(13)	0.0000
							-					

in Redhammer and Roth (2002)]. This indicates that substitution occurs at the O4 hydroxyl site. The shortening of the *c* parameter is partly due to OH  $\leftrightarrow$  F substitution (Boukili et al. 2001), but it has been shown that substitutions involving a deprotonation mechanism generally play a major role (Cesare et al. 2003; Scordari et al. 2006 and references therein). In particular, it is recognized that Ti-oxy substitutions lead to diagnostic values for some structural features, such as (1) shortening of the *c*parameter as well as of the K-O4 distance; (2) high values of bond length distortions (BLD) for M2, see Table 8; (3) high values of the shift of the M2 cation from the geometric center of the octahedron toward the O4 oxygen; and (4) low values for  $\Delta_{K-O4}$  and t<sub>int</sub> (Cruciani and Zanazzi 1994; Cesare et al. 2003; Schingaro et al. 2005a; Scordari et al. 2006). In addition, the occurrence of  $M^{3+,4+}$ -oxy type substitutions, with  $M^{3+,4+} = Al^{3+}$ , Fe<sup>3+</sup>, Ti<sup>4+</sup>, has been assessed in other Vulture micas (Scordari et al. 2006; Matarrese et al. 2006). In the case of the VUT191 phlogopites, however, not all samples follow the trends typical of oxy-substituted micas (see below).

Individual tetrahedra are regular and slightly elongated in the direction of the T-O<sub>apical</sub> bond. The tetrahedral mean bond lengths and parameters describing geometrical features of the tetrahedron (Tables 7, 8, and 10) do not vary much from sample to sample, consistent with the limited compositional variation of the T sites (see Table 9). The examined crystals exhibit an inverse dependence of the  $\alpha$ -values relative to their Fe contents, as expected

from comparison to literature data (Cruciani and Zanazzi 1994; Brigatti and Guggenheim 2002; Brigatti et al. 2005). Variations of octahedral bond distances, with <M1-O> in the range 2.074–2.082 Å, and <M2-O> in the range 2.064–2.072 Å (Table 7), and of the octahedral mean atomic numbers (m.a.n.), 14.13  $\leq$  m.a.n. (M1)  $\leq$  16.77 e<sup>-</sup>, 14.35  $\leq$  m.a.n. (M2)  $\leq$  17.04 e<sup>-</sup> (Table 10), reflect the variations of the annite component and of the Ti content (see Fig. 1 and accompanying discussion).

The analyzed micas are meso-octahedral from a geometrical viewpoint (Table 7), with the exception of VUT191 1 and VUT191 2 single crystals, which are homo-octahedral (Weiss et al. 1992). From a chemical viewpoint (i.e., considering the mean atomic numbers at the M1 and M2 sites; Table 10), the samples are homo-octahedral (Ďurovíč 1994). Indeed, the average error associated with the refined site-scattering power (here referred to as mean atomic number) is 0.5 e-, whereas the differences between M1 and M2 mean atomic numbers are within 1 e- (see Table 10). Systematic differences in structural parameters between samples VUT191\_1 and VUT191\_2 on the one hand and the remaining single crystals on the other, are apparent from the analyses reported in Tables 7 and 8. They are relevant to the <K-O>outer (Table 7), tint, shift<sub>M2</sub> (Table 8) parameters and are probably related to the different Ti and Fe<sup>3+</sup> substitutions affecting the two groups of samples, as well as to the different degrees of hydrogenation of the two groups of samples (see next section for further details).

## **Crystal chemistry**

Figure 3 illustrates the variation of the *c*-parameter of the micas as a function of the water content as determined by SIMS. It is apparent that micas with the shortest *c*-parameter are more dehydrogenated. However, micas with intermediate *c* values may have remarkably different degrees of dehydrogenation. Inspection of Figures 4 and 5, which plot the out-of-center-shift parameters and the  $\langle \text{K-O} \rangle_{\text{outer}}$  distances, respectively, vs. Ti content, show clearly that the samples can be divided into two distinct groups (see section Structural features) that are expected

to be affected by different substitution mechanisms. This was confirmed by the comparison of the relevant crystal chemical formulae (Table 9), whose calculation is detailed below.

The crystal-chemical formulae reported on Table 3 were obtained by combining electron microprobe, SIMS, and Mössbauer investigations, using the 12(O, OH, Cl, F) basis. When SIMS analyses where not available, the C-H-N value was used. However, the latter is an average value (see section Micas analyses) and proved not to be suitable for every single crystal, in particular for those characterized by low *c*-values. In the latter cases, indeed, micas should be more strongly dehydrogenated, and the average C-H-N value invariably leads to an overestimation of octahedral vacancies, providing unacceptable discrepancies (in the range  $3-5 e^{-}$ ) between EPMA- and SCXRD-derived mean atomic numbers. In those cases, the OH contents were estimated from structural parameters (*c*, <K-O><sub>outer</sub>, see Cruciani and Zanazzi 1994; Cesare et al. 2003), also taking into account the contribution of F to the shortening of the *c*-parameter as well



**FIGURE 3.** A plot of the crystallographic *c* parameter vs.  $H_2O$  content. Unless explicitly stated,  $H_2O$  (data) are from SIMS measurements.

TABLE 7. Results of structure refinement in space group C2/m: selected bond distances (Å)

Samples	VUT191_1	VUT191_2	VUT191_10	VUT191_11	VUT191_13	VUT191_19
T-O(1)	1.662(3)	1.663(3)	1.660(2)	1.660(2)	1.661(2)	1.657(2)
T-O(1')	1.664(3)	1.664(3)	1.665(2)	1.662(2)	1.662(2)	1.662(2)
T-O(2)	1.661(2)	1.663(2)	1.663(1)	1.666(2)	1.660(1)	1.661(2)
T-O(3)	1.656(3)	1.664(3)	1.675 (1)	1.679(2)	1.677(2)	1.677(3)
<t-0></t-0>	1.661	1.664	1.666	1.667	1.665	1.664
M1-O(4)(×2)	2.050(4)	2.037(5)	2.047(2)	2.042(3)	2.043(3)	2.038(4)
M1-O(3)(×4)	2.094(3)	2.092(3)	2.098 (2)	2.097(2)	2.102(2)	2.097(2)
<m1-0></m1-0>	2.079	2.074	2.081	2.079	2.082	2.077
M2-O(4)(×2)	2.034(3)	2.021(4)	2.010(2)	1.999(3)	1.986(2)	1.983(3)
M2-O(3)(×2)	2.088(3)	2.087(3)	2.086(1)	2.084(2)	2.085(2)	2.086(2)
M2-O(3')(×2)	2.094(3)	2.098(3)	2.110(2)	2.116(3)	2.122(2)	2.123(3)
<m2-0></m2-0>	2.072	2.069	2.069	2.066	2.064	2.064
<m-0></m-0>	2.074	2.071	2.073	2.070	2.070	2.068
K-O(1)(×4)	2.947(3)	2.939(3)	2.944(2)	2.942(2)	2.945(2)	2.943(3)
K-O(1')(×4)	3.386(3)	3.392(3)	3.364(2)	3.355(2)	3.343(2)	3.346(3)
K-O(2)(×2)	2.948(5)	2.939(5)	2.948(2)	2.941(3)	2.949(3)	2.943(4)
K-O(2')(×2)	3.396(4)	3.398(5)	3.374(2)	3.368(3)	3.361(3)	3.362(4)
<k-o><sub>inner</sub></k-o>	2.947	2.939	2.945	2.942	2.946	2.943
<k-o><sub>outer</sub></k-o>	3.389	3.394	3.367	3.359	3.349	3.351
<k-o></k-o>	3.169	3.167	3.156	3.151	3.148	3.147

<b>TABLE 8.</b> Selected distortional parameters derived from the structure refinements in space droup C2/	d from the structure refinements in space group $C_2/m$
--	---

	VUT 191_1	VUT 191_2	VUT 191_10	VUT 191_11	VUT 191_13	VUT 191_19
t <sub>tet</sub> (Å)	2.231	2.239	2.247	2.254	2.250	2.249
BLDT	0.142	0.030	0.252	0.356	0.342	0.387
Volume <sub>⊤</sub> (ų)	2.349	2.361	2.370	2.376	2.368	2.365
TQE	1.000	1.000	1.000	1.000	1.000	1.000
TAV	0.972	0.880	0.789	0.963	0.908	0.760
τ (°)	110.257	110.208	110.159	110.253	110.218	110.133
α (°)	9.63	10.02	9.24	9.09	8.74	8.90
Δz (Å)	0.008	0.004	0.012	0.011	0.019	0.014
D.M. (Å)	0.555	0.582	0.571	0.576	0.569	0.576
ψ <sub>M1</sub> (°)	58.91	59.09	59.28	59.38	59.46	59.47
ψ <sub>M2</sub> (°)	58.79	59.02	59.08	59.18	59.17	59.24
BLD <sub>M1</sub>	0.942	1.164	1.087	1.179	1.264	1.268
ELD <sub>M1</sub>	5.024	5.235	5.452	5.567	5.663	5.668
BLD <sub>M2</sub>	1.206	1.549	1.885	2.156	2.527	2.614
ELD <sub>M2</sub>	4.879	5.137	5.207	5.323	5.301	5.390
Shift <sub>M2</sub> (Å)	0.012	0.025	0.044	0.055	0.068	0.072
Volume <sub>M1</sub> (ų)	11.800	11.671	11.783	11.732	11.790	11.703
OQE <sub>M1</sub>	1.011	1.012	1.013	1.014	1.014	1.014
OAV <sub>M1</sub>	36.557	39.593	42.743	44.565	46.108	46.117
Volume <sub>M2</sub> (ų)	11.680	11.597	11.592	11.542	11.506	11.486
OQE <sub>M2</sub>	1.011	1.012	1.013	1.013	1.014	1.014
OAV <sub>M2</sub>	34.814	38.826	40.299	42.600	43.216	44.844
$e_u(M1)/e_s(M1)$	1.106	1.111	1.115	1.118	1.120	1.120
$e_u(M2)/e_s(M2)$	1.103	1.108	1.110	1.112	1.112	1.114
t <sub>oct</sub> (Å)	2.147	2.130	2.126	2.117	2.116	2.111
t <sub>int</sub> (Å)	3.424	3.415	3.360	3.339	3.329	3.332
∆ <sub>K-04</sub> (Å)	0.442	0.455	0.422	0.417	0.403	0.408
t <sub>K-O4</sub> (Å)	3.967	3.978	3.948	3.948	3.945	3.944

Notes:  $t_{tet}$  = tetrahedral sheet thickness calculated from z coordinates of basal and apical O atoms; TQE = tetrahedral quadratic elongation (Robinson et al. 1971); TAV = tetrahedral angle variance (Robinson et al. 1971);  $\tau$  = tetrahedral flattening angle;  $\alpha$  = tetrahedral rotation angle (Hazen and Burnham. 1973);  $\Delta z$  = departure from co-planarity of the basal O atoms (Güven 1971); D.M. = dimensional misfit between tetrahedral and octahedral sheets (Toraya. 1981);  $\psi$  = octahedral flattening angles (Donnay et al. 1964); BLD = bond-length distortions (Renner and Lehmann 1986); ELD = edge-length distortion (Renner and Lehman. 1986); Shift<sub>M12</sub> = off-center shift of the M2 cation defined as the distance between the refined position of cation and the geometrical center of M2 site (coordinates: x/a = 0.0, y/b= 0.8333, z/c = 0.5); OQE = octahedral quadratic elongation (Robinson et al. 1971); OAV = octahedral angle variance (Robinson et al. 1971); et al. 1971); OAV = octahedral angle variance (Robinson et al. 1971); et al. 1971); oAV = octahedral angle variance (Robinson et al. 1971); et al. 1971); oAV = octahedral angle variance (Robinson et al. 1971); et al. 1971); et al. 1971); OAV = octahedral angle variance (Robinson et al. 1971); et al. 1971); oAV = octahedral angle variance (Robinson et al. 1971); et al. 19

#### TABLE 9. Final crystal chemical formulae for VUT191 phlogopite

VUT191_1	$(K_{0.78}Na_{0.09}Ba_{0.09}\_b_{1.0})(AI_{0.14}Mg_{2.19}Fe_{0.30}^{+1}Fe_{0.30}^{+1}Fi_{0.02}C_{10.02}=0_{0.12})(Si_{2.68}AI_{1.32})O_{10.06}F_{0.10}OH_{1.84}$
VUT191_2	$(K_{0.82}Na_{0.09}Ba_{0.03} - 1,_{0.6})(AI_{0.12}Mg_{2.11}Fe_{0.11}^{2+1}Fe_{0.13}^{3+1}Ti_{0.15}Cr_{0.01} - 1,_{0.19})(Si_{2.72}AI_{1.28})O_{10}F_{0.15}OH_{1.86}$
VUT191_10	(K <sub>0.81</sub> Na <sub>0.09</sub> Ba <sub>0.07</sub> D <sub>0.03</sub> )(Al <sub>0.08</sub> Mg <sub>1.81</sub> Mn <sub>0.01</sub> Fe <sup>3+</sup> <sub>0.19</sub> Fe <sup>3+</sup> <sub>0.56</sub> Ti <sub>0.27</sub> D <sub>0.08</sub> )(Si <sub>2.62</sub> Al <sub>1.38</sub> )O <sub>10.69</sub> F <sub>0.15</sub> Cl <sub>0.01</sub> OH <sub>1.15</sub>
VUT191_11	$(K_{0.78}Na_{0.09}Ba_{0.09}-B_{0.04})(AI_{0.13}Mg_{1.73}Mn_{0.01}Fe_{0.20}^{+2.5}Fe_{0.29}^{-3.5}Ti_{0.28}-B_{0.08})(Si_{2.67}AI_{1.33})O_{10.78}F_{0.24}CI_{0.01}OH_{0.97}$
VUT191_13	$(K_{0.77}Na_{0.09}Ba_{0.08}-b_{0.09})(Al_{0.19}Mg_{1.69}Fe_{0.19}^{2+}Fe_{0.15}^{3+}Fi_{0.27}-b_{0.10})(Si_{2.71}Al_{1.29})O_{10.79}F_{0.22}Cl_{0.01}OH_{0.98}$
VUT191_19	$(K_{0.78}Na_{0.09}Ba_{0.09}\Box_{0.04})(AI_{0.17}Mg_{1.70}Mn_{0.01}Fe_{0.20}^{+3}Fe_{0.10}^{+3}Fi_{0.30}Li_{0.01}\Box_{0.08})(Si_{2.66}AI_{1.34})O_{10.79}F_{0.31}CI_{0.01}OH_{0.89}$

TABLE 10. Mean atomic numbers (e<sup>-</sup>) of cation sites and octahedral and tetrahedral mean distances (Å), as determined from structure refinement (X-ref) and chemical analyses (EPMA) (see text for details)

	VUT191_1	VUT191_2	VUT191_10	VUT191_11	VUT191_13	VUT191_19
M1 e <sup>-</sup> <sub>X-ref</sub>	14.15	14.13	16.77	16.35	16.10	15.99
M2 e <sup>-</sup> <sub>X-ref</sub>	14.35	14.65	17.00	16.96	17.04	16.95
(M1+2M2) e <sup>-</sup> X-ref	42.85	43.43	50.77	50.27	50.18	49.89
(M1+2M2) e <sup>-</sup> <sub>EMPA</sub>	41.90	41.34	48.45	48.88	48.06	49.25
K e <sup>-</sup> <sub>X-ref</sub>	19.09	19.36	21.07	21.31	21.45	21.44
K e <sup>-</sup> <sub>EMPA</sub>	17.49	18.25	20.30	20.85	20.10	20.85
T e <sup>-</sup> <sub>X-ref</sub>	13.86	13.95	13.90	13.91	13.90	13.86
T e <sup>−</sup> <sub>EMPA</sub>	13.68	13.68	13.66	13.67	13.68	13.67
<t-0></t-0>	1 661	1 664	1 666	1 667	1 665	1 664
$\langle T - O \rangle_{X-ref}$	1.667	1.667	1.670	1.668	1.667	1.669
<m-o> X-ref</m-o>	2.074	2.071	2.073	2.070	2.070	2.068
<m-o> EMPA</m-o>	2.072	2.075	2.061	2.061	2.058	2.060

as charge-balance considerations. The final crystal-chemical formulae are compiled in Table 9.

For sample VUT191-1, the average water content provided by C-H-N analysis was initially used for the formula recalculation (see Table 3). This value gave a sum of octahedral cations greater than 3 apfu. Considering the close similarity of this sample to sample VUT191 2, both with respect to chemistry (Fe/Mg, Ti

content) and structural features (Tables 7 and 8 and Figs. 4 and 5), it was concluded that the water contents of these two samples should also be similar. As a result, the final formula listed in Table 9 for VUT191\_1 was calculated using the same water content as sample VUT191\_2. For VUT191\_2 and VUT191\_19 (i.e., for samples for which SIMS data were available), the formulae in Tables 3 and 9 coincide.



**FIGURE 4.** A plot of the parameter  $shift_{M2}$  vs. Ti content. High values of the  $shift_{M2}$  parameter are associated with the occurrence of Ti-oxy substitutions.VUT191 phlogopite samples (filled diamond) plot in two separate groups (see text). SA1 and VUT 187 samples references as in Figure 1.

For samples from VUT191\_10 to VUT191\_19 in Table 9, apart from the M<sup>3+</sup>-Tschermak substitutions ( $^{VI}M^{2+} + ^{IV}Si^{4+} \leftrightarrow$  $^{VI}M^{3+} + ^{IV}Al^{3+}$ , with M<sup>3+</sup> = Al, Fe<sup>3+</sup>), the remaining M<sup>3+</sup> cations and the Ti<sup>4+</sup> cations are involved in M<sup>3+</sup>-oxy [ $^{VI}M^{2+} + (OH)^- \leftrightarrow$ M<sup>3+</sup> + O<sup>2</sup> +  $^{I}_{2}H_{2}$ ] and Ti-oxy substitutions [ $^{VI}M^{2+} + 2(OH)^- \leftrightarrow$  $^{VI}Ti^{4+} + 2O^{2-} + H_{2}$ ], respectively. Low concentrations of octahedral vacancies are present and are charge-balanced through the Ti-vacancy substitution (2  $^{VI}M^{2+} \leftrightarrow ^{VI}Ti^{4+} + ^{VI}\Box$ ).

Samples VUT191\_1 and VUT191\_2 instead have negligible or no oxy component. Sample VUT191\_1 contains a high concentration of octahedral vacancies that are balanced through the Ti-vacancy mechanism. In sample VUT191\_2, both Ti-vacancy and the dioctahedral-trioctahedral  $(3^{VI}M^{2+} \leftrightarrow 2^{VI}M^{3+} + {}^{VI}\square$ mechanism seem to be active.

The formulae in Table 9 generally demonstrate good agreement between observed and calculated mean atomic numbers, as well as between observed octahedral average bond distances and those calculated from the chemical molar fraction and atomic radii in Shannon (1976). In all the Fe- and Ti-rich samples (see Table 10, from sample VUT191 10 to VUT191 19), a tendency toward a systematic underestimation of the average octahedral bond distances is apparent. Such systematic variations of the observed bond lengths with respect to the radii of Shannon (1976) are already known (Bailey 1984; Gibbs et al. 1997; Mercier et al. 2006). In particular, Mercier et al. (2006) observed that for non end-member single-crystal biotite-1M structure, bond lengths calculated from Shannon's radii tend to overestimate octahedral mean bond lengths between the Fe<sup>2+</sup> and Mg<sup>2+</sup> end-member, whereas the opposite is true when substitution by smaller +3, +4 cations occurs. Our data are consistent with those findings. Unfortunately, the new cation and coordination-specific bond lengths that these authors provided, apart from not being complete, are more suitable for trioctahedral micas with simpler compositions close to the M<sup>2+</sup>-end members.

Another explanation for the discrepancies in observed vs. calculated average octahedral bond lengths in our samples is that the  $Fe^{2+}/Fe^{3+}$  ratio provided by Mössbauer spectroscopy could



**FIGURE 5.** A plot of the parameter  $\langle K-O \rangle_{outer}$  vs. Ti content. The decrease of  $\langle K-O \rangle_{outer}$  distance as a function of the Ti content is associated with an increase in the amount of the oxy-component. VUT191 phlogopite samples (filled diamonds) plot in two separate groups (see text). SA1 and VUT 187 samples references as in Figure 1.

be an "average" value, so that single crystals might depart from this value to different extents.

## **FURTHER REMARKS**

Petrographic and chemical evidence for VUT191 phlogopite shows a complete range of composition originating close to mantle micas and progressing up to high-*T*/low-*P* subvolcanic micas. This variation indicates that micas can be passed through different stages of magmatic evolution, starting from a foiditic (nephelinitic) and/or melilititic primary magma, to a tephrifoiditic, and then to a thepri-fonolitic term.

Chemical and structural data revealed that VUT191 phlogopite clusters into two groups, each characterized by different H contents and different Ti substitutions, in particular <sup>VI</sup>M<sup>2+</sup> + 2(OH)<sup>-</sup>  $\leftrightarrow$  <sup>VI</sup>Ti<sup>4+</sup> + 2O<sup>2-</sup> + H<sub>2</sub> (Ti-oxy) and <sup>VI</sup>M<sup>2+</sup>  $\leftrightarrow$  <sup>VI</sup>Ti<sup>4+</sup> + <sup>VI</sup> $\square$  (Ti-vacancy). This is a crucial point because the assessment of correct substitution mechanisms in biotite not only clarifies the behavior of petrologically interesting species (such as Fe, Ti) when they are incorporated into the mica structure, but also affects the choice of the annite activity model from which several petrogenetic parameters may be derived (see for instance Fabbrizio et al. 2006).

Finally, Mössbauer results of the phlogopite analyzed here, as well as those analyzed in previous works (Matarrese et al. 2005; Scordari et al. 2006; Matarrese 2007) suggest variable  $f_{O_2}$  and  $f_{H_2O}$  during the crystallization of micas of intermediate Monte Vulture pyroclastics. Thus, crystal settling of mafic minerals cannot account entirely for magma differentiation. Variable phreatomagmatic phenomena could help to explain not only such broad variations but also slightly different evolution patterns for the primary foiditic, melilititic melt.

Mica chemistry and structure can be used to predict bulk-rock chemistry, with the latter being modified by abundant analcime (plus cancrinite) formation during late-magmatic stages (Stoppa et al. 2006). The evolutionary stages experienced by the micas (see above) could parallel a possible polybaric/polythermal evolution from melilititic, nephelinitic, and phonolitic-foidite to phonolite, which is more commonly observed at Mt. Vulture. According to a previous hypothesis, all rock types of the Vulture may derive from two rock series, one melilite-bearing and one melilite-free (Melluso et al. 1996). However, melilite-bearing series share the same geological setting as melilite-free series and occur together, with melilite-bearing and melilite-free rocks being persistently similar in petrochemistry and geochemistry (e.g., isotopes). Trace-element distributions in carbonatites/melilities and foidites are also very similar. These features are difficult to reconcile with a hypothesis of two distinct parental magmas. Open-system crystallization conditions and variable  $H_2O/CO_2$ may have affected the composition of the primary magma found in melt inclusions in olivine, clinopyroxene, and apatite from the Vulture phonolitic foidite (Sololova et al. 2005).

The combination of analytical methods used in the present paper has shown that volcanic micas are able to provide information related to their geological history even in complex cases such as those considered here, where the micas are characterized by notable chemical variability. However, other than complete chemical and structural analysis, the use of single-crystal techniques, both for hydrogen determinations (such as SIMS) and for Fe speciation (such as micro-XANES or other element-sensitive techniques) could help to shed light into the uncertainties related to the results of the bulk analytical methods.

## **ACKNOWLEDGMENTS**

The authors are grateful to Marcello Serracino for assistance during electron probe microanalyses at the Istitutto di Geologia Ambientale e Geoingegneria, CNR, Rome. Antonio Giaretta is thanked for C-H-N measurements at CNR-IGC, Padova. Detailed reviews by G. Redhammer and an anonymous reviewer greatly improved the final manuscript. This work was supported by the COFIN-MIUR.

#### **REFERENCES CITED**

- Alietti, E., Brigatti, M.F., and Poppi, L. (1995) The crystal structure and chemistry of high-aluminum phlogopite. Mineralogical Magazine, 59, 149–157.
- Bailey, S.W. (1984) Micas, 13, 584 p. Reviews in Mineralogy and Geochemistry, Mineralogical Society of America, Chantilly, Virginia.
- Betteridge, P.W., Carruthers, J.R., Copper, R.I., Prouth, K., and Watkin, D.J. (2003) Crystals version 12: Software for guided crystal structure analysis. Journal of Applied Crystallography, 36, 1487–1492.
- Bohlen, S.R., Peacor, D.R., and Essene, E.J. (1980) Crystal chemistry of a metamorphic biotite and its significance in water barometry. American Mineralogist, 65, 55–62.
- Boukili, B., Robert, J.L., Beny, J.M., and Holtz, F. (2001) Structural effects of OH → F substitution in trioctahedral micas of the system: K<sub>2</sub>O-FeO-Fe<sub>2</sub>O<sub>3</sub>-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>-H<sub>2</sub>O-HF. Schweizerische Mineralogische und Petrographische Mitteilungen, 81, 55–67.
- Brigatti, M.F. and Guggenheim, S. (2002) Mica crystal chemistry and the influence of pressure, temperature, and solid solution on atomistic models. In A. Mottana, F.P. Sassi, J.B. Thompson, and S. Guggenheim, Eds., Micas: Crystal Chemistry and Metamorphic Petrology, 46, p. 1–100. Reviews in Mineralogy and Geochemistry, Mineralogical Society of America, Chantilly, Virginia.
- Brigatti, M.F., Caprilli, E., Funiciello, R., Giordano, G., Mottana, A., and Poppi, L. (2005) Crystal chemistry of ferroan phlogopites from the Albano maar lake zone (Colli Albani volcano, central Italy). European Journal of Mineralogy, 17, 611–621.
- Brocchini, D., La Volpe, L., Laurenzi, M.A., and Principe, C. (1994) Storia evolutiva del Monte Vulture. Plinius, 12, 22–25.
- Bruker (2003a) SAINT. Bruker AXS Inc., Madison, Wisconsin.

(2003b) APEX2. Bruker AXS Inc., Madison, Wisconsin.

- Cesare, B., Cruciani, G., and Russo, U. (2003) Hydrogen deficiency in Ti-rich biotite from anatectic metapelites (El Joyazo, SE Spain): Crystal-chemical aspects and implications for high-temperature petrogenesis. American Mineralogist, 88, 583–595.
- Chakhmouradian, A.R. (2006) High-field-strength elements in carbonatitic rocks: Geochemistry, crystal chemistry and significance for constraining the sources of carbonatites. Chemical Geology, 235, 138–160.
- Cruciani, G. and Zanazzi, P.F. (1994) Cation partitioning and substitution mechanism in 1M phlogopite: A crystal chemical study. American Mineralogist,

79, 289–301.

- De La Roche, H., Leterrier, J., Grande, C.P., and Marchal, M. (1980) A classification of volcanic and plutonic rocks using R1-R2 diagrams and major elements analyses: Its relationship and current nomenclature. Chemical Geology, 29, 183–210.
- Donnay, G., Morimoto, N., Takeda, H., and Donnay, J.D.H. (1964a) Trioctahedral one-layer micas. I. Crystal structure of a synthetic iron mica. Acta Crystallographica, 17, 1369–1373.
- Donnay, G., Donnay, J.D.H., and Takeda, H. (1964b) Trioctahedral one-layer micas. II Prediction of the structure from composition and cell dimensions. Acta Crystallographica, 17, 1374–1381.
- Ďurovíč, S. (1994) Classification of phyllosilicates according to the symmetry of their octahedral sheets. Ceramics—Silikàty, 38, 81–84.
- Dyar, M.D. (2002) Optical and Mössbauer spectroscopy of iron in micas. In A. Mottana, F.P. Sassi, J.B. Thompson, and S. Guggenheim, Eds., Micas: Crystal Chemistry and Metamorphic Petrology, 46, p. 313–340. Reviews in Mineralogy and Geochemistry, Mineralogical Society of America, Chantilly, Virginia.
- Fabbrizio, A., Rouse, J.P., and Carroll, M.R. (2006) New experimental data on biotite+magnetite+sanidine saturated phonolitic melts and application to the estimation of magmatic water fugacity. American Mineralogist, 91, 1863–1870.
- Feldstein, S.N., Lang, R.A., Vennemann, T., and O'Neil, J.R. (1996) Ferric-ferrous ratios, H<sub>2</sub>O contents and D/H ratios of phlogopite and biotite from lavas of different tectonic regimes. Contributions to Mineralogy and Petrology, 126, 51–66.
- Forbes, W.C. and Flower, M.F.J. (1974) Phase relations of titan-phlogopite, K<sub>2</sub>Mg<sub>4</sub>TiAl<sub>2</sub>Si<sub>6</sub>O<sub>20</sub>(OH)<sub>4</sub>: a refractory in the upper mantle? Earth and Planetary Science Letters, 22, 60–66.
- Giannandrea, P., La Volpe, L., Principe, C., and Schiattarella, M. (2006) Unità stratigrafiche a limiti inconformi e storia evolutiva del vulcano medio-pleistocenico del Monte Vulture (Appennino Meridionale, Italia). Bollettino della Societa' Geologica Italiana, 125, 67–92.
- Gibbs, G.V., Tamada, O., and Boisen, Jr., M.B. (1997) Atomic and ionic radii: a comparison with radii derived from electron density distributions. Physics and Chemistry of Minerals, 24, 432–439.
- Güven, N. (1971) The crystal structure of 2*M*<sub>1</sub> phengite and 2*M*<sub>1</sub> muscovite. Zeischrift für Kristallographie, 134, 196–212.
- Hawthorne, F.C., Ungaretti, L., and Oberti, R. (1995) Site populations in minerals: terminology and presentation of results. Canadian Mineralogist, 33, 907–911.
- Hazen, R.M. and Burnham, C.W. (1973) The crystal structure of one layer phlogopite and annite. American Mineralogist, 58, 889–900.
- Henry, D.J. and Guidotti, C.V. (2002) Titanium in biotite from metapelitic rocks: Temperature effects, crystal-chemical controls, and petrologic applications. American Mineralogist, 87, 375–382.
- Henry, D.J., Guidotti, C.V., and Thompson, J. (2005) The Ti-saturation surface for low-to-medium pressure metapelitic biotites: implications for geothermometry and Ti-substitution mechanisms. American Mineralogist, 90, 316–328.
- Irvine, T.N. and Bargar, W.R.A. (1971) A guide to the chemical classification of the common volcanic rocks. Canadian Journal of Earth Science, 8, 523–548.
- Lagarec, K. and Rancourt, D.G. (1997) Extended Voigt-based analytic lineshape method for determining N-dimensional correlated hyperfine parameter distributions in Mössbauer spectroscopy. Nuclear Instruments and Methods in Physics Research, B129, 266–280.
- (1998) RECOIL. Mössbauer spectral analysis software. University of Ottawa, Canada (www.isapps.ca/recoil).
- Matarrese, S. (2007). Cristallochimica comparativa di flogopiti del Monte Vulture, 170 p. Ph.D. thesis, University of Bari, Italy.
- Matarrese, S., Mesto, E., Pedrazzi, G., Schingaro, E., and Scordari, F. (2005) Trioctahedral micas from the youngest volcanics of Mt. Vulture (Pz, Italy): a Crystal chemical study. International workshop Micas@Italy, Rimini(Italy), February 9–11, 2005, Book of Abstract, 33.
- Matarrese, S., Schingaro E., Scordari, F., Rosatelli, G., Stoppa, F., and Pedrazzi, G. (2006) Micas as indicators of geological processes: The case of Mt. Vulture (Italy). Annual meeting of the Geological Society of America, Philadelphia, 22–25 October 2006. Geological Society of America, Abstracts with Programs, Vol. 38, No. 7, p. 292.
- Melluso, L., Morra, V., and Di Girolamo, P. (1996) The Mt. Vulture volcanic complex (Italy): Evidence for distinct parental magmas and for residual melts wirh melilite. Mineralogy and Petrology, 56, 225–250.
- Mercier, P.H.J., Rancourt, D.G., Redhammer, G.J., Lalonde, A.E., Robert, J-L., Berman, R.G., and Kodama, H. (2006) Upper limit of tetrahedral rotation angle and factors affecting octahedral flattening in synthetic and natural 1*M* polytype *C2lm* space group micas. American Mineralogist, 91, 831–849.
- Mesto, E., Schingaro, E., Scordari, F., and Ottolini, L. (2006) Electron probe microanalysis, secondary ion mass spectrometry and single crystal X-ray diffraction study of phlogopites from Mt. Vulture, Potenza, Italy: Consideration of cation partitioning. American Mineralogist, 91, 182–190.
- Nespolo, M. and Ferraris, G. (2001) Effects of the stacking faults on the calculated

electron density of mica polytypes—The Ďurovíč effect. European Journal of Mineralogy, 13, 1035–1045.

- Ottolini, L. and Hawthorne, F.C. (2001) SIMS ionization of hydrogen in silicates: a case study of kornerupine. Journal of Analytical Atomic Spectrometry, 6, 1266–1270.
- Ottolini, L., Bottazzi, P., and Vannucci, R. (1993) Quantification of lithium, beryllium and boron in silicates by secondary ion mass spectrometry using conventional energy filtering. Analytical Chemistry, 65, 1960–1968.
- Ottolini, L., Bottazzi, P., Zanetti, A., and Vannucci, R. (1995) Determination of hydrogen in silicates by secondary ion mass spectrometry. Analyst, 120, 1309–1313.
- Ottolini, L., Camara, F., Hawthorne, F.C., and Stirling, J. (2002) SIMS matrix effects in the analysis of light elements in silicate minerals: Comparison with SREF and EMPA data. American Mineralogist, 87, 1477–1485.
- Pouchou, J.L. and Pichoir, F. (1985) 'PAP' Φ(ρZ) procedure for improved quantitative micro-analysis. Microbeam Analysis, 104–160.
- Rancourt, D.G. and Ping, J.Y. (1991) Voigt-based methods for arbitrary-shape static hyperfine parameter distribution in Mössbauer spectroscopy. Nuclear Instruments and Methods in Physics Research, B58, 85–97.
- Rancourt, D.G., Tume, P., and Lalonde, A.E. (1993) Kinetics of the (Fe<sup>2+</sup>OH)<sub>mica</sub> → (Fe<sup>3+</sup>O<sup>2-</sup>)<sub>mica</sub> + H oxidation reaction in bulk single crystal biotite studied by Mössbauer spectroscopy. Physics and Chemistry of Minerals, 20, 276–284.
- Rancourt, D.C., Christie, I.A.D., Royer M., Kodama, H., Robert, J.-L., Lalonde, A.E., and Murad, E. (1994a) Determination of accurate <sup>4</sup>Fe<sup>3+</sup>, <sup>6</sup>Fe<sup>3+</sup>, <sup>6</sup>Fe<sup>3+</sup>, and <sup>6</sup>Fe<sup>2+</sup> site populations in synthetic annite by Mössbauer spectroscopy. American Mineralogist, 79, 51–62.
- Rancourt, D.G., Ping, J.Y., and Berman, R.G. (1994b) Mössbauer Spectroscopy of Minerals III. Octahedral-site Fe<sup>2+</sup> quadrupole splitting distributions in the phlogopite-annite series. Physics and Chemistry of Minerals, 21, 258–267.
- Redhammer, G.J. (1998) Characterization of synthetic trioctahedral micas by Mössbauer spectroscopy. Hyperfine Interactions, 117, 85–115.
- Redhammer, G.J. and Roth, G. (2002) Single-crystal structure refinements and crystal chemistry of synthetic trioctahedral micas KM<sub>3</sub>(Al<sup>3+</sup>, Si<sup>4+</sup>)<sub>4</sub>O<sub>10</sub>(OH)<sub>2</sub>, where M = Ni<sup>2+</sup>, Mg<sup>2+</sup>, Co<sup>2+</sup>, Fe<sup>2+</sup>, or Al<sup>3+</sup>. American Mineralogist, 87, 1464–1476.
- Redhammer, G.J., Amthauer, G., Lottermoser, W., Bernroider, M., Tippelt, G., and Roth, G. (2005) X-ray powder diffraction and <sup>57</sup>Fe-Mössbauer spectroscopy of synthetic trioctahedral micas {K}(Me<sub>3</sub>) <TSi<sub>3</sub>>O<sub>10</sub>(OH)<sub>2</sub>, Me = Ni<sup>2+</sup>, Mg<sup>2+</sup>, Co<sup>2+</sup>, Fe<sup>2+</sup>; T = Al<sup>3+</sup>, Fe<sup>3+</sup>. Mineralogy and Petrology, 85, 89–115.
- Renner, B. and Lehmann, G. (1986) Correlation of angular and bond length distortions in TO<sub>4</sub> units in crystals. Zeitschrift f
  ür Kristallographie, 175, 43–59.
- Righter, K., Dyar, M.D., Delaney, J.S., Vennemann, T.W., Hervig, R.L., and King, P.L. (2002) Correlations of octahedral cations with OH, O<sup>2</sup>, Cl, and F in biotite from volcanic rocks and xenoliths. American Mineralogist, 87, 142–153.
- Robert, J.L. (1976) Titanium solubility in synthetic phlogopite solid solution. Chemical Geology, 17, 213–227.
- Robinson, K., Gibbs, G.V., and Ribbe, P.H. (1971) Quadratic elongation, a quantitative measure of distortion in coordination polyhedra. Science, 172, 567–570.

- Salvador, A. (1987) Unconformity-bounded stratigraphic units. Geological Society of American Bulletin, 98, 232–237.
- Schingaro, E., Scordari, F., and Ventruti, G. (2001) Trioctahedral micas-1*M* from Mt. Vulture (Italy): Structural disorder and crystal chemistry. European Journal of Mineralogy, 13, 1057–1069.
- Schingaro, E., Scordari, F., Mesto, E., Brigatti, M.F., and Pedrazzi, G. (2005a) Cation site partitioning in Ti-rich micas from Black Hill (Australia): A multitechnical approach. Clays and Clay Minerals, 53, 179–189.
- Schingaro, E., Scordari, F., and Ottolini, L. (2005b) Ti and Fe rich phlogopites from Mt. Vulture (Potenza, Italy): A combined EPMA, SIMS and SCXRD study. GEOITALIA 2005, Spoleto (Italy), September 21–23, 2005, EPITOME, 61.
- Schingaro, E., Scordari, F., Matarrese, S., Stoppa, F., Rosatelli, G., and Pedrazzi, G. (2006) Trioctahedral micas from Mt. Vulture (Potenza, Italy): chemical and structural trends. XXXV AIC meeting, Ferrara, september 18–21, 2006. Collected Abstracts, MS1-O3.
- Scordari, F., Ventruti, G., Sabato, A., Bellatreccia, F., Della Ventura, G., and Pedrazzi, G. (2006) Ti-rich phlogopite from Monte Vulture (Potenza, Italy) investigated by a multianalytical approach: Substitutional mechanisms and orientation of the OH dipoles. European Journal of Mineralogy, 18, 379–391.
- Shabani, A.A.T. (1999) Mineral Chemistry and Mössbauer Spectroscopy of Micas from Granitic Rocks of the Canadian Appalachians. Ph.D. thesis, University of Ottawa, Canada.
- Shannon, R.D. (1976) Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. Acta Crystallographica, A32, 751–767.
- Sheldrick, G.M. (2003) SADABS, Program for empirical absorption correction of area detector data. University of Göttingen, Germany.
- Sololova, I.P., Girnis, A.V., Kogarko, L.N., Kononkova, N.N., Stoppa, F., and Rosatelli, G. (2005) Compositions of magmas and carbonate-silicate liquid immiscibility in the Vulture alkaline igneous complex, Italy. Lithos, Special Issue Eurocarb, 85, 113–128.
- Stoppa, F., Rosatelli, G., and Principe, C. (2006) Classificazione modale delle vulcaniti del Monte Vulture. In C. Principe, Ed., La geologia del Monte Vulture, Regione Basilicata, p. 87–103. CNR, Potenza, Italy.
- Toraya, H. (1981) Distortions of octahedra and octahedral sheets in 1*M* micas and the relation to their stability. Zeitschrift für Kristallographie, 157, 173–190.
- Virgo, D. and Popp, R.K. (2000) Hydrogen deficiency in mantle-derived phlogopites. American Mineralogist, 85, 753–759.
- Watkin, D.J. (1994) The control of difficult refinements. Acta Crystallographica, A50, 411–437.
- Weiss, Z., Rieder, M., and Chmíelovà, M. (1992) Deformation of coordination polyhedra and their sheets in phyllosilicates. European Journal of Mineralogy, 4, 665–682.

MANUSCRIPT RECEIVED FEBRUARY 1, 2007

MANUSCRIPT ACCEPTED JULY 20, 2007

MANUSCRIPT HANDLED BY FRANCESCO SASSI