

Gamagarite

Ba₂(Fe³⁺, Mn³⁺)(VO₄)₂(OH)

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic. *Point Group:* 2/m. Needlelike crystals, elongated along [010], to more than 1 cm, and as prisms, flattened on [001]; as anhedral grains.

Physical Properties: *Cleavage:* {001} and {100}, distinct; {101} difficult, indices probable. Hardness = 4.5–5 D(meas.) = 4.62–4.68 D(calc.) = [4.72] Moderately magnetic.

Optical Properties: Transparent in thin fragments. *Color:* Dark brown to almost black.

Streak: Reddish brown. *Luster:* Adamantine.

Optical Class: Biaxial (+). *Pleochroism:* X = red-brown; Y = very deep red-brown; Z =

light salmon-buff. *Orientation:* Y = b; X ∧ c = 39°–43°. *Dispersion:* r < v, fairly strong.

Absorption: Y > X > Z. α = 2.016(10) β = 2.040(10) γ = 2.130(10) 2V(meas.) = 46°–62°

Cell Data: *Space Group:* P2₁/m. a = 9.121–9.15 b = 6.142–6.17 c = 7.838–7.88
β = 112.7°–112.88° Z = 2

X-ray Powder Pattern: Gloucester Farm, South Africa.

3.309 (100), 3.051 (80), 2.805 (80), 2.349 (50), 4.969 (10), 4.627 (10), 3.711 (10)

Chemistry:

	(1)	(2)	(3)
As ₂ O ₅			3.47
V ₂ O ₅	31.9	31.9	29.07
Al ₂ O ₃	0.0	0.8	
Fe ₂ O ₃	8.4	8.1	7.80
Mn ₂ O ₃		5.7	6.11
MnO	5.5		
SrO			3.60
BaO	52.4	53.5	48.37
H ₂ O ⁺	1.06	[1.91]	[1.58]
Total	99.3	[101.9]	[100.00]

(1) Gloucester Farm, South Africa; recalculated from original analysis. (2) Do.; by electron microprobe, total Fe as Fe₂O₃, total Mn as Mn₂O₃, H₂O calculated for charge balance; corresponds to Ba_{1.99}(Fe_{0.58}³⁺Mn_{0.41}³⁺Al_{0.09})_{Σ=1.08}(VO₄)₂(OH)_{1.22}. (3) Molinello mine, Italy; by electron microprobe, total Fe as Fe₂O₃, total Mn as Mn₂O₃, H₂O calculated from crystal-structure analysis; corresponding to (Ba_{1.80}Sr_{0.20})_{Σ=2.00}(Fe_{0.56}³⁺Mn_{0.44}³⁺)_{Σ=1.00}[(VO₄)_{1.83}(AsO₄)_{0.17}]_{Σ=2.00}(OH)_{1.00}.

Mineral Group: Brackebuschite group.

Occurrence: Very rare, in banded manganese ore replacing shale (Gloucester Farm, South Africa); in veinlets cutting manganese ore (Molinello mine, Italy).

Association: Bixbyite, hematite, ephesite, diaspore, amesite (Gloucester Farm, South Africa); braunite, saneroite, bementite, caryopilite, calcite, quartz (Molinello mine, Italy).

Distribution: Along Gamagara Ridge, on Gloucester Farm, about 27 km north of Postmasburg, Cape Province, South Africa. In the Molinello manganese mine, near Chiavari, Val Graveglia, Liguria, Italy.

Name: For the original occurrence on Gamagara Ridge, South Africa.

Type Material: The Natural History Museum, London, England, 1958,553; National Museum of Natural History, Washington, D.C., USA, 105142.

References: (1) de Villiers, J.E. (1943) Gamagarite, a new vanadium mineral from the Postmasburg manganese deposits. *Amer. Mineral.*, 28, 329–335. (2) Harlow, G.E., P.J. Dunn, and G.R. Rossman (1984) Gamagarite: a re-examination and comparison with brackebuschite-like minerals. *Amer. Mineral.*, 69, 803–806. (3) Basso, R., A. Palenzona, and L. Zefiro (1987) Gamagarite: new occurrence and crystal structure refinement. *Neues Jahrb. Mineral., Monatsh.*, 295–305.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.