

# Manganese valency and the colour of the $\text{Mn}_2\text{AsO}_4(\text{OH})$ polymorphs eveite and sarkinite

ULF HÄLENIUS

Department of Mineralogy, Swedish Museum of Natural History, P.O. Box 50007, SE-10405 Stockholm, Sweden

AND

ERIKA WESTLUND

Department of Geology and Geochemistry, Stockholm University, SE-10691 Stockholm, Sweden

## ABSTRACT

Polarized optical absorption spectra of single crystals of the two dimorphic  $\text{Mn}_2\text{AsO}_4(\text{OH})$ -minerals eveite and sarkinite show that minor concentrations of  $\text{Mn}^{3+}$  at the octahedral site determine the colour and cause the distinct pleochroism in green and yellow of eveite. In the sarkinite spectra only absorption bands due to spin-forbidden  $d-d$  transitions in  $\text{Mn}^{2+}$  at six- and five-coordinated sites are observed, which produce a weak flesh-red mineral colour and a very faint pleochroism.

**KEY WORDS:** eveite, sarkinite, optical spectroscopy, Långban, Sweden.

## Introduction

EVEITE and sarkinite, two naturally occurring polymorphs of  $\text{Mn}_2\text{AsO}_4(\text{OH})$ , display contrasting properties which reflect their different structural character. Eveite, which is a member of the orthorhombic adamite group and thus is structurally related to the rock-forming silicate andalusite (Moore and Smythe, 1968) has, e.g. considerably lower density than the closer packed monoclinic sarkinite (Fig. 1), which is structurally related to wagnerite (Dal Negro *et al.*, 1974). However, the causes for the contrasting colours observed for the two minerals are less obvious. In spite of large similarities with respect to  $\text{Mn}^{2+}$  coordination, sarkinite is faintly pleochroic in flesh-red hues while eveite displays a distinct pleochroism with  $X(a) = Z(b) = \text{green}$  and  $Y(c) = \text{yellow}$ . In eveite there is one Mn-centred octahedral site and one independent five-coordinated Mn-site, while in sarkinite there are four independent octahedral Mn-sites as well as four unique five-coordinated Mn-centered sites. The individual octahedral sites in the two minerals are comparable, each displaying two OH-ligands in cis-configuration, and mean  $M-L$  distances in the

range 2.19–2.23 Å. The trigonal bipyramidal sites in the two minerals also show comparable features including the presence of one OH-ligand and mean  $M-L$  distances ranging from 2.11 to 2.15 Å. The observed difference in colour was commented on by Moore (1968), who ascribed it to a possibly higher crystal field strength at the Mn-sites in eveite as compared to the Mn-sites in sarkinite. In the present study polarized crystal field spectra of chemically well-characterized single crystals of eveite and sarkinite have been recorded with the aim to shed some light on the effects responsible for the observed colour difference.

## Materials and methods

Natural single crystals of eveite (NRM # 390271) and sarkinite (NRM # 741004) from Långban, Sweden were ground and polished to produce self-supporting double-sided polished thin sections. The crystals used were selected on the basis of their crystal morphology, transparency and lack of inclusions. Two eveite crystal sections (cut perpendicular to (010) and (001)) were prepared, while three sections (cut




FIG. 1. The crystal structure of eveite and sarkinite projected on (001). Strongly shaded, intermediately shaded and open polyhedra represent five-coordinated Mn-centred sites, Mn-octahedra and As-tetrahedra, respectively.

perpendicular to (100), (010) and (001)) were prepared from three sarkinite crystals. The orientation of the crystal sections were confirmed by means of crystal morphology and conoscopic microscopy. As determined by means of digital micrometer measurements the thickness of the eveite absorbers were 265 and 266  $\mu\text{m}$  and the sarkinite absorbers were 530, 560 and 753  $\mu\text{m}$ . In addition to this, approximately 2 mg of each mineral was separated under a binocular microscope and ground to a powder suitable for FTIR-measurements.

The self-supporting crystal sections were measured in polarized light by optical microscope-spectrometry in the range 333–2,000 nm ( $30\,000\text{--}5\,000\text{ cm}^{-1}$ ) using a Zeiss MPM 800-instrument. The general instrumental set up (light sources, lenses, monochromators, detectors and polarizers) in the different wavelength regions has been described by Hålenius and Lindqvist (1996). In the present study the spectral slit widths adopted were 1 nm in the range 333–800 nm and 5 nm in the range 800–2,000 nm. The field aperture and the object aperture were in all measurements 100 and 60  $\mu\text{m}$ , respectively. The measurements were performed at ambient temperature and air served as a reference medium. Raw spectra were analysed by means of peak resolution using the Peak Fit 4 computer

program under the assumption of Gaussian band shapes.

Subsequent to the optical absorption measurements the five absorbers were analysed at the spots used for these recordings by means of a Cameca SX50-microprobe running at an acceleration voltage of 20 kV and a sample current of 12.0 nA. At each spot 7–8 analyses were performed. Standard samples included synthetic  $\text{CaSiO}_3$  (Ca), Cu-metal (Cu),  $\text{Fe}_2\text{O}_3$  (Fe), GaAs (As), MgO (Mg),  $\text{MnTiO}_3$  (Mn, Ti), ZnS (Zn) and natural vanadinite (Cl). Neither Ti nor Cl were found in detectable concentrations. Corrections of raw data were performed by means of the ZAF-related PAP-program (Pouchou and Pichoir, 1984).

Fourier Transform Infrared (FTIR) spectra of the separated mineral powders were obtained with a Biorad-instrument using Nujol emulsions placed on a transparent  $\text{CaF}_2$  disc. The spectra were recorded during 64 cycles at a spectral resolution of  $2\text{ cm}^{-1}$ .

## Results

The chemical analyses of the two eveite and three sarkinite crystals are summarized in Table 1. The obtained analyses show that the crystals used in the present study are compositionally very close

TABLE 1. Electron microprobe analyses of sarkinite and eveite

	Sarkinite A N = 8	Sarkinite B N = 8	Sarkinite C N = 8	Eveite A N = 7	Eveite B N = 8
As <sub>2</sub> O <sub>5</sub>	44.09	44.41	44.23	43.34	42.44
MgO	0.19	0.15	0.19	1.78	0.50
CaO	0.29	0.25	0.29	0.32	0.68
MnO	51.77	51.52	51.49	49.26	51.32
FeO	0.02	0.02	0.02	0.02	0.00
CuO	0.01	0.00	0.01	0.01	0.00
ZnO	0.15	0.08	0.10	0.00	0.02
H <sub>2</sub> O <sub>calc</sub>	3.40	3.40	3.40	3.38	3.34
Total	99.93	99.84	99.72	98.10	98.29
Cations on the basis of 9 negative charges					
As	1.015	1.022	1.019	1.006	0.994
Mg	0.013	0.010	0.013	0.118	0.033
Ca	0.014	0.012	0.014	0.015	0.032
Mn	1.931	1.920	1.922	1.852	1.948
Fe	0.001	0.001	0.001	0.001	0.000
Cu	0.000	0.000	0.000	0.000	0.000
Zn	0.005	0.003	0.003	0.000	0.001

to the nominal mineral chemistry. The analyses also confirm the chemical homogeneity of the individual crystals. Substitutions at the Mn-sites are limited and mainly restricted to Ca and Mg proxying for Mn. The total extent of the substitutions is less than 7% in the present samples. The eveite analyses also show a low oxide sum, apart from the presence of detectable concentrations of Mg, which were not observed in previous analyses of the mineral (Moore, 1968). The only microprobe analyses of eveite so far published (Moore, 1968) similarly display low totals. A series of WDS-scans on the present eveite crystals down to  $Z = 10$  did not reveal detectable concentrations of any elements other than those reported in Table 1.

The recorded polarized optical absorption spectra of eveite (E//X(a), E//Y(c) and E//Z(b)) and sarkinite (E//X, E//Y(b), E//Z, E//a and E//c) are illustrated in Figs 2 and 3. Due to the non-orthogonality of the sarkinite cell five polarized spectra were recorded in order to obtain a complete absorption character (see, e.g. Dowty, 1978). The sarkinite spectra which are weakly polarized show a number of sharp to relatively sharp absorption bands in the UV-VIS spectral range. A very similar set of absorption bands are also observed in the eveite spectra, although the

absorption bands in the UV-region are less well-resolved due to stronger L-M absorption, which pushes the UV-absorption edge further into the visible spectral region. In addition to these absorption bands, which are common to the two




Fig. 2. Polarized absorption spectra of eveite. Arrows indicate absorption bands assigned to trivalent manganese. The X- and Z-spectra are for the sake of clarity offset with respect to the absorption baseline.




FIG. 3. Polarized absorption spectra of sarkinite. All spectra except E//c are offset with respect to the absorption baseline.

minerals, the eveite spectra reveal two absorption bands at 22 160 and 16 485  $\text{cm}^{-1}$ . These two broader bands are distinctly anisotropic and they determine the colour and pleochroism of eveite. In addition to the bands in the UV and visible spectral ranges there are two strongly polarized narrow absorption features in the NIR-range of the eveite spectra at 6965  $\text{cm}^{-1}$  and at 5200  $\text{cm}^{-1}$ . In the sarkinite spectra just one polarized NIR-absorption feature at approximately 6920  $\text{cm}^{-1}$  is recorded. This feature is, from detailed recordings at a spectral resolution of 2 nm, demonstrated to be composed of at least four absorption bands at 6855, 6875, 6915 and 6940  $\text{cm}^{-1}$ . All the NIR-bands in the spectra of the two minerals are distinctly anisotropic.

The FTIR-spectrum of eveite (Fig. 4) is characterized by one sharp absorption band at 3560  $\text{cm}^{-1}$  and two broad and less intense bands at 3340 and 3450  $\text{cm}^{-1}$ . In the FTIR-spectrum of sarkinite (Fig. 4) a set of overlapping sharp absorption bands centred at 3505, 3515, 3525 and 3535  $\text{cm}^{-1}$  are evident.

## Discussion

In view of the nominally identical set-up of transition metal cations ( $\text{Mn}^{2+}$ ) and the similarities in symmetry and geometry of the Mn-centred sites in the two minerals one would expect




FIG. 4. Unpolarized powder FTIR absorption spectra of eveite and sarkinite.

absorption spectra of the two minerals to be very similar, with possibly only minor shifts in peak positions and band intensities. With the exception of the absorption bands at 22160 and 16485  $\text{cm}^{-1}$ , and a stronger UV-absorption in the eveite spectra, the strong similarities in the spectral character of the two minerals are indeed evident. The absorption bands in common to the two minerals are all relatively weak ( $\epsilon < 0.8$  l/mole·cm) and have energies and band widths comparable to those recorded in a large number of  $\text{Mn}^{2+}$ -bearing minerals (Rossman, 1988a; Burns, 1993). These bands are tentatively assigned to different spin-forbidden  $d-d$  transitions in divalent manganese at six- and five-coordinated sites (Fig. 5 and Table 2). In this assignment scheme, the Racah B- and C-parameters for six-coordinated  $\text{Mn}^{2+}$  in eveite are calculated at 639 and 3583  $\text{cm}^{-1}$ , respectively, and in sarkinite at 646 and 3578  $\text{cm}^{-1}$ , respectively. For the five-coordinated divalent manganese, B- and C-values are determined to be 623 and 3550  $\text{cm}^{-1}$  in eveite and 619 and 3563  $\text{cm}^{-1}$  in sarkinite. The crystal field splitting parameter Dq for divalent manganese at the five-coordinated sites in eveite and sarkinite, 556 and 569  $\text{cm}^{-1}$ , are found to be considerably smaller than for the six-coordinated manganese, 907 and 902  $\text{cm}^{-1}$ , in the two minerals. For this assignment the resulting Dq-ratio for  $\text{Mn}^{2+}$  at the two different types of coordination sites is  $\sim 0.62$ , which is very close to the theoretical ratio of 0.52 (Burns, 1993). This theoretical value is based on coordination sites having equal  $M-L$  distances. As the  $M-L$  distances of the five-coordinated sites in eveite




FIG. 5. Energy level diagram of Mn<sup>2+</sup> (d<sup>5</sup>) with B = 664 and C = 3560 cm<sup>-1</sup> (Modified from Moore and White, 1972). Vertical lines and open and filled circles indicate the approximate positions of bands caused by six- and five-coordinated Mn<sup>2+</sup>, respectively, in eveite and sarkinite.

and sarkinite are slightly shorter than those of the six-coordinated sites, the D<sub>q</sub>/D<sub>q<sub>VI</sub></sub>-ratios in the present cases are expected to be 10–30 % higher. The lower B-values found for Mn<sup>2+</sup> at the five-coordinated sites as compared to the octahedrally

bonded divalent manganese, indicate a slightly higher degree of covalent bonding, which is consistent with the shorter M–L distances for the five-coordinated sites.

Alternative Mn<sup>2+</sup>-assignment models, including the eveite spectral bands at 16485 and 22160 cm<sup>-1</sup> were explored in addition to the one presented in Table 2 and Fig. 5. These additional schemes yielded unreasonably high D<sub>q</sub>-values for five-coordinated Mn<sup>2+</sup> in eveite, which equalled or even exceeded those obtained for divalent manganese at the six-coordinated sites of the two arsenates. Considering the strongly similar Mn<sup>2+</sup>-coordination in terms of site geometries, ligand types and next-nearest neighbours in the two minerals, as well as the theoretically predicted D<sub>q<sub>V</sub></sub>/D<sub>q<sub>VI</sub></sub>-relationship, the parameters obtained from the alternative assignment models seem unrealistic and so these models are judged to be less likely than the one presented in Table 2 and Fig. 5.

The two additional absorption bands observed at 22160 and 16485 cm<sup>-1</sup> in the eveite spectra, and which cause the colour and pleochroism of the mineral, are thus obviously not related to divalent manganese at the five- or six-coordinated site. Furthermore, the EMP-analyses of the present eveite crystals show that no other 3d-elements other than Mn are present in concentrations likely to produce detectable absorption bands. Even an assumed and highly overestimated ε-value of 200 l/mole·cm for a potential Fe<sup>2+</sup>-band in eveite, in combination with the observed maximum FeO-concentration of 0.02 wt.% would result in bands with extinction coefficients of < 2.0 cm<sup>-1</sup>.

TABLE 2. Band positions and assignments

Band energy (cm <sup>-1</sup> )		Band assignment	
Sarkinite	Eveite	Cation	Transition
28870	28770	Mn <sup>2+</sup> [VI]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> E(D)
28330	28340	Mn <sup>2+</sup> [V]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> E(D)
27610	27630	Mn <sup>2+</sup> [V]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>2</sub> (D)
27070	26950	Mn <sup>2+</sup> [VI]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>2</sub> (D)
24350	24300	Mn <sup>2+</sup> [VI]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> A <sub>1</sub> <sup>4</sup> E(G)
24000	23980	Mn <sup>2+</sup> [V]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> A <sub>1</sub> <sup>4</sup> E(G)
23150	23570	Mn <sup>2+</sup> [V]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>2</sub> (G)
–	22160	Mn <sup>3+</sup> [VI]	<sup>5</sup> B <sub>1g</sub> (D) → <sup>5</sup> B <sub>2g</sub> (D)
21800	–	Mn <sup>2+</sup> [VI]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>2</sub> (G)
21000	21470	Mn <sup>2+</sup> [V]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>1</sub> (G)
18190	18130	Mn <sup>2+</sup> [VI]	<sup>6</sup> A <sub>1g</sub> (S) → <sup>4</sup> T <sub>1</sub> (G)
–	16485	Mn <sup>3+</sup> [VI]	<sup>5</sup> B <sub>1g</sub> (D) → <sup>5</sup> A <sub>1g</sub> (D)

Obviously the cause for the two strong (approaching  $10\text{ cm}^{-1}$ ), relatively broad ( $\omega_{1/2}$  approximately  $2000\text{ cm}^{-1}$ ) and anisotropic bands in the visible region of the eveite spectra must be sought elsewhere. In the absence of significant concentrations of additional 3d-elements only one likely alternative remains, i.e. a fraction of the manganese is present in the trivalent state. The broadness of the two bands, their strong polarization as well as their energies are also consistent with an assignment to spin-allowed  $d-d$  transitions in trivalent manganese. The band energies are comparable to those observed for spin-allowed  $d-d$  bands in  $\text{Mn}^{3+}$  at distorted octahedral sites in a number of minerals (Burns, 1993), and in particular to the  $\text{Mn}^{3+}$ -bands ( $22\,100$  and  $16\,000\text{ cm}^{-1}$ ) recorded in the spectra of the isostructural mineral kanonaite (Smith *et al.*, 1982). In analogy with the observed ordering of the Jahn-Teller cation  $\text{Cu}^{2+}$  ( $3d^9$ ) in isostructural Cu-substituted adamite (Chisholm, 1985) it is proposed that  $\text{Mn}^{3+}$  ( $3d^4$ ), which is also susceptible to large Jahn-Teller distortions, has a strong preference for the octahedral site in eveite. Taking an  $\varepsilon$ -value of  $200\text{ l/mole}\cdot\text{cm}$ , which corresponds to the upper limit of  $\varepsilon$ -values observed for spin-allowed  $\text{Mn}^{3+}$   $d-d$  bands at the octahedral site in the isostructural mineral kanonaite (Smith *et al.*, 1982), the concentration of  $\text{Mn}_2\text{O}_3$  in the present eveite samples is calculated to be  $\sim 0.10\text{ wt.}\%$ . Based on the present assignment of trivalent manganese exclusively to the octahedral site in eveite, this implies that only  $0.4\%$  of the octahedra are  $\text{Mn}^{3+}$ -centred and the remaining  $99.6\%$  are mainly occupied by  $\text{Mn}^{2+}$ . A number of coupled cation substitutions for incorporating trivalent cations under charge balance conditions in eveite are possible. These alternatives involve the presence of reduced As-cation species, dehydroxylation and cation vacancies. The IR-spectrum of eveite does not show any evidence for  $\text{As}^{3+}$ -clusters, which possess IR-active modes that should give rise to absorption bands at energies lower than bands caused by  $\text{As}^{5+}$ -clusters. On the other hand it is questionable if the low concentrations of  $\text{As}^{3+}$  required to enable the minor degree of  $\text{Mn}^{3+}$ -incorporation would be detectable in the IR-spectrum. It is concluded that it is not possible on the basis of the present data sets to unequivocally assign a mechanism for the substitution of trivalent Mn in eveite. An on-going laser Raman study with the aim to obtain information on the As valence state distribution in eveite may give some further useful results.

The sharp IR-band recorded at  $3560\text{ cm}^{-1}$  in the powder spectrum of eveite is in accordance with previous studies on the isostructural Zn-arsenate adamite assigned to a stretching mode of the structural OH-group (Braithwaite, 1983). The slightly higher energy of the OH-stretching band in eveite as compared to adamite ( $3540\text{ cm}^{-1}$ , from Braithwaite, 1983) is ascribed to the lower electronegativity of Mn relative to Zn. The absorption bands found between  $3400$  and  $3600\text{ cm}^{-1}$  in the sarkinite IR-spectrum are also assigned to OH-stretching modes. The more complex IR-absorption in sarkinite, which displays a number of overlapping OH-bands, is explained by the fact that in sarkinite there are four independent OH-ligands as compared to one in adamite. The absorption bands recorded in the NIR-spectra of both minerals between  $6800$  and  $7000\text{ cm}^{-1}$  represent overtones of the fundamental OH-stretching modes. The very pronounced pleochroism of the OH-stretching overtone in eveite (Ella>>Ellb>Ellc) is in agreement with the major alignment of the O-H vector along the  $a$ -direction determined by X-ray structural refinement of the isostructural mineral adamite (Hill, 1976).

In addition to the sharp bands representing fundamental frequencies and overtones of the OH-stretching modes there are two absorption features observed in the NIR and IR-spectra of eveite that suggest that structurally bound  $\text{H}_2\text{O}$ -molecules are present. The distinctly polarized absorption band at  $5200\text{ cm}^{-1}$  observed in the single crystal spectra and the two broad bands at  $3340$  and  $3450\text{ cm}^{-1}$  in the eveite powder spectrum display band features (energies and band widths) typical for  $\text{H}_2\text{O}$ -combination modes (bend+stretch) and OH-stretching modes, respectively, in water (Rossman, 1988b). This indication of additional water molecules in the structure may be one explanation for the low oxide sums obtained from the EMP-analyses of eveite.

## Conclusions

The colour and pleochroism of eveite is mainly determined by the presence of two distinctly anisotropic absorption bands at  $22\,160$  and  $16\,485\text{ cm}^{-1}$ , which are caused by spin-allowed  $d-d$  transitions in six-coordinated trivalent Mn. Remaining absorption bands in the UV-VIS spectral region in eveite are caused by spin-forbidden  $d-d$  transitions in  $\text{Mn}^{2+}$  at the five- and

six-coordinated sites. In the optical absorption spectra of sarkinite all bands in the same spectral range are assignable to divalent manganese at comparable five- and six-coordinated cation positions. The main cause for the distinct difference in colour between the two Mn<sub>2</sub>AsO<sub>4</sub>(OH)-polymorphs is the presence of small amounts of Mn<sup>3+</sup> in eveite.

The presence of structurally bound H<sub>2</sub>O in eveite has been indicated by NIR-spectra of single crystals as well as by IR-spectra of powdered samples. The low analytical sums recorded for the mineral by EMP-methods may be partially due to the presence of water molecules.

### Acknowledgements

We thank the Swedish Natural Science Research Council (NFR) for a grant to one of us (UH). Our thanks are also due to H. Harrysson, Uppsala University for microprobe analyses and J. Lindgren for providing access to the FTIR facility at Uppsala University. The manuscript was improved by a constructive review by Prof. G. Rossman, which we greatly appreciate.

### References

- Braithwaite, R.S.W. (1983) Infrared spectroscopic analysis of the olivenite-adamite series, and of phosphate substitution in olivenite. *Mineral. Mag.*, **47**, 51–57.
- Burns, R.G. (1993) *Mineralogical Applications of Crystal Field Theory*. Second edition. Cambridge University Press, Cambridge.
- Chisholm, J.E. (1985) Cation segregation and the O-H stretching vibration of the olivenite–adamite series. *Phys. Chem. Minerals*, **12**, 185–90.
- Dal Negro, A., Giuseppetti, G. and Pozas, J.M.M. (1974) The crystal structure of sarkinite, Mn<sub>2</sub>AsO<sub>4</sub>(OH). *Tscherm. Mineral. Petrol. Mitt.*, **21**, 246–60.
- Dowty, E. (1978) Absorption optics of low-symmetry crystals - applications to titanian clinopyroxene spectroscopy. *Phys. Chem. Minerals*, **3**, 173–81.
- Hålenius, U. and Lindqvist, B. (1996) Chromophoric divalent iron in optically anisotropic magnussonite. *Eur. J. Mineral.*, **8**, 25–34.
- Hill, R.J. (1976) The crystal structure and infrared properties of adamite. *Amer. Mineral.*, **61**, 979–86.
- Moore, P.B. (1968) Eveite, a new mineral from Långban. *Ark. Mineral. Geol.*, **4**, 473–6.
- Moore, P.B. and Smythe, J.R. (1968) Crystal chemistry of the basic arsenates: III. The crystal chemistry of eveite, Mn<sub>2</sub>(OH)(AsO<sub>4</sub>). *Amer. Mineral.*, **53**, 1841–6.
- Moore, R.K. and White, W.B. (1972) Electronic spectra of transition metal ions in silicate garnets. *Canad. Mineral.*, **11**, 791–811.
- Pouchou, J.L. and Pichoir, F. (1984) A new model for quantitative X-ray microanalysis. I. Application to the analysis of homogeneous samples. *La Recherche Aérospatiale*, **3**, 13–36.
- Rossmann, G.R. (1988a) Optical Spectroscopy. In *Spectroscopic Methods in Mineralogy and Geology* (F.C. Hawthorne, ed.). Mineral. Soc. Amer., *Rev. Mineral.*, **18**, 207–54.
- Rossmann, G.R. (1988b) Vibrational spectroscopy of hydrous components. In *Spectroscopic Methods in Mineralogy and Geology* (F.C. Hawthorne, ed.). Mineral. Soc. Amer., *Rev. Mineral.*, **18**, 193–206.
- Smith, G., Hålenius, U. and Langer, K. (1982) Low temperature spectral studies of Mn<sup>3+</sup>-bearing andalusite and epidote type minerals in the range 30,000–5,000 cm<sup>-1</sup>. *Phys. Chem. Minerals*, **8**, 136–42.

[Manuscript received 8 January 1997;  
revised 2 April 1997]